

**NIGHTTIME COLD AND FLU- acetaminophen, dextromethorphan hbr,
doxylamine succinate liquid
WALMART INC. (see also Equate)**

Drug Facts

Active ingredients (in each 30 mL)

Acetaminophen 650 mg

Dextromethorphan HBr 30 mg

Doxylamine succinate 12.5 mg

Purposes

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Uses

- temporarily relieves these common cold/flu symptoms
 - minor aches and pains
 - headache
 - sore throat
 - fever
 - runny nose and sneezing
 - cough due to minor throat and bronchial irritation

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks everyday while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

if a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is

accompanied or is followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist

Ask a doctor before use if you have

- liver disease
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
- trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking

- sedatives or tranquilizers
- the blood thinning drug warfarin

When using this product

- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- pain or cough gets worse or last more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts.

These could be a signs of a serious conditions.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed (see Overdose warning)
- do not take more than 4 doses in any 24-hour period
- measure only with dosing cup provided. Do not use any other dosing device.
- mL = milliliter
- keep dosing cup with product
- adults and children 12 years and over: 30 mL every 6 hours
- children under 12 years of age: do not use
- **when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing**

Other information

- **each 30 mL contains:** potassium 5 mg
- **each 30 mL contains:** sodium 24 mg
- store between 20-25°C (68-77°F). Do not refrigerate

Inactive ingredients

acesulfame potassium, alcohol, citric acid, FD&C blue1, FD&C red 40, Flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions or comments?

1-888-287-1915

Principal Display Panel

Compare to Vicks® NyQuil™ Cold & Flu active ingredients*

NIGHTTIME

Cold & Flu

Nighttime relief

- Acetaminophen 650 mg

Pain reliever/Fever reducer

- Dextromethorphan HBr 30 mg

Cough suppressant

- Doxylamine succinate 12.5 mg

Antihistamine

Relieves

- Headache, fever, sore throat, minor aches & pains
- Sneezing

- Runny nose
- Cough

For ages 12 years and over

Cherry flavor

10% ALCOHOL

FL OZ (mL)

*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® and NyQuil™ are registered trademarks of The Procter & Gamble Company.

TAMPER EVIDENT: DO NOT USE IF PRINTED SAFETY SEAL AROUND DOSAGE CUP OR UNDER CAP IS BROKEN OR MISSING.

DISTRIBUTED BY: Walmart Inc.,

Bentonville, AR 72716

Product Label

2 PACK

equate

NIGHTTIME
Cold & Flu
Nighttime relief

- Acetaminophen 650 mg
Pain reliever/Fever reducer
- Dextromethorphan HBr 30 mg
Cough suppressant
- Doxylamine succinate 12.5 mg
Antihistamine

Relieves

- Headache, fever, sore throat, minor aches & pains
- Sneezing
- Runny nose
- Cough

For ages 12 years and over

2 - 12 FL OZ (355mL) BOTTLES
24 FL OZ (1.5 PT) 710mL TOTAL

10% ALCOHOL

Drug Facts

Active Ingredients (in each 30 mL)
Acetaminophen 650 mg Pain reliever/fever reducer
Dextromethorphan HBr 30 mg Cough suppressant
Doxylamine succinate 12.5 mg Antihistamine

Purposes
■ temporarily relieves these common cold/flu symptoms
■ minor aches and pains ■ headache ■ sore throat
■ runny nose and sneezing ■ fever
■ cough due to minor throat and bronchial irritation

Uses
■ temporarily relieves these common cold/flu symptoms
■ minor aches and pains ■ headache ■ sore throat
■ runny nose and sneezing ■ fever
■ cough due to minor throat and bronchial irritation

Warnings
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take:
■ more than 4,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product
Severe allergic reactions: Acetaminophen may cause severe skin reactions. Symptoms may include:
■ skin redness ■ hives ■ rash ■ a skin reaction occurs, stop use and seek medical help right away.
See throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use
■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have
■ liver disease ■ diabetes
■ cough that occurs with too much phlegm (mucus)
■ a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema
■ trouble urinating due to enlarged prostate gland

Ask a doctor or pharmacist before use if you are
■ taking sedatives or tranquilizers
■ taking the blood thinning drug warfarin

Drug Facts (continued)

When using this product
■ excitability may occur, especially in children
■ marked drowsiness may occur
■ avoid alcoholic drinks
■ be careful when driving a motor vehicle or operating machinery
■ alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if
■ pain or cough gets worse or lasts more than 7 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present
■ new symptoms occur
■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions
■ do not take more than directed (see Overdose warning)
■ do not take more than 4 doses in any 24-hour period
■ measure only with dosing cup provided. Do not use any other dosing device.
■ mL = milliliter ■ keep dosing cup with product
■ adults and children 12 years and over: 30 mL every 6 hours
■ children under 12 years of age: do not use
■ when using other daytime or nighttime products, carefully read each label to ensure correct dosing

Other information
■ each 30 mL contains: potassium 5 mg
■ each 30 mL contains: sodium 24 mg
■ store between 20°-25°C (68°-77°F). Do not refrigerate.

Inactive ingredients acacia, aluminum potassium, alcohol, citric acid, FD&C Blue #1, FD&C Red #40, flavor, high fructose corn syrup, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium citrate

Questions or comments?
1-888-261-1915

EQUATE Nighttime Cold & Flu Cherry Flavor

NIGHTTIME COLD AND FLU

acetaminophen, dextromethorphan hbr, doxylamine succinate liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:79903-173
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
		650 mg

ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	30 mg in 30 mL
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL

Inactive Ingredients

Ingredient Name	Strength
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ALCOHOL (UNII: 3K9958V90M)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
HIGH FRUCTOSE CORN SYRUP (UNII: XY6UN3QB6S)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	CHERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:79903-173-08	237 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/30/2022	03/30/2027
2	NDC:79903-173-12	355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/30/2022	03/30/2027
3	NDC:79903-173-24	2 in 1 PACKAGE	12/30/2022	03/30/2027
3		355 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/30/2022	03/30/2027

Labeler - WALMART INC. (see also Equate) (051957769)

Revised: 12/2025

WALMART INC. (see also Equate)