

ACETAMINOPHEN- acetaminophen tablet
Akron Pharma Inc.

Acetaminophen 325 mg Tablets
Regular Strength

Active ingredient (in each tablet)

Acetaminophen 325 mg

Purpose

Pain Reliever/Fever Reducer

Uses

To reduce fever and for the temporary relief of minor aches and pains due to:

- headache
- muscular aches
- backache
- minor pain of arthritis
- the common cold
- toothache
- premenstrual and menstrual cramps.
- Temporarily reduces fever.

Warnings

Liver warning:

This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if the user has ever had an allergic reaction to this product or any of its ingredients.

Ask a doctor before use if the user has

- has liver disease
- is a child with pain of arthritis

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin

Stop use and ask a doctor if:

- pain gets worse or lasts more than 10 days (for adults) or 5 days (for children)
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose Warning:

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away.(1-800-222-1222)

Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions - do not take more than directed (see overdose warning)

adults & children 12 years and over	<ul style="list-style-type: none">• take 2 tablets every 4 to 6 hours while symptoms last• do not take more than 10 tablets in 24 hours• do not use for more than 10 days unless directed by a doctor
children 6-11 years	<ul style="list-style-type: none">• take 1 tablet every 4 to 6 hours while symptoms last• do not take more than 5 tablets in 24 hours• do not use for more than 5 days unless directed by a doctor
child under 6 years	ask a doctor

Other information

- store at 20° to 25°C (68° to 77°F)

Inactive Ingredients:

Gelatinized starch, magnesium stearate, povidone

Questions or Comments?

Call toll-free 1-877-225-6999

Manufactured for
Akron Pharma, Inc.,

373 RT US46 W Building E,

Suite 117, Fairfield, NJ - 07004

Mfg. Lic. No: TN/DRUGS/558/1997

* This product is not manufactured or distributed by Johnson and Johnson, consumer inc., distributor of regular Tylenol Tablets.

NDC- 71399-1025-6

Acetaminophen
325MG TABLETS

REGULAR STRENGTH

50 TABLETS

ASPIRIN FREE
PAIN RELIEVER
FEVER REDUCER

*Compare to the active ingredient
in Regular Strength Tylenol® Tablets

DO NOT USE WITH
OTHER MEDICINES
CONTAINING
ACETAMINOPHEN

Akron Pharma

Drug Facts	Purpose Pain reliever/ Fever reducer
Active ingredient (in each tablet) Acetaminophen 325 mg	
Uses	temporarily relieves minor aches and pains due to: <ul style="list-style-type: none"> ■ headache ■ backache ■ the common cold ■ premenstrual and menstrual cramps ■ temporarily reduces fever
Warnings	Liver warning: This product contains acetaminophen. Severe liver damage may occur if: <ul style="list-style-type: none"> ■ adult takes more than 4,000 mg of acetaminophen in 24 hours ■ child takes more than 5 doses in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: <ul style="list-style-type: none"> ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away.
Do not use	with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. If the user has ever had an allergic reaction to this product or any of its ingredients.
Ask a doctor before use if the user	has liver disease is a child with pain of arthritis
Ask a doctor or pharmacist before use if the user is	taking the blood thinning drug warfarin.

Child-Resistant Packaging

Drug Facts (Continued under label)

Do not use if imprinted safety seal under cap is broken or missing

*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol® Regular Strength

KOT-L214012-01

KTK/25/666/2017

Akron Pharma
Manufactured for:
Akron Pharma, Inc.
373 RT US46 W Building E,
Suite 117, Fairfield, NJ 07004
www.akronpharma.com

Rev: 08/24

No Varnish

PEEL HERE FOR MORE DRUG FACTS

Drug Facts (continued)

Stop use and ask a doctor if:

- pain gets worse or lasts more than 10 days (for adults) or 5 days (for children)
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present.

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (right away) (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions ■ do not take more than directed (see overdose warning)

adults & children 12 years and over	<ul style="list-style-type: none"> ■ take 2 tablets every 4 to 6 hours while symptoms last ■ do not take more than 10 tablets in 24 hours ■ do not use for more than 10 days unless directed by a doctor
children 6-11 years	<ul style="list-style-type: none"> ■ take 1 tablet every 4 to 6 hours while symptoms last ■ do not take more than 5 tablets in 24 hours ■ do not use for more than 5 days unless directed by a doctor
child under 6 years	ask a doctor

Other Information: store at 20° to 25° C (68° to 77° F)

Inactive ingredients: Gelatinized starch, magnesium stearate, povidone

Questions or comments? Call toll-free 1-877-225-6999

NDC 71399-1025-1

Acetaminophen 325MG TABLETS

REGULAR STRENGTH

100
TABLETS

ASPIRIN FREE
PAIN RELIEVER
FEVER REDUCER

*Compare to the active ingredient
in Regular Strength Tylenol® Tablets

DO NOT USE WITH
OTHER MEDICINES
CONTAINING
ACETAMINOPHEN



N 3 71399 10251 15

Drug Facts	Purpose Pain reliever/ Fever reducer
Active ingredient (in each tablet) Acetaminophen 325 mg	
Uses	temporarily relieves minor aches and pains due to: <ul style="list-style-type: none"> headache muscular aches backache minor pain of arthritis the common cold toothache premenstrual and menstrual cramps temporarily reduces fever
Warnings	Liver warning: This product contains acetaminophen. Severe liver damage may occur if: <ul style="list-style-type: none"> adult takes more than 4,000 mg of acetaminophen in 24 hours child takes more than 5 doses in 24 hours taken with other drugs containing acetaminophen adult has 3 or more alcoholic drinks every day while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: <ul style="list-style-type: none"> skin reddening blisters rash If a skin reaction occurs, stop use and seek medical help right away.
Do not use	with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. <ul style="list-style-type: none"> if the user has ever had an allergic reaction to this product or any of its ingredients.
Ask a doctor before use if the user	has liver disease is a child with pain of arthritis
Ask a doctor or pharmacist before use if the user is	taking the blood thinning drug warfarin

Drug Facts (continued under label)

Child-Resistant Packaging

Do not use if imprinted safety seal under cap is broken or missing

*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol® Regular Strength

KOT-L24002-01

KTK/25/666/2017

Akron Pharma
Manufactured for:
Akron Pharma, Inc.
373 RT US 46 W Building E,
Fairfield, NJ 07004
www.akronpharma.com

Rev. 08/24

No Varnish

PEEL HERE FOR MORE DRUG FACTS

Drug Facts (continued)
Stop use and ask a doctor if: <ul style="list-style-type: none"> pain gets worse or lasts more than 10 days (for adults) or 5 days (for children) fever gets worse or lasts more than 3 days new symptoms occur redness or swelling is present. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.
Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away, (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions do not take more than directed (see overdose warning)
adults & children 12 years and over <ul style="list-style-type: none"> take 2 tablets every 4 to 6 hours while symptoms last do not take more than 10 tablets in 24 hours do not use for more than 10 days unless directed by a doctor
children 6-11 years <ul style="list-style-type: none"> take 1 tablet every 4 to 6 hours while symptoms last do not take more than 5 tablets in 24 hours do not use for more than 5 days unless directed by a doctor
child under 6 years ask a doctor
Other information: store at 20° to 25° C (68° to 77° F)
Inactive ingredients: Gelatinized starch, magnesium stearate, povidone
Questions or comments? Call toll-free 1-877-225-6999

NDC 71399-1025-2

Acetaminophen 325MG TABLETS

REGULAR STRENGTH

1000
TABLETS

ASPIRIN FREE
PAIN RELIEVER
FEVER REDUCER

*Compare to the active ingredient
in Regular Strength Tylenol® Tablets

DO NOT USE WITH
OTHER MEDICINES
CONTAINING
ACETAMINOPHEN



N 3 71399 10252 12

Drug Facts	Purpose Pain reliever/ Fever reducer
Active ingredient (in each tablet) Acetaminophen 325 mg	
Uses	temporarily relieves minor aches and pains due to: <ul style="list-style-type: none"> headache muscular aches backache minor pain of arthritis the common cold toothache premenstrual and menstrual cramps temporarily reduces fever
Warnings	Liver warning: This product contains acetaminophen. Severe liver damage may occur if: <ul style="list-style-type: none"> adult takes more than 4,000 mg of acetaminophen in 24 hours child takes more than 5 doses in 24 hours taken with other drugs containing acetaminophen adult has 3 or more alcoholic drinks every day while using this product Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: <ul style="list-style-type: none"> skin reddening blisters rash If a skin reaction occurs, stop use and seek medical help right away.
Do not use	with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. <ul style="list-style-type: none"> if the user has ever had an allergic reaction to this product or any of its ingredients.
Ask a doctor before use if the user	has liver disease is a child with pain of arthritis
Ask a doctor or pharmacist before use if the user is	taking the blood thinning drug warfarin

Child-Resistant Packaging

Drug Facts (continued)
Stop use and ask a doctor if: <ul style="list-style-type: none"> pain gets worse or lasts more than 10 days (for adults) or 5 days (for children) fever gets worse or lasts more than 3 days new symptoms occur redness or swelling is present. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children.
Overdose warning: Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away, (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions do not take more than directed (see overdose warning)
adults & children 12 years and over <ul style="list-style-type: none"> take 2 tablets every 4 to 6 hours while symptoms last do not take more than 10 tablets in 24 hours do not use for more than 10 days unless directed by a doctor
children 6-11 years <ul style="list-style-type: none"> take 1 tablet every 4 to 6 hours while symptoms last do not take more than 5 tablets in 24 hours do not use for more than 5 days unless directed by a doctor
child under 6 years ask a doctor
Other information: store at 20° to 25° C (68° to 77° F)
Inactive ingredients: gelatinized starch, magnesium stearate, povidone
Questions or comments? Call toll-free 1-877-225-6999

Do not use if imprinted safety seal under cap is broken or missing

*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Tylenol® Regular Strength

KTK/25/666/2017

Akron Pharma
Manufactured for:
Akron Pharma, Inc.
373 RT US 46 W Building E,
Fairfield, NJ 07004
www.akronpharma.com

Rev. 08/24

No Varnish

PEEL HERE FOR MORE DRUG FACTS

LBH : 73 X 60 X 115

NDC 71399-1025-3

**Acetaminophen
325MG TABLETS**

REGULAR STRENGTH

**ASPIRIN FREE
PAIN RELIEVER
FEVER REDUCER**

*Compare to the active ingredient
in Regular Strength Tylenol® Tablets

**DO NOT USE WITH
OTHER MEDICINES
CONTAINING
ACETAMINOPHEN**



**100
TABLETS**

Unit Dose

**Acetaminophen 325 mg
REGULAR STRENGTH TABLETS**

Drug Facts

Active Ingredient (in each tablet) Purpose
Acetaminophen 325 mg Pain reliever/fever reducer

Uses Temporarily reduces fever and relieves minor aches and pains due to:
■ headache ■ muscular aches ■ backache ■ minor pain of arthritis
■ common cold ■ toothache ■ premenstrual and menstrual cramps

Warnings

Liver warning: This product contains acetaminophen. The maximum daily dose of this product is 10 tablets (3,250 mg) in 24 hours for adults or 5 tablets (1,625 mg) in 24 hours for children. Severe liver damage may occur if ■ adult takes more than 4,000 mg of acetaminophen in 24 hours ■ child takes more than 5 doses in 24 hours, which is the maximum daily amount ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product.

Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are allergic to acetaminophen or any of the inactive ingredients in this product.

Ask a doctor before use if you have ■ liver disease

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin.

Stop use and ask a doctor if ■ pain gets worse or lasts more than 10 days in adults; ■ pain gets worse or lasts more than 5 days in children under 12 years ■ fever gets worse or lasts more than 3 days ■ new symptoms occur ■ redness or swelling is present. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use. **Keep out of reach of children.**

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

This unit dose package is not child resistant and is intended for institutional use only.

READ AND KEEP CARTON FOR WARNINGS AND INFORMATION

**100
TABLETS**

REGULAR STRENGTH

**Acetaminophen
325MG
TABLETS**

NDC 71399-1025-3



NDC 71399-1025-3

**Acetaminophen
325MG TABLETS**

REGULAR STRENGTH

**ASPIRIN FREE
PAIN RELIEVER
FEVER REDUCER**

*Compare to the active ingredient
in Regular Strength Tylenol® Tablets

**100
TABLETS**

Unit Dose

**Acetaminophen 325 mg
REGULAR STRENGTH TABLETS**

Drug Facts (Continued)

Directions ■ do not take more than directed (see overdose warning)

adults & children 12 years and over ■ take 2 tablets every 4 to 6 hours while symptoms last
■ do not use for more than 10 tablets in 24 hours unless directed by a doctor

children 6 years to under 12 years ■ take 1 tablet every 4 to 6 hours while symptoms last
■ do not take more than 5 tablets in 24 hours
■ do not use for more than 3 days unless directed by a doctor

child under 6 years ask a doctor

Other information store at 20° to 25°C (68° to 77°F)

Inactive ingredients: Gelatinized starch, magnesium stearate, povidone

Questions or comments? Call toll-free 1-877-226-6999

Tamper Evident: Do not use if sealed blister units are broken or damaged.

*This product is not manufactured or distributed by McNeil Consumer Healthcare, distributor of Regular Strength Tylenol® Tablets

No Varnish

KT025/666/2017



Manufactured for:
Akron Pharma, Inc.
Fairfield, NJ 07004
www.akronpharma.com



ACETAMINOPHEN

acetaminophen tablet

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:71399-1025
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg

Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
POVIDONE (UNII: FZ989GH94E)	

Product Characteristics

Color	white	Score	2 pieces
Shape	ROUND	Size	4mm
Flavor		Imprint Code	A325

Contains**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:71399-1025-3	100 in 1 PACKAGE	12/06/2024	
1		1 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:71399-1025-6	50 in 1 BOTTLE; Type 0: Not a Combination Product	12/06/2024	
3	NDC:71399-1025-1	100 in 1 BOTTLE; Type 0: Not a Combination Product	12/06/2024	
4	NDC:71399-1025-2	1000 in 1 BOTTLE; Type 0: Not a Combination Product	12/06/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	12/06/2024	

Labeler - Akron Pharma Inc. (067878881)

Revised: 1/2025

Akron Pharma Inc.