MOTION SICKNESS RELIEF- dimenhydrinate 50 mg tablet TARGET CORPORATION

352R Target Dimenhydrinate 50 mg Tablets

Drug Facts

Active ingredient (in each tablet)

Dimenhydrinate 50 mg

Purpose

Antiemetic

Uses

for prevention and treatment of these symptoms associated with motion sickness:

- nausea
- vomiting
- dizziness

Warnings

Do not give to children under 2 years of age unless directed by a doctor.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

When using this product

- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

If pregnant or breast-feeding, ask a doctor before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- to prevent motion sickness, the first dose should be taken ½ to 1 hour before starting activity
- to prevent or treat motion sickness, see below:
 - take 1 to 2 tablets every 4-6 hours
 - do not take more than 8 tablets in 24 hours, or as

years and over

directed by a doctor

Children 6 to under 12 years

- give ½ to 1 tablet every 6-8 hours
- do not give more than 3 tablets in 24 hours, or as directed by a doctor
- give ½ tablet every 6-8 hours

Children 2 to under 6 years

• do not give more than 1 $\frac{1}{2}$ tablets in 24 hours, or as directed by a doctor

Other information

- SODIUM FREE
- store at room temperature 20-25°C (68-77°F)
- use by expiration date on package

Inactive ingredients

colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, mineral oil

Questions or comments?Call 1-800-910-6874

Coating/Print Omit Area

24 Tablets **Actual Size**

24 TABLETS

dnydn

gnitimov bns ssənizzib , səsu sn Dual action: Prevents and relieves

Direnhydrinate 50 mg Tablets/Antiemetic

Compare to active ingredient in Dramamine Original Formula*

RETAIN CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION TAMPER EVIDENT: DO NOT USE IF INDIVIDUAL BLISTER UNIT IS OPEN OR TORN 'This product is not manufactured or distributed by Medtech Products Inc., owner of the registered trademark Dramamine^{to} Original Formula.

Drug Facts

Active ingredient (in each tablet)

Purpose .Antiemetic ▶

■ for prevention and treatment of these symptoms associated

Dimenhydrinate 50 mg..

with motion sickness: ■ nausea ■ vomiting ■ dizziness

Warnings

TM & ©2025 Target Brands, Inc

NDC 11673-354

Coating/Print Omit Area

Do not give to children under 2 years of age unless directed by

Ask a doctor before use if you have

Drug Facts (continued)

■ a breathing problem such as emphysema or chronic bronchitis ■ glaucoma introuble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers.

When using this product

- marked drowsiness may occur avoid alcoholic drinks
- alcohol, sedatives, and franquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

If pregnant or breast-feeding, ask a doctor before use. Keep out of reach of children.

Drug Facts (continued)

Drug Facts (continued)

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

- to prevent motion sickness, the first dose should be taken ½ to 1 hour before starting activity
- to prevent or treat motion sickness, see below:

Adults and children 12 years and over

■ take 1 to 2 tablets every 4-6 hours ■ do not take more than 8 tablets in 24 hours, or as directed by a doctor

Children 6 to under 12 years

■ give ½ to 1 tablet every 6-8 hours ■ do not give more than 3 tablets in 24 hours, or as directed by a doctor

Children 2 to under 6 years

■ give ½ tablet every 6-8 hours ■ do not give more than 1 ½ tablets in 24 hours, or as directed by a doctor

-OI-034 -0000 C-002262-I 2 0726 ಕ 994

Drug Facts (continued)

Other information

- SODIUM FREE
- Store at room temperature 20-25℃ (68-77°F)
- use by expiration date on package

Drug Facts (continued)

Inactive ingredients colloidal silicon dioxide, croscarmellose sodium, hypromellose, lactose anhydrous, magnesium stearate, microcrystalline cellulose, mineral oil

Questions or comments? Call 1-800-910-6874

MOTION SICKNESS RELIEF

dimenhydrinate 50 mg tablet

Product Information

HUMAN OTC DRUG Product Type Item Code (Source) NDC:11673-354

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIMENHYDRINATE (UNII: JB937PER5C) (8-CHLOROTHEOPHYLLINE - UNII:GE2UA340FM)	DIMENHYDRINATE	50 mg

Inactive Ingredients			
Ingredient Name	Strength		
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)			
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)			
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
CROSCARMELLOSE SODIUM (UNII: M28OL1HH48)			
MAGNESIUM STEARATE (UNII: 70097M6I30)			
MINERAL OIL (UNII: T5L8T28FGP)			
ANHYDROUS LACTOSE (UNII: 3SY5LH9PMK)			

Product Characteristics					
Color	white	Score	2 pieces		
Shape	ROUND	Size	9mm		
Flavor		Imprint Code	TCL352		
Contains					

Packaging					
	#	Item Code	Package Description	Marketing Start Date	Marketing End Date
	1	NDC:11673-354- 24	4 in 1 CARTON	03/31/2025	
	1		6 in 1 BLISTER PACK; Type 0: Not a Combination Product		

Marketing Information				
Marketing Application Number or Monograp Category Citation		Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M009	03/31/2025		

Labeler - TARGET CORPORATION (006961700)

Registrant - TIME CAP LABORATORIES INC (037052099)

Establishment				
Name	Address	ID/FEI	Business Operations	
TIME CAP LABORATORIES, INC		037052099	manufacture(11673-354)	

Revised: 1/2025 TARGET CORPORATION