

RESFRIDOL ANTITUSSIVE- dextromethorphan hbr, guaifenesin syrup
Laboratorios Alfa

Resfridol Antitussive (NEW)

Drug Facts

Active ingredients and Purposes

Active ingredients (in each 10 mL)	Purposes
Dextromethorphan HBr 5 mg	Cough suppressant
Guaifenesin 200 mg.....	Expectorant

Uses

Temporarily relieves common cold symptoms:

- cough due to minor throat & bronchial irritation
- helps loosen phlegm (mucus) and thin bronchial secretions to:
- drain bronchial tubes
- make coughs more productive

Warnings

Do not use

- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Stop use and ask a doctor if

■ a cough persists for more than 7 days, tends to recur, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

Ask a doctor before use if you have

- a cough that is accompanied by excessive phlegm (mucus)
- a persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of accidental overdose, get medical help or contact a Poison Control Center right away.

Directions

- only use the dosage cup provided
- shake well before use
- do not exceed 6 doses per 24-hour period

Age

adults and children 12 years and older
children 6 to under 12 years
children under 6 years

Dose

20 mL, every 4 hours
10 mL, every 4 hours
do not use

Other information

- store between 68-77°F (20-25°C)
- do not use if safety seal over cap is torn, broken, or missing

Inactive ingredients

anhydrous citric acid, FD&C red #40, flavors, sodium benzoate, sorbitol, sucralose, water

Questions?

+1-809-544-0222

(M-F) 9 AM - 5 PM EST or www.laboratoriosalfa.com

Principal Display Panel

NDC 72483-101-04



Resfridol[®]
Antitussive
Syrup

Dextromethorphan HBr
(Cough suppressant)
Guaifenesin
(Expectorant)



4 fl oz (120 mL)

Drug Facts

Active ingredients (in each 10 mL)
Dextromethorphan HBr 5 mg.....Cough suppressant
Guaifenesin 200 mg.....Expectorant

Purposes

Uses Temporarily relieves common cold symptoms:
 ■ cough due to minor throat & bronchial irritation
 ■ helps loosen phlegm (mucus) and thin bronchial secretions to:
 ■ drain bronchial tubes ■ make coughs more productive

Warnings
Do not use ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
Ask a doctor before use if you have ■ a cough that is accompanied by excessive phlegm (mucus) ■ a persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Stop use and ask a doctor if ■ a cough persists for more than 7 days, tends to recur, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away.

Directions ■ only use the dosage cup provided ■ shake well before use ■ do not exceed 6 doses per 24-hour period

Age	Dose
adults and children 12 years and older	20 mL, every 4 hours
children 6 to under 12 years	10 mL, every 4 hours
children under 6 years	do not use

Other information ■ store between 77-86°F (25-30°C) ■ do not use if safety seal over cap is torn, broken, or missing

Inactive ingredients anhydrous citric acid, FD&C red #40, flavors, sodium benzoate, sorbitol, sucralose, water

NDC 72483-101-04



Resfridol[®]
Antitussivo
Jarabe

Dextrometorfano HBr
(Supresor de la tos)
Guaifenesina
(Expectorante)



4 oz fl (120 mL)

Datos Farmacológicos

Ingredientes activos (en cada 10 mL)
Dextrometorfano HBr 5 mg.....Supresor de la tos
Guaifenesina 200 mg.....Expectorante

Propósitos

Usos Alivia temporalmente los síntomas de un resfriado común:
 ■ tos debido a irritación leve de garganta y bronquios
 ■ ayuda a aliviar la flema (moco) y diluir las secreciones bronquiales para:
 ■ drenar los tubos bronquios ■ hacer que la tos sea más productiva

Advertencias
No use ■ si usted ahora está tomando un inhibidor de la monoamino oxidasa (MAO) recetado (ciertos medicamentos para la depresión, condiciones psiquiátricas o emocionales, o la enfermedad de Parkinson), o durante 2 semanas después de suspender el medicamento MAO. Si no sabe si su medicamento recetado contiene un MAO, consulte a un médico o farmacéutico antes de tomar este producto.
Pregúntele a un médico antes de usar si usted tiene ■ una tos que se acompaña de flema excesiva (moco) ■ una tos persistente o crónica como la que ocurre con el tabaquismo, el asma, bronquitis crónica, o el enfisema.
Pare el uso y pregúntele a un médico si ■ una tos persiste o los síntomas no mejoran en 7 días, tiende a reaparecer o se acompaña de fiebre, sarpullido o dolor de cabeza persistente. Estos podrían ser signos de una condición seria.
Si usted está embarazada o lactando, pregúntele a un profesional de la salud antes de usar.
Mantener fuera del alcance de los niños. En caso de sobredosis accidental, obtenga ayuda médica o llame a un Centro de Control de Intoxicaciones de inmediato.
Instrucciones ■ use únicamente la taza dosificadora proporcionada ■ agite bien antes de usar ■ no exceda 6 dosis en un periodo de 24 horas

Edad	Dosis
adultos y niños de 12 años y mayores	20 mL, cada 4 horas
niños de 6 a menos de 12 años	10 mL, cada 4 horas
niños menores de 6 años	no use

Otra información ■ almacenar entre 77-86°F (25-30°C) ■ no utilice si el sello de seguridad sobre la tapa está rasgado, roto o falta

Ingredientes inactivos ácido cítrico anhidro, FD&C rojo #40, sabores, benzoato de sodio, sorbitol, sucralosa, agua

RESFRIDOL ANTITUSSIVE

dextromethorphan hbr, guaifenesin syrup

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:72483-101
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	5 mg in 10 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
SUCRALOSE (UNII: 96K6UQ3ZD4)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	

Product Characteristics

Color	red	Score	
Shape		Size	
Flavor	STRAWBERRY	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:72483-101-04	1 in 1 BOX	01/31/2025	
1		120 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	01/31/2025	

Labeler - Laboratorios Alfa (815941244)

Revised: 12/2024

Laboratorios Alfa