

ASPIRIN PACKS- aspirin tablet, film coated
Sina Health Inc

Medifirst Aspirin

Drug Facts

Active ingredient (in each tablet)

Aspirin 325 mg (NSAID*)

*nonsteroidal anti-inflammatory drug

Purpose

Pain reliever/fever reducer

Uses

Temporarily relieves

- headache
- toothache
- pain and fever of colds
- muscle pain
- menstrual pain
- minor pain of arthritis

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning:

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcohol drinks every day while using this product

- take more or for a longer time than directed

Do not use if you are allergic to aspirin or to any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are taking a prescription drug for

- gout
- diabetes
- arthritis

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:

■ feel faint ■ vomit blood ■ have bloody or black stools ■ have stomach pain that does not get better

- pain gets worse or lasts more than 10 days
- fever gets worse or lasts more than 3 days
- if ringing in the ears or loss of hearing occurs
- redness or swelling is present
- new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- drink a full glass of water with each dose

Adults and children: Take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours.

(12 years and over)

Children under 12 years: Consult a doctor

Other information

- store at room temperature
- avoid excessive heat and humidity
- tamper evident sealed packets
- do not use any opened or torn packets

Inactive ingredients

corn starch, hypromellose*, microcrystalline cellulose*, polyethylene glycol*, povidone*, propylene glycol*

*m ay contain

Questions or comments? 1-800-634-76870

Medique Products Aspirin Label

Medique®

Aspirin

5 Grain (325 mg)

Pain Reliever/Fever Reducer

(NSAID)

This Package is for Households without Young Children.

Pull to Open

Aches/Fever • Aspirin 325 mg

200 Tablets (100 x 2)

Tamper Evident Unit Dose Packets



Medi-First Aspirin Label

100 Tablets

(50 x 2)

Medi-First®

Aspirin 5 Grain (325 mg)

Compare Active Ingredients to:

Genuine Bayer®

Registered Trademark of Pfizer Consumer

Pull To Open

Aches, Fever • Aspirin 325 mg

Pain Reliever/Fever Reducer (NSAID)

This Package is for Households without Young Children.

Tamper Evident Unit Dose Packets

MEDI-FIRST®
Aspirin 5 Grain (325 mg)
Compare active ingredient to:
Genuine Bayer®
Registered Trademark of Pfizer Consumer
✓ Aches, Fever • Aspirin 325 mg
Pain Reliever/Fever Reducer (NSAID)
THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.
Tamper Evident Unit Dose Packets

Drug Facts
Active ingredient (in each tablet) Purpose
Aspirin 325 mg (NSAID)¹..... Pain reliever/fever reducer
¹nonsteroidal anti-inflammatory drug

Uses
temporarily relieves
■ headache ■ muscle pain
■ toothache ■ menstrual pain
■ pain and fever of colds ■ minor pain of arthritis

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:
■ hives ■ facial swelling ■ asthma (wheezing) ■ shock
if an allergic reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)
Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcohol drinks every day while using this product
■ take more or for a longer time than directed

Do not use if you are allergic to aspirin or to any other pain reliever/fever reducer.

Ask a doctor before use if
■ stomach bleeding warning applies to you
■ you have a history of stomach problems such as heartburn
■ you have high blood pressure, heart disease, liver disease, or kidney disease
■ you are taking a diuretic
■ you have asthma
(Continued on opposite panel)

MEDI-FIRST®
Aspirin 5 Grain (325 mg)
Compare active ingredient to:
Genuine Bayer®
Registered Trademark of Pfizer Consumer
Pull To Open
✓ Aches, Fever • Aspirin 325 mg
Pain Reliever/Fever Reducer (NSAID)
THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.
Tamper Evident Unit Dose Packets

Drug Facts (continued)
Ask a doctor or pharmacist before use if you are taking a prescription drug for:
■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if
■ you experience any of the following signs of stomach bleeding:
■ feel faint ■ vomit blood
■ have bloody or black stools
■ have stomach pain that does not get better
■ pain gets worse or lasts more than 10 days
■ fever lasts more than 3 days
■ if ringing in the ears or loss of hearing occurs
■ redness or swelling is present
■ new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Drug Facts (continued)
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions
■ drink a full glass of water with each dose
Adults and children (12 years and over) Take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours, not to exceed 12 tablets in 24 hours.
Consult a doctor.
Children under 12 years: Consult a doctor.

Other information
■ store at room temperature
■ avoid excessive heat and humidity
■ tamper evident sealed packets
■ do not use any opened or torn packets

Inactive ingredients
corn starch, hypromellose¹, microcrystalline cellulose², polyethylene glycol³, povidone⁴, polyethylene glycol⁵
¹may contain

Questions or comments? 1-800-634-7680

9053300L
Rev. 5/6/22

MEDI-FIRST®
Aspirin 5 Grain (325 mg)
Pain Reliever/Fever Reducer (NSAID)

4768280533

Reorder #05533
Manufactured for
Medique Products
Fort Myers, FL 33907
1-800-634-7680
www.mediqproducts.com

RETAIN CARTON FOR COMPLETE PRODUCT INFORMATION

Medi-First Plus Aspirin Label

250 Tablets

(125 x 2's)

Medi First® Plus

Aspirin

This Package is for Households without Young Children.

Pull To Open

Aspirin 325 mg (NSAID)

Pain Reliever/Fever Reducer

Compare active ingredient to:

Genuine Bayer®

Registered Trademark of Bayer Corporation
 Tamper Evident Unit Dose Packets



Medique®
 Aspirin
 5 Grain (325 mg)
 Pain Reliever/Fever Reducer
 (NSAID)
 This Package is for Households without Young Children.
 Pull to Open
 Aches/Fever • Aspirin 325 mg
 200 Tablets (100 x 2)
 Tamper Evident Unit Dose Packets



100 Tablets

(50 x 2)

Medi-First®

Aspirin 5 Grain (325 mg)

Compare Active Ingredients to:

Genuine Bayer®

Registered Trademark of Pfizer Consumer


Pull To Open

Aches, Fever • Aspirin 325 mg

Pain Reliever/Fever Reducer (NSAID)

This Package is for Households without Young Children.

Tamper Evident Unit Dose Packets



250 Tablets

(125 x 2's)

Medi First® Plus

Aspirin

This Package is for Households without Young Children.

Pull To Open

Aspirin 325 mg (NSAID)

Pain Reliever/Fever Reducer

Compare active ingredient to:

Genuine Bayer®

Registered Trademark of Bayer Corporation

Tamper Evident Unit Dose Packets



90548001
Rev. 5/16/22
Pkg. 14,24

Drug Facts

Active ingredient (in each tablet) Purpose
Aspirin 325 mg (NSAID) Pain reliever/fever reducer
nonsteroidal anti-inflammatory drug

Uses
Temporarily relieve
■ headache ■ muscle pain
■ minor pain of arthritis ■ pain and fever of colds
■ toothache ■ menstrual pain

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Warnings (continued)
Allergic alert: Aspirin may cause a severe allergic reaction which may include:
■ hives ■ facial swelling
■ asthma (wheezing) ■ shock
If an allergic reaction occurs, stop use and seek medical help right away.

Drug Facts (continued)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:
■ are age 65 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
■ have 3 or more alcohol drinks every day while using this product
■ take more for a longer time than directed

Do not use if you are allergic to aspirin or to any other pain reliever/fever reducer.

Ask a doctor before use if
■ stomach bleeding warning applies to you
■ you have a history of stomach problems such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
■ you are taking a diuretic
■ you have asthma

(continued on opposite panel)



250 Tablets (125 x 2's)
Aspirin 325 mg (NSAID)
Pain Reliever/Fever Reducer

THIS PACKAGE IS FOR HOUSEHOLDS WITHOUT YOUNG CHILDREN.
Pull To Open
Compare active ingredient to:
Genuine Bayer® Tamper Evident Unit Dose Packets
Registered Trademark of Bayer Corporation

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking a prescription drug for: ■ gout ■ diabetes ■ arthritis

Stop use and ask a doctor if
■ an allergic reaction occurs. Seek medical help right away.
■ you experience any of the following signs of stomach bleeding:
■ dark, tarry ■ vomit blood
■ new bloody or black stools
■ new stomach pain that does not get better

■ pain gets worse or lasts more than 10 days
■ fever lasts more than 3 days
■ ringing in the ears or loss of hearing occurs
■ redness or swelling is present
■ new symptoms appear

If pregnant or breast feeding, ask a health professional before use. It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Drug Facts (continued)

Directions
■ drink a full glass of water with each dose

Adults and children 10 years and over: Take 1 or 2 tablets every 4 hours or 3 tablets every 6 hours. Not to exceed 12 tablets in 24 hours.
Children under 12 years: Consult a doctor.

Other information
■ store at room temperature
■ avoid excessive heat and humidity
■ tamper evident sealed packets
■ do not use any opened or torn packets

Inactive ingredients
corn starch, hypromellose, microcrystalline cellulose, polyethylene glycol, potassium, propylene glycol
*may contain:
Questions or comments? 1-800-634-7680



Recorder #90548
Manufactured for Medi First Plus
Fort Myers, FL 33967
1-800-634-7680
www.medicapproducts.com

ASPIRIN PACKS			
aspirin tablet, film coated			
Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70385-2006(NDC:47682-617)
Route of Administration	ORAL		
Active Ingredient/Active Moiety			
Ingredient Name		Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)		ASPIRIN	325 mg
Inactive Ingredients			
Ingredient Name			Strength
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)			
STARCH, CORN (UNII: O8232NY3SJ)			
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)			
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)			
Product Characteristics			
Color	white	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	44;157;ASPIRIN
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70385-2006-2	2 in 1 PACKAGE	07/20/2020	
1		2 in 1 PACKET; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	07/20/2020	

Labeler - Sina Health Inc (047161553)

Establishment

Name	Address	ID/FEI	Business Operations
Sina Health Inc		047161553	repack(70385-2006)

Revised: 1/2026

Sina Health Inc