

**NIGHT TIME COLD AND FLU RELIEF MULTI SYMPTOM- acetaminophen,  
dextromethorphan hbr, doxylamine succinate capsule, liquid filled  
Rite Aid Corporation**

-----  
**Drug Facts**

**Active ingredients (in each softgel)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

**Purpose**

**Pain reliever/Fever Reducer**

Cough suppressant

Antihistamine

**Uses**

- temporarily relieves common cold and flu symptoms:
  - sore throat
  - headache
  - minor aches and pains
  - fever
  - runny nose and sneezing
  - cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks everyday while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists more than 2 days, is

accompanied or followed by fever, headache rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

### **Ask a doctor before use if you have**

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

### **Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

### **When using this product**

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

### **Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as

for children even if you do not notice any signs or symptoms.

### **Directions**

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- **when using other Daytime or Nighttime products, carefully read each label to insure correct dosing**

### **Other information**

- store between 15-30°C (59-86°F)
- avoid excessive heat

### **Inactive ingredients**

D&C yellow #10, FD&C blue #1, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

### **Questions or comments?**

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

### **Principal Display Panel**

Compare to the active ingredients in Vicks® Nyquil® Cold & Flu LiquiCaps®\*

MULTI-SYMP TOM

COLD & FLU RELIEF

NIGHTTIME

ACETAMINOPHEN 325 mg

DEXTROMETHORPHAN HBr 15 mg

DOXYLAMINE SUCCINATE 6.25 mg

PAIN RELIEVER / FEVER REDUCER

COUGH SUPPRESSANT • ANTIHISTAMINE

Reliever aches, fever, sore throat, cough, runny nose & sneezing

Alcohol-free

SOFTGELS\*\*

\*\*LIQUID FILLED CAPSULES

\*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks® , NyQuil® , and LiquiCaps® are registered trademark of The Procter & Gamble Company.

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

**DISTRIBUTED BY:**

RITE AID

30 HUNTER LANE

CAMP HILL, PA 17011

[www.riteaid.com](http://www.riteaid.com)

**Package Label**

**Questions or comments?**  
Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

**Inactive ingredients**  
D&C yellow #10, FD&C blue #1, gelatin, glycerin, mannitol, polyethylene glycol, povidone, purified water, sorbitan, sorbitol, white ink

**Other information**  
■ store between 15-30°C (59-86°F) ■ avoid excessive heat

**Directions**  
■ do not take more than directed (see Overdose warning)  
■ adults and children 12 years and over: take 2 softgels with water every 6 hours  
■ swallow whole; do not crush, chew, or dissolve  
■ children under 12 years: do not use  
■ when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing

**Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Keep out of reach of children.**  
**Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Do not use**  
■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
■ liver disease ■ glaucoma  
■ trouble urinating due to an enlarged prostate gland  
■ a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis or emphysema  
■ cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are**  
■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers  
■ do not exceed recommended dosage

■ be careful when driving a motor vehicle or operating machinery  
■ alcohol, sedatives, and tranquilizers may increase drowsiness  
■ avoid alcoholic drinks ■ marked drowsiness may occur  
■ excitability may occur, especially in children

**Drug Facts (continued)**

**Warnings**  
Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions.  
Symptoms may include: ■ skin reddening ■ blisters ■ rash  
If a skin reaction occurs, stop use and seek medical help right away.  
**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**  
■ with any other drug containing acetaminophen (prescription or nonprescription), if you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.  
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug, if you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
■ liver disease ■ glaucoma  
■ trouble urinating due to an enlarged prostate gland  
■ a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis or emphysema  
■ cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are**  
■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers  
■ do not exceed recommended dosage

■ be careful when driving a motor vehicle or operating machinery  
■ alcohol, sedatives, and tranquilizers may increase drowsiness  
■ avoid alcoholic drinks ■ marked drowsiness may occur  
■ excitability may occur, especially in children

**Drug Facts (continued)**

**Uses**  
■ temporarily relieves common cold and flu symptoms: ■ minor aches and pains ■ sore throat ■ headache ■ runny nose and sneezing ■ fever ■ cough due to minor throat and bronchial irritation

**Drug Facts**

**Active ingredients (in each softgel)**  
Acetaminophen 325 mg.....Pain reliever/fever reducer  
Dextromethorphan HBr 15 mg.....Cough suppressant  
Doxylamine succinate 6.25 mg.....Antihistamine

**Purposes**

NDC 11822-3335-5 Compare to the active ingredients in Vicks® NyQuil® Cold & Flu LiquiCaps®\*

# MULTI-SYMPOTOM COLD & FLU RELIEF

**NIGHTTIME**

**ACETAMINOPHEN 325 mg**  
**DEXTROMETHORPHAN HBr 15 mg**  
**DOXYLAMINE SUCCINATE 6.25 mg**

PAIN RELIEVER / FEVER REDUCER  
COUGH SUPPRESSANT • ANTIHISTAMINE

Relieves aches, fever, sore throat,  
cough, runny nose & sneezing

Alcohol-free

**24**

**RITE AID**



TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.  
KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

PHARMACY

SOFTGELS

\*\*LIQUID FILLED CAPSULES

Lot No  
Exp Dt

\*This product is not manufactured or distributed by The Procter & Gamble Company.  
Vicks®, NyQuil®, and LiquiCaps® are registered trademarks of The Procter & Gamble Company.

PLD-E569B FC005808



**DISTRIBUTED BY:**  
RITE AID  
30 HUNTER LANE  
CAMP HILL, PA 17011  
www.riteaid.com

**SATISFACTION  
GUARANTEE**  
If you're not satisfied, we'll  
happily refund your money.

**PARENTS:**  
Learn about teen medicine abuse  
www.StopMedicineAbuse.org

**RITE AID PHARMACY Multi-Symptom Cold & Flu Relief**

**NIGHT TIME COLD AND FLU RELIEF MULTI SYMPTOM**

acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:11822-3335
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RT19KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	15 mg
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg

**Inactive Ingredients**

Ingredient Name	Strength
<b>D&amp;C YELLOW NO. 10</b> (UNII: 35SW5USQ3G)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>GELATIN</b> (UNII: 2G86QN327L)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ05DW1A)	
<b>POVIDONE</b> (UNII: FZ989GH94E)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SORBITAN</b> (UNII: 6O92ICV9RU)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>MANNITOL</b> (UNII: 3OWL53L36A)	

**Product Characteristics**

<b>Color</b>	green	<b>Score</b>	no score
<b>Shape</b>	OVAL	<b>Size</b>	20mm
<b>Flavor</b>		<b>Imprint Code</b>	P30
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:11822-3335-5	24 in 1 CARTON	05/29/2020	12/29/2026
1		1 in 1 BLISTER PACK; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	05/29/2020	12/29/2026

**Labeler** - Rite Aid Corporation (014578892)

Revised: 8/2025

Rite Aid Corporation