**NIGHT TIME COLD AND FLU RELIEF MULTI SYMPTOM- acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled**
**Rite Aid Corporation**

---

**Drug Facts**

**Active ingredients (in each softgel)**

**Acetaminophen 325 mg**

Dextromethorphan HBr 15 mg

Doxylamine succinate 6.25 mg

**Purpose**

**Pain reliever/Fever Reducer**

Cough suppressant

Antihistamine

**Uses**

- temporarily relieves common cold and flu symptoms:
  - sore throat
  - headache
  - minor aches and pains
  - fever
  - runny nose and sneezing
  - cough due to minor throat and bronchial irritation

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take:

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks everyday while using this product

**Allergy alert**: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning**: If sore throat is severe, persists more than 2 days, is

accompanied or followed by fever, headache rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor of pharmacist before taking this product.

**Ask a doctor before use if you have**

- liver disease
- glaucoma
- trouble urinating due to an enlarged prostate gland
- a breathing problem or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are**

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

**When using this product**

- **do not exceed recommended dosage**
- excitability may occur, especially in children
- avoid alcoholic drinks
- marked drowsiness may occur
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- pain or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as

for children even if you do not notice any signs or symptoms.

**Directions**

- **do not take more than directed (see Overdose warning)**
- do not take more than 4 doses in 24 hours
- adults and children 12 years and over: take 2 softgels with water every 6 hours
- swallow whole; do not crush, chew, or dissolve
- children under 12 years: do not use
- **when using other Daytime or Nighttime products, carefully read each label to insure correct dosing**

**Other information**

- store between 15-30ºC (59-86ºF)
- avoid excessive heat

**Inactive ingredients**

D&C yellow #10, FD&C blue #1, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

**Questions or comments?**

Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

**Principal Display Panel**

Compare to the active ingredients in Vicks® Nyquil® Cold & Flu LiquiCaps®*

MULTI-SYMPTOM

COLD & FLU RELIEF

NIGHTTIME

ACETAMINOPHEN 325 mg

DEXTROMETHORPHAN HBr 15 mg

DOXYLAMINE SUCCINATE 6.25 mg

PAIN RELIEVER / FEVER REDUCER

COUGH SUPPRESSANT • ANTIHISTAMINE

Reliever aches, fever, sore throat, cough, runny nose & sneezing

Alcohol-free

SOFTGELS**

**LIQUID FILLED CAPSULES

*This product is not manufactured or distributed by The Procter & Gamble Company. Vicks®, NyQuil®, and LiquiCaps® are registered trademark of The Procter & Gamble Company.

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**

**DISTRIBUTED BY:**

RITE AID

30 HUNTER LANE

CAMP HILL, PA 17011

www.riteaid.com

**Package Label**

NDC 11822-3335-5

Compare to the active ingredients in **Vicks® NyQuil® Cold & Flu LiquiCaps®***

# MULTI-SYMPTOM COLD & FLU RELIEF

## NIGHTTIME

**ACETAMINOPHEN 325 mg**
**DEXTROMETHORPHAN HBr 15 mg**
**DOXYLAMINE SUCCINATE 6.25 mg**

PAIN RELIEVER / FEVER REDUCER
COUGH SUPPRESSANT • ANTIHISTAMINE

**Relieves aches, fever, sore throat, cough, runny nose & sneezing**

Alcohol-free

ACTUAL SIZE

**24**

RITE AID

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

## Drug Facts

| Active ingredients (in each softgel) | Purposes |
|---|---|
| Acetaminophen 325 mg | Pain reliever/fever reducer |
| Dextromethorphan HBr 15 mg | Cough suppressant |
| Doxylamine succinate 6.25 mg | Antihistamine |

## Drug Facts (continued)

### Uses

- temporarily relieves common cold and flu symptoms:
  - sore throat ■ headache ■ minor aches and pains
  - fever ■ runny nose and sneezing
  - cough due to minor throat and bronchial irritation

## Drug Facts (continued)

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take: ■ more than 4,000 mg of acetaminophen in 24 hours
■ with other drugs containing acetaminophen
■ 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions.
Symptoms may include: ■ skin reddening ■ blisters ■ rash
If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### Do not use

■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

**Ask a doctor before use if you have** ■ liver disease ■ glaucoma
■ trouble urinating due to an enlarged prostate gland
■ a breathing problem or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis or emphysema
■ cough that occurs with too much phlegm (mucus)

**Ask a doctor or pharmacist before use if you are**
■ taking the blood thinning drug warfarin ■ taking sedatives or tranquilizers

**When using this product** ■ **do not exceed recommended dosage**
■ excitability may occur, especially in children
■ avoid alcoholic drinks ■ marked drowsiness may occur
■ alcohol, sedatives, and tranquilizers may increase drowsiness
■ be careful when driving a motor vehicle or operating machinery

## Drug Facts (continued)

### Stop use and ask a doctor if
■ pain or cough gets worse or lasts more than 7 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present ■ new symptoms occur
■ cough comes back or occurs with rash or headache that lasts.
These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** Taking more than the recommended dose can cause liver damage. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### Directions
■ **do not take more than directed (see Overdose warning)**
■ do not take more than 4 doses in 24 hours
■ adults and children 12 years and over: take 2 softgels with water every 6 hours
■ swallow whole; do not crush, chew, or dissolve
■ children under 12 years: do not use
■ **when using other Daytime or Nighttime products, carefully read each label to ensure correct dosing**

### Other information
■ store between 15-30°C (59-86°F) ■ avoid excessive heat

### Inactive ingredients
D&C yellow #10, FD&C blue #1, gelatin, glycerin, mannitol, polyethylene glycol, povidone, propylene glycol, purified water, sorbitan, sorbitol, white ink

### Questions or comments?
Call **1-877-753-3935** Monday-Friday 9AM-5PM EST

**RITE AID PHARMACY Multi-Symptom Cold & Flu Relief**

# NIGHT TIME COLD AND FLU RELIEF MULTI SYMPTOM

acetaminophen, dextromethorphan hbr, doxylamine succinate capsule, liquid filled

## Product Information

| | | | |
|---|---|---|---|
| **Product Type** | HUMAN OTC DRUG | **Item Code (Source)** | NDC:11822-3335 |
| **Route of Administration** | ORAL | | |

## Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|---|---|
| **ACETAMINOPHEN** (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D) | ACETAMINOPHEN | 325 mg |
| **DEXTROMETHORPHAN HYDROBROMIDE** (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS) | DEXTROMETHORPHAN HYDROBROMIDE | 15 mg |
| **DOXYLAMINE SUCCINATE** (UNII: V9BI9B5YI2) (DOXYLAMINE - UNII:95QB77JKPL) | DOXYLAMINE SUCCINATE | 6.25 mg |

## Inactive Ingredients

| Ingredient Name | Strength |
|---|---|
| **D&C YELLOW NO. 10** (UNII: 35SW5USQ3G) | |
| **FD&C BLUE NO. 1** (UNII: H3R47K3TBD) | |
| **GELATIN** (UNII: 2G86QN327L) | |
| **GLYCERIN** (UNII: PDC6A3C0OX) | |
| **POLYETHYLENE GLYCOL, UNSPECIFIED** (UNII: 3WJQ0SDW1A) | |
| **POVIDONE** (UNII: FZ989GH94E) | |
| **PROPYLENE GLYCOL** (UNII: 6DC9Q167V3) | |
| **WATER** (UNII: 059QF0KO0R) | |
| **SORBITAN** (UNII: 6O92ICV9RU) | |
| **SORBITOL** (UNII: 506T60A25R) | |
| **MANNITOL** (UNII: 3OWL53L36A) | |

## Product Characteristics

| | | | |
|---|---|---|---|
| **Color** | green | **Score** | no score |
| **Shape** | OVAL | **Size** | 20mm |
| **Flavor** | | **Imprint Code** | P30 |
| **Contains** | | | |

## Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|---|---|---|---|
| **1** | NDC:11822-3335-5 | 24 in 1 CARTON | 05/29/2020 | |
| **1** | | 1 in 1 BLISTER PACK; Type 0: Not a Combination Product | | |

## Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---|---|---|---|
| OTC Monograph Drug | M012 | 05/29/2020 | |

## Labeler - Rite Aid Corporation (014578892)

Revised: 5/2024

Rite Aid Corporation