

**WOMENS GENTLE LAXATIVE- bisacodyl tablet, coated  
Preferred Plus (Kinray)**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Drug Facts**

**Active ingredient (in each tablet)**

Bisacodyl USP 5 mg

**Purpose**

Stimulant laxative

**Use**

- for temporary relief of occasional constipation and irregularity
- this product generally produces a bowel movement in 6 to 12 hours

**Warnings**

**Do not use**

if you cannot swallow without chewing.

**Ask a doctor before use if you have**

- stomach pain, nausea or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

**When using this product**

- do not chew or crush tablet(s)
- do not use within 1 hour after taking an antacid or milk
- it may cause stomach discomfort, faintness and cramps

**Stop use and ask a doctor if**

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions**

- take with a glass of water

---

adults and children 12 years of age and over | take 1 to 3 tablets in a single daily dose

children 6 to under 12 years of age	take 1 tablet in a single daily dose
children under 6 years of age	ask a doctor

### Other information

- **each tablet contains:** calcium 15 mg
- store between 20°-25°C (68°-77°F)

### Inactive ingredients

acacia, calcium sulfate anhydrous, carnauba wax, colloidal silicon dioxide, corn starch, D&C red #7 calcium lake, hypromellose phthalate, iron oxide, iron oxide black, iron oxide yellow, kaolin, magnesium stearate, microcrystalline cellulose, polyethylene glycol (PEG) 400, povidone, shellac, sodium starch glycolate, stearic acid, sugar, talc, titanium dioxide, and triacetin

### Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

### Principal Display Panel

\*Compare to the Active Ingredient in Correctol®

### Women's *Gentle Laxative*

*Bisacodyl Stimulant Laxative*

For Dependable, Natural Overnight Relief

Tablets

Each Enteric Coated Tablet Contains 5 mg Bisacodyl

**TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.**

**KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.**


\*This product is not manufactured or distributed by MSD Consumer Care Inc., distributor of Correctol®.

**Distributed by: Kinray, LLC.**

**152-35 Tenth Avenue**

**Whitestone, NY 11357**

### Package Label



<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>■ a sudden change in bowel habits that lasts more than 2 weeks</li> <li>■ When using this product <ul style="list-style-type: none"> <li>■ do not chew or crush tablet(s)</li> <li>■ do not use within 1 hour after taking an antacid or milk</li> <li>■ it may cause stomach discomfort, faintness and cramps</li> </ul> </li> <li>■ Stop use and ask a doctor if <ul style="list-style-type: none"> <li>■ you have rectal bleeding or no bowel movement after</li> </ul> </li> </ul>	
<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>■ adults and children 12 years of age and over take 1 to 3 tablets in a single daily dose</li> <li>■ children 6 to under 12 take 1 tablet in a single daily dose</li> <li>■ children under 6 years of age ask a doctor</li> </ul>	
<p><b>Other information</b></p> <ul style="list-style-type: none"> <li>■ each tablet contains calcium 15 mg</li> <li>■ store between 20°C-25°C (68°F-77°F)</li> </ul>	

<p><b>Drug Facts</b></p> <p><b>Active ingredient (in each tablet)</b> Bisacodyl USP 5 mg.....Stimulant laxative</p> <p><b>Purpose</b> ■ for temporary relief of occasional constipation and irregularity</p>	
<p><b>Warnings</b></p> <p>Do not use if you cannot swallow without chewing.</p> <p>Ask a doctor before use if you have stomach pain, nausea or vomiting.</p>	
<p><b>Drug Facts (continued)</b></p> <ul style="list-style-type: none"> <li>■ this product generally produces a bowel movement in 6 to 12 hours</li> </ul>	

NDC 61715-039-30



**\*Compare to the Active Ingredient in Correctol®**

# Women's Gentle Laxative

*Bisacodyl Stimulant Laxative*

For Dependable, Natural Overnight Relief

**30 Tablets** 

Each Enteric Coated Tablet Contains 5 mg Bisacodyl



TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF BLISTER UNIT IS TORN, BROKEN OR SHOWS ANY SIGNS OF TAMPERING.

KEEP OUTER CARTON FOR COMPLETE WARNINGS AND PRODUCT INFORMATION.

Lot No.:  
Exp. Date:

Distributed by: Kinray, LLC.  
152-35 Tenth Avenue  
Whitestone, NY 11357.

©2014 Kinray LLC., All Rights Reserved, PREFERRED PLUS, PREFERRED PLUS++++ and the PREFERRED PLUS logo are trademarks and/or registered trademarks of Kinray LLC. All other marks are property of their respective owners.  
All Preferred Plus Pharmacy products are satisfaction guaranteed or your money back from retailer where purchased.  
PLD-C14M FC002117

\*This product is not manufactured or distributed by MSD Consumer Care Inc., distributor of Correctol®.  
KIN# 410-159



Preferred Plus Women's Gentle Laxative

# WOMENS GENTLE LAXATIVE

bisacodyl tablet, coated

## Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:61715-039
Route of Administration	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BISACODYL (UNII: 10X0709Y6I) (DEACETYLBISACODYL - UNII:R09078E41Y)	BISACODYL	5 mg

## Inactive Ingredients

Ingredient Name	Strength
ACACIA (UNII: 5C5403N26O)	
CALCIUM SULFATE ANHYDROUS (UNII: E934B3V59H)	
CARNAUBA WAX (UNII: R12CBM0EIZ)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
STARCH, CORN (UNII: O8232NY3SJ)	
D&C RED NO. 7 (UNII: ECW0LZ41X8)	
HYPROMELLOSE PHTHALATE (24 % PHTHALATE, 55 CST) (UNII: 87Y6436BKR)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERROSFERRIC OXIDE (UNII: XM0M87F357)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
KAOLIN (UNII: 24H4NWX5CO)	
MAGNESIUM STEARATE (UNII: 70097M6I30)	
CELLULOSE, MICROCRYSTALLINE (UNII: OP1R32D61U)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
POVIDONE (UNII: FZ989GH94E)	
SHELLAC (UNII: 46N107B71O)	
SODIUM STARCH GLYCOLATE TYPE A CORN (UNII: AG9B65PV6B)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
SUCROSE (UNII: C151H8M554)	
TALC (UNII: 7SEV7J4R1U)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TRIACETIN (UNII: XHX3C3X673)	

## Product Characteristics

Color	PINK	Score	no score
Shape	ROUND	Size	11mm
Flavor		Imprint Code	TCL057
Contains			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	----------------------	--------------------

1	NDC:61715-039-30	2 in 1 CARTON	08/02/2010	
1		15 in 1 BLISTER PACK; Type 0: Not a Combination Product		

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC MONOGRAPH NOT FINAL	part334	08/02/2010	

**Labeler** - Preferred Plus (Kinray) (012574513)

Revised: 11/2019

Preferred Plus (Kinray)