

**ASPIRIN 325 MG- aspirin tablet**  
**Reliable 1 Laboratories LLC**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**ASPIRIN USP 325 MG TABLET**

***Drug Facts***

Aspirin USP, 325 mg (NSAID)\*

\*nonsteroidal anti-inflammatory drug

Pain Reliever/Fever Reducer

temporarily relieves

- headache
- menstrual pain
- minor pain of arthritis
- muscle pain
- toothache
- pain and fever of colds

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include

- hives
- facial swelling
- asthma (wheezing)
- shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (ibuprofen, naproxen, or others)
- take more or for a longer time than directed
- have 3 or more alcoholic drinks every day while using this product

***Drug Facts*** continued on back of label

**Do not use** if you have ever had an allergic reaction to any other pain reliever/fever reducer

**Ask a doctor before use if**

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

- you have asthma

### **Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for diabetes, gout or arthritis
- under a doctor's care for any serious condition
- taking any other drug

### **Stop use and ask a doctor if**

- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
  - pain gets worse or lasts more than 10 days
  - fever gets worse or lasts more than 3 days
  - redness or swelling is present in the painful area
  - any new symptoms appear
  - ringing in the ears or loss of hearing occurs

These could be signs of a serious condition

**If pregnant or breast-feeding**, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control right away.

### **Directions**

DRINK A FULL GLASS OF WATER WITH EACH DOSE

- **adults and children 12 years and over:** 1-2 tablets every 4 hours while symptoms last, not more than 12 tablets in 24 hours
- **children under 12 years:** ask a doctor

### **Other information**

- store at room temperature 15°-30°C (59°-86°F)

**Inactive ingredients** corn starch, povidone

### **Questions or comments**

**Questions or comments ?** call **516-341-0666**, 8:30 am - 4:30 pm ET, Monday - Friday

**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

\*Reliable 1 Laboratories LLC is not affiliated with the owner of the trademark GENUINE BAYER®.

Distributed by: **Reliable 1 Laboratories LLC, Valley Stream, NY 11580**

**www.reliable1labs.com**

NDC 69618-016-01

Aspirin USP 325 mg (NSAID)

PAIN RELIEVER

FEVER REDUCER

100 TABLETS

\*Compare to Active Ingredient in GENUINE BAYER®

NDC 69618-016-01



**Aspirin USP**  
325 mg (NSAID)

**PAIN RELIEVER  
FEVER REDUCER**

**100 TABLETS**

†Compare to Active Ingredient in GENUINE BAYER®

**Drug Facts**

**Active ingredient (in each tablet)** Purpose  
Aspirin USP 325 mg (NSAID)\* ..... Pain reliever/Fever reducer  
\*nonsteroidal anti-inflammatory drug

**Uses** temporarily relieves ■ headache ■ menstrual pain  
■ minor pain of arthritis ■ muscle pain ■ toothache  
■ pain and fever of colds

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include ■ hives ■ facial swelling ■ asthma (wheezing) ■ shock

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding.  
The chance is higher if you  
■ are age 60 or older  
■ have had stomach ulcers or bleeding problems  
■ take a blood thinning (anticoagulant) or steroid drug  
■ take other drugs containing prescription or nonprescription NSAIDs (ibuprofen, naproxen, or others)  
■ take more or for a longer time than directed  
■ have 3 or more alcoholic drinks every day while using this product

**Drug Facts** continued on back of label  
**TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING**

†Reliable 1 Laboratories LLC is not affiliated with the owner of the trademark GENUINE BAYER®.

Distributed by: **Reliable 1 Laboratories LLC, Valley Stream, NY 11580**  
[www.reliable1labs.com](http://www.reliable1labs.com)

Rev. 0715



Lot No.:  
Exp. Date:  
**PEEL HERE FOR MORE DRUG FACTS**

**Drug Facts (continued)**

**Do not use** if you have ever had an allergic reaction to any other pain reliever/fever reducer

**Ask a doctor before use if**  
■ stomach bleeding warning applies to you  
■ you have a history of stomach problems, such as heartburn  
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease  
■ you are taking a diuretic  
■ you have asthma

**Ask a doctor or pharmacist before use if you are**  
■ taking a prescription drug for diabetes, gout or arthritis  
■ under a doctor's care for any serious condition  
■ taking any other drug

**Stop use and ask a doctor if**  
■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood  
■ have bloody or black stools  
■ have stomach pain that does not get better  
■ pain gets worse or lasts more than 10 days  
■ fever gets worse or lasts more than 3 days  
■ redness or swelling is present in the painful area  
■ any new symptoms appear  
■ ringing in the ears or loss of hearing occurs  
These could be signs of a serious condition

**If pregnant or breast-feeding,** ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.  
**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions** DRINK A FULL GLASS OF WATER WITH EACH DOSE ■ **adults and children 12 years and over:** 1-2 tablets every 4 hours while symptoms last, not more than 12 tablets in 24 hours  
■ **children under 12 years:** ask a doctor

**Other information** ■ store at room temperature 15°-30°C (59°-86°F)

**Inactive ingredients** corn starch, povidone

**Questions or comments?** call 516-341-0666, 8:30 am - 4:30 pm ET, Monday - Friday

### ASPIRIN 325 MG

aspirin tablet

#### Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:69618-016

Route of Administration

ORAL

**Active Ingredient/Active Moiety**

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

**Inactive Ingredients**

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
POVIDONE (UNII: FZ989GH94E)	

**Product Characteristics**

Color	white	Score	no score
Shape	ROUND	Size	10 mm
Flavor		Imprint Code	AP;0 10
Contains			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69618-016-01	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/01/2015	
2	NDC:69618-016-10	1000 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/01/2015	



# ASPIRIN 325MG Tablet

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part343	11/01/2015	

**Labeler** - Reliable 1 Laboratories LLC (079718111)

Revised: 2/2016

Reliable 1 Laboratories LLC