

COYDEN TYPE A MEDICATED ARTICLE- clopidol powder
Huvepharma, Inc.

Approved by the FDA under NADA # 034-393

COYDEN® 25%

(clopidol)

Type A Medicated Article

ACTIVE DRUG INGREDIENT: Clopidol.....113.5 gm/lb (25%)

INGREDIENTS: Rice Hulls, Mineral Oil, Fatty Acid Glycerides

BROILER/REPLACEMENT CHICKENS:

As an aid in the prevention of cecal and intestinal coccidiosis caused by *E. tenella*, *E. necatrix*,
E. acervulina, *E. maxima*, *E. brunetti*, *E. mivati*.

TURKEYS:

As an aid in the prevention of leucocytozoonosis caused by *Leucocytozoon smithi*.

SEE DIRECTIONS ON BACK PANEL

FOR THE MANUFACTURE OF COMPLETE FEEDS ONLY.

Manufactured by:

HUVEPHARMA, Inc. Peachtree City, GA 30269

Huvepharma, Inc.

525 Westpark Drive, Suite 230

Peachtree City, GA 30269

Distributed by:

HUVEPHARMA, Inc. Peachtree City, GA 30269

For questions, call 1-877-994-4883

® COYDEN is a registered trademark of Huvepharma, Inc.

Net Wt.: 50 lb (22.68 kg)

INDICATIONS FOR USE:

BROILER AND REPLACEMENT CHICKENS

As an aid in the prevention of coccidiosis in broilers and replacements for caged layers caused by

E. tenella, *E. necatrix*, *E. acervulina*, *E. maxima*, *E. brunetti*, *E. mivati*.

TURKEYS GROWN FOR MEAT PURPOSES ONLY

As an aid in the prevention of leucocytozoonosis caused by *Leucocytozoon smithi*.

USER SAFETY WARNING:

Not for Human Use. Keep Out of Reach of Children.

Contact Huvepharma Inc. at 1-877-994-4883 or <http://www.huvepharma.us>. For additional

information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-VETS or

<http://www.fda.gov/reportanimalae>.

WITHDRAWAL PERIODS AND RESIDUE WARNINGS:

CHICKENS: When feeding 0.025% clopidol, reduce dosage to 0.0125% or withdraw

medication 5 days before slaughter. Do not feed to chickens over 16 weeks of age.

TURKEYS: Withdraw medication 5 days before slaughter.

CAUTION: Consult a veterinarian or poultry pathologist if losses exceed 0.5% in a 2-day period.

MIXING DIRECTIONS:

For Complete Rations - Type C Medicated Products

COYDEN® 25% (clopidol) should be thoroughly and evenly mixed in the feed in accordance with

current good manufacturing practice for medicated feed. COYDEN 25% may be used to manufacture

a Type C medicated feed in the concentration range of 0.0125% to 0.025% clopidol.

To aid in even distribution of COYDEN 25% in the finished feed, prepare a mixture of COYDEN 25% in

a portion of complete feed ingredient before mixing into the finished ration. Blend the mixture with

the remainder of the finished feed and mix thoroughly.

DIRECTIONS FOR USE:

Must be mixed with other ingredients before feeding.

Mixing and conveying equipment should be properly grounded.

Use 1 pound of COYDEN 25% per ton of feed to produce a feed containing 0.0125% clopidol.

Use 2 pounds of COYDEN 25% per ton of feed to produce a feed containing 0.025% clopidol.

FEEDING DIRECTIONS:

Feed continuously, as the sole ration, either the 0.0125% or 0.025% Type C medicated feed

depending upon management practices, degree of exposure, and amount of feed eaten.


STORE IN A COOL PLACE

KEEP BAG TIGHTLY CLOSED

Restricted Drug (California)

Use only as Directed

TAKE TIME



OBSERVE LABEL DIRECTIONS

<p>INDICATIONS FOR USE: BROILERS AND REPLACEMENT CHICKENS: As an aid in the prevention of coccidiosis in broilers and replacement birds caused by <i>E. tenella</i>, <i>E. necatrix</i>, <i>E. oocervulina</i>, <i>E. maxima</i>, <i>E. brunetti</i>, <i>E. mivati</i>.</p> <p>TURKEYS GROWN FOR MEAT PURPOSES ONLY: As an aid in the prevention of leucocytozoosis caused by <i>Leucocytozoon smithi</i>.</p> <p>USER SAFETY WARNINGS: Not for Human Use. Keep Out of Reach of Children. Contact Huvepharma Inc. at 1-877-994-4383 or http://www.huvepharma.us. For additional information about reporting side effects for animal drugs, contact FDA at 1-888-FDA-1085 or http://www.fda.gov/medwatch.</p> <p>WITHDRAWAL PERIODS AND RESIDUE WARNINGS: CHICKENS: When feeding 0.025% clopidol, reduce dosage to 0.0125% or withdraw medication 5 days before slaughter. Do not feed to chickens over 16 weeks of age. TURKEYS: Withdraw medication 5 days before slaughter.</p> <p>CAUTION: Consult a veterinarian or poultry pathologist if losses exceed 0.5% in a 3-day period.</p> <p>MIXING DIRECTIONS: For Complete Rations - Type C Medicated Products COYDEN® 25% (clopidol) should be thoroughly and evenly mixed in the feed in accordance with current good manufacturing practice for medicated feed. COYDEN 25% may be used to manufacture a Type C medicated feed in the concentration range of 0.025% to 0.032% clopidol. To aid in even distribution of COYDEN 25% in the finished feed, prepare a mixture of COYDEN 25% in a portion of complete feed ingredient before mixing into the finished ration. Blend the mixture with the remainder of the finished feed and thoroughly.</p> <p>DIRECTIONS FOR USE: Must be mixed with other ingredients before feeding. Mixing and conveying equipment should be properly grounded. Use 1 pound of COYDEN 25% per ton of feed to produce a feed containing 0.025% clopidol. Use 2 pounds of COYDEN 25% per ton of feed to produce a feed containing 0.032% clopidol.</p> <p>FEEDING DIRECTIONS: Feed continuously, as the sole ration, using the 0.025% or 0.032% Type C medicated feed depending upon management practices, degree of exposure, and amount of feed eaten.</p> <p>STORE IN A COOL PLACE KEEP BAG TIGHTLY CLOSED</p> <p>RESTRICTED DRUG (California) Use only as Directed</p> <p>TAKE TIME OBSERVE LABEL DIRECTIONS</p> <p>COYDEN® 25% (clopidol) Type A Medicated Article</p> <p>9011-02 Rev. 03-2024</p>	<p>COYDEN® 25% (clopidol) Type A Medicated Article</p> <p>HUVEPHARMA</p>	<p>Approved by FDA under NADA # 034-393</p> <p>COYDEN® 25% (clopidol) Type A Medicated Article</p> <p>ACTIVE DRUG INGREDIENT: Clopidol.....113.5 gm/lb (25%)</p> <p>INGREDIENTS: Rice Hulls, Mineral Oil, Fatty Acid Glycerides</p> <p>BROILERS/REPLACEMENT CHICKENS: As an aid in the prevention of cecal and intestinal coccidiosis caused by <i>E. tenella</i>, <i>E. necatrix</i>, <i>E. oocervulina</i>, <i>E. maxima</i>, <i>E. brunetti</i>, <i>E. mivati</i>.</p> <p>TURKEYS: As an aid in the prevention of leucocytozoosis caused by <i>Leucocytozoon smithi</i>.</p> <p>SEE DIRECTIONS ON BACK PANEL.</p> <p>FOR THE MANUFACTURE OF COMPLETE FEEDS ONLY.</p> <p>Manufactured by: HUVEPHARMA, Inc., Peachtree City, GA 30269 Distributed by: HUVEPHARMA, Inc., Peachtree City, GA 30269 For questions, call 1-877-994-4383</p> <p>© COYDEN is a registered trademark of Huvepharma, Inc.</p> <p>Net Wt.: 50 lb (22.68 kg)</p> <p>Huvepharma, Inc. 225 Westpark Drive, Suite 230 Peachtree City, GA 30269</p> <p>HUVEPHARMA</p>	<p>COYDEN® 25% (clopidol) Type A Medicated Article</p> <p>HUVEPHARMA</p> <p>LOT NO & EXP DATE</p> <p>000000000111 0 04 1807-0000</p>
---	--	--	---

COYDEN TYPE A MEDICATED ARTICLE

clopidol powder

Product Information

Product Type	OTC TYPE A MEDICATED ARTICLE ANIMAL DRUG	Item Code (Source)	NDC:23243-9568
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CLOPIDOL (UNII: 8J763HFF5N) (CLOPIDOL - UNII:8J763HFF5N)	CLOPIDOL	250.2 g in 1 kg

Inactive Ingredients

Ingredient Name	Strength
MINERAL OIL (UNII: T5L8T28FGP)	
RICE BRAN (UNII: R60QEP13IC)	

Product Characteristics

Color	brown ((Light Tan to Brown))	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:23243-9568-5	22.68 kg in 1 BAG		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
NADA	NADA034393	05/29/2009	

Labeler - Huvepharma, Inc. (619153559)

Registrant - Huvepharma EOOD (552671651)

Establishment

Name	Address	ID/FEI	Business Operations
Huvepharma, Inc.		883128204	manufacture, analysis, pack, label, medicated animal feed manufacture

