

ALKA-SELTZER HEARTBURN AND PAIN- anhydrous citric acid, aspirin, and sodium bicarbonate tablet, effervescent
Bayer HealthCare LLC.

Alka-Seltzer® Heartburn + Pain UI1600130

Drug Facts

Active ingredients (in each tablet)	Purpose
Anhydrous citric acid 1000 mg	Antacid
Aspirin 325 mg (NSAID) *	Analgesic
Sodium bicarbonate (heat-treated) 1916 mg	Antacid

* nonsteroidal anti-inflammatory drug

Uses

- For the temporary relief of minor aches and pains with heartburn, sour stomach, or acid indigestion and upset stomach associated with these symptoms
- for temporary relief of headaches, body aches, backache and pain alone

Warnings

Reye's syndrome

Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert

Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning

This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

STOMACH WARNING: TO AVOID SERIOUS INJURY, DO NOT TAKE UNTIL TABLET IS COMPLETELY DISSOLVED. IT IS VERY IMPORTANT NOT TO TAKE THIS PRODUCT WHEN OVERLY FULL FROM FOOD OR DRINK. Consult a doctor if severe stomach pain occurs after taking this product.

Do not use

- if you have ever had an allergic reaction to any other pain reliever/fever reducer
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma
- you have a sodium-restricted diet

Ask a doctor or pharmacist before use if you are

- presently taking a prescription drug. Antacids may interact with certain prescription drugs.
- taking a prescription drug for diabetes, gout, or arthritis

When using this product do not take more than the maximum daily dosage in a 24-hour period, or use the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a physician.

Stop use and ask a doctor if

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
- ringing in the ears or a loss of hearing occurs
- pain get worse or last more than 10 days
- redness or swelling is present
- any new symptoms appear

If pregnant or breast-feeding, ask a health professional before use. **It is especially important not to use aspirin at 20 weeks or later in pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.**

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center right away.

Directions

Directions

● Completely dissolve 2 tablets in 4 ounces of water and be sure bubbling has stopped before drinking

● Do not exceed recommended dose. **SEE WARNINGS**

adults and children 12 years and over	2 tablets every 4 hours, or as directed by a doctor	do not exceed 8 tablets in 24 hours
adults 60 years and over	2 tablets every 4 hours, or as directed by a doctor	do not exceed 4 tablets in 24 hours
children under 12 years	consult a doctor	

Other information

- **each tablet contains:** sodium 568 mg
- store at room temperature. Avoid excessive heat.
- Alka-Seltzer Heartburn + Pain in water contains principally the antacid sodium citrate and the analgesic sodium acetylsalicylate

Inactive ingredients

none

Questions or comments?

1-800-986-0369 (Mon - Fri 9AM - 5PM EST)

Dist. by: Bayer HealthCare LLC

Whippany, NJ 07981

PRINCIPAL DISPLAY PANEL - 24 Tablet Carton



ALKA-SELTZER HEARTBURN AND PAIN

anhydrous citric acid, aspirin, and sodium bicarbonate tablet, effervescent

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0280-0160
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL) (ANHYDROUS CITRIC ACID - UNII:XF417D3PSL)	ANHYDROUS CITRIC ACID	1000 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg
SODIUM BICARBONATE (UNII: 8MDF5V39QO) (BICARBONATE ION - UNII:HN1ZRA3Q20)	SODIUM BICARBONATE	1916 mg

Product Characteristics

Color	white	Score	no score
--------------	-------	--------------	----------

Shape	ROUND	Size	25mm
Flavor		Imprint Code	ASP;SAC
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0280-0160-01	58 in 1 CARTON	02/03/2025	
1		2 in 1 POUCH; Type 0: Not a Combination Product		
2	NDC:0280-0160-02	6 in 1 CARTON	02/03/2025	
2		2 in 1 POUCH; Type 0: Not a Combination Product		
3	NDC:0280-0160-03	12 in 1 CARTON	02/03/2025	
3		2 in 1 POUCH; Type 0: Not a Combination Product		
4	NDC:0280-0160-04	18 in 1 CARTON	02/03/2025	
4		2 in 1 POUCH; Type 0: Not a Combination Product		
5	NDC:0280-0160-05	36 in 1 CARTON	02/03/2025	
5		2 in 1 POUCH; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	02/03/2025	

Labeler - Bayer HealthCare LLC. (112117283)

Revised: 12/2024

Bayer HealthCare LLC.