### SUKANG BZK ANTISEPTIC TOWELETTE - benzalkonium chloride liquid Taizhou Kanglin Health Protection Products Co Ltd

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

-----

### SUKANG BZK Antiseptic Towelette

### SUKANG BZK Antiseptic Towelette

#### **Active Ingredient**

Benzalkonium Chloride, 0.13% v/v

#### Purpose

Antiseptic

#### Uses

First aid antiseptic to help prevent skin infection in: minor cuts, scrapes, burns

Antiseptic cleansing

Perineal and maternity care

#### Warnings

#### For external use only.

#### Do not use

- as an antiseptic for more than 1 week
- in the eyes
- over large areas of the body

#### Ask a doctor before use if you have

- deep or puncture wounds
- animal bites
- serious burns

#### Stop use

- if irritation and redness develops
- if condition persists for more than 72 hours, consult a physician

#### Keep out of reach of children

• If swallowed, get medical help or contact a Poison Control Center right away.

#### Directions

Tear at notch

Remove towelette Use only once **As a first aid antiseptic** Clean affected area Apply 1 to 3 times daily May be covered with a sterile bandage If bandaged, let dry first

### **Other Information**

- Store at room temperature
- Avoid excessive heat

#### **Inactive Ingredient**

water

## SUKANG BZK Antiseptic Towelette 100 count (22408-001-01)



Drug Facts	100 5" x 7"	100 Packets 5" x 7"				
Active IngredientPurpose Benzalkonium Chloride, 0.13% v/vAntiseptic	Packets	Packets				
Uses First aid antiseptic to help prevent skin infection in: • Minor Cuts • Scrapes • Burns • Antiseptic Cleansing • Perineal and maternity care	BZK					
Warnings	the second se	BZK 💦 🛞				
For External Use Only.	Antiseptic	Antiseptic				
Do not use • as an antiseptic for more than 1 week • in the eyes • over large areas of the body	Towelettes	Towelettes				
Ask a doctor before use if you have • deep or puncture wounds • animal bites • serious burns	And Hospital Use	For Professional SUKANG				
Stop use • if irritation and redness develop • If condition persists more than 72 hours, consult a physician.	47%	And Hospital Osc				
Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.						
Directions • Tear at notch • Remove towelette • Use only once	SUKANG					
As a first aid antiseptic • Clean affected area • Apply 1 to 3 times daily • May be covered with a sterile bandage • If bandaged, let dry first	Manufactured by: Taizhou Kanglin Health					
Other information • Store at room temperature • Avoid excessive heat	Protection Products Co.Ltd.					
Inactive Ingredients Water						
-01						
ade in China						

SUKANG BZK a		TIC TOWELE	ГТЕ						
	. iiquiu								
Product Information	on								
Product TypeHUMAN OTC DRUGItem			Ite m C	Code (Source)			NDC:22408-001		
Route of Administrati	on	TOPICAL							
Active Ingredient/A	Active Moi	etv							
ficure ingreatent		redient Name			Basis of Strength			Strength	
BENZALKONIUM CHLORIDE (UNII: F5UM2KM3W7) (BENZALKONIUM - UNII:7N6JUD5X6Y)				BENZALKONIUM CHLORIDE			0.13 g in 100 g		
Inactive Ingredien	ts								
Ingredient Name						Strength			
WATER (UNII: 059QF0K	(00R)								
Packaging									
# Item Code	Pac	kage Description	Marketing Start Da		Date	Marketing		End Date	
<b>1</b> NDC:22408-001-01	100 in 1 E	BOX							
1	1.7 g in 1	PACKET							
Marketing Info	rmation								
Marketing Category				Marketing Start Date			Marketing End Date		
OTC monograph final	part333E		-	02/13/20	-			2	

# Labeler - Taizhou Kanglin Health Protection Products Co Ltd (530576131)

## **Registrant** - Taizhou Kanglin Health Protection Products Co Ltd (530576131)

# Establishment

Name	Address	ID/FEI	<b>Business Operations</b>
Taizhou Kanglin Health Protection Products Co Ltd		530576131	manufacture (22408-001)

Revised: 2/2013

Taizhou Kanglin Health Protection Products Co Ltd