

SIELKIN QUIT SMOKING NICOTINE PATCHES- nicotine patch
YITONGBADA (SHENZHEN) INTERNATIONAL TRADE CO., LTD

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click [here](#).

Drug Facts

Drug Facts

Active ingredients (in each patch)

Nicotine 7mg delivered over 24 hours

Purpose

Stop smoking aid

Use reduces withdrawal symptoms, including nicotine cravings associated with quitting smoking.

Warnings

For external use only

Ask a doctor before use if you have

■ heart disease ■ high blood pressure ■ skin allergies ■ diabetes ■ history of seizures

Ask a doctor or pharmacist before use if you are ■ using a non-nicotine stop smoking drug ■ pregnant or breast-feeding

When using this product if you have vivid dreams or other sleep disturbances remove this patch at bedtime

Stop use and ask a doctor if irritation or allergic reactions occur

Keep out of reach of children and pets. In case of accidental ingestion, seek medical advice immediately or contact a Poison Control Center (1-800-222-1222)

Directions

■ if you are under 18 years of age, ask a doctor before use.

■ Before use, carefully read the insert for complete instructions.

■ If you smoke more than 10 cigarettes per day, follow this 10-week schedule:

	Step 1		Step 2	Step 3
Cigarettes per Day	Week 1 (Key Adjustment Phase)	Weeks 2-6	Weeks 7-8	Weeks 9-10
10-20 cigarettes	two 21mg patch/day	one 21mg patch/day	one 14mg patch/day	one 7mg patch/day
20+ cigarettes	three 21mg patch/day	two 21mg patch/day	one 14mg patch/day	one 7mg patch/day

■ If you smoke 10 or fewer cigarettes per day, skip **STEP 1 (21 mg)**. Start with **STEP 2 (14 mg)** for 6 weeks—using two 14 mg patches per day for the first week, then one 14 mg patch per day for weeks 2 to 6. Then move to **STEP 3 (7 mg)** for 2 weeks, and stop.

Other information Store at 20-25°C (68-77°F)

Inactive ingredients camphor, ethanol, acrylate adhesive, aluminized polyester, cellulose paper, methacrylic acid copolymer

Questions or comments? Contact us at: cs@sielkin.com

Active ingredients (in each patch)
Nicotine 7mg delivered over 24 hours
Stop smoking aid

reduces withdrawal symptoms, including nicotine cravings associated with quitting smoking.

Ask a doctor before use if you have

■heart disease ■high blood pressure ■skin allergies ■diabetes ■history of seizures

Ask a doctor or pharmacist before use if you are■using a non-nicotine stop smoking drug■pregnant or breast-feeding

When using this product

if you have vivid dreams or other sleep disturbances remove this patch at bedtime.

Stop use and ask a doctor if irritation or allergic reactions occur

Keep out of reach of children and pets. In case of accidental ingestion, seek medical advice immediately or contact a Poison Control Center(1-800-222-1222)

Warnings

For external use only

Directions

■ **if you are under 18 years of age, ask a doctor before use.**

■Before use, carefully read the insert for complete instructions.

■If you smoke more than 10 cigarettes per day, follow this 10-week schedule:

	Step 1		Step 2	Step 3
Cigarettes per Day	Week 1(Key Adjustment Phase)	Weeks 2-6	Weeks 7-8	Weeks 9-10
10-20 cigarettes	two 21mg patch/day	one 21mg patch/day	one 14mg patch/day	one 7 mg patch/day
20+ cigarettes	three 21mg patch/day	two 21mg patch/day	one 14mg patch/day	one 7 mg patch/day

■If you smoke 10 or fewer cigarettes per day, skip **STEP 1(21mg)**. Start with **STEP 2(14mg)** for 6 weeks-using two14mg patches per day for the first week, then one 14mg patch per day for weeks 2 to 6. Then move to **STEP 3(7mg)** for 2weeks, and stop.

Other information Store at 20-25°C (68-77°F)

Inactive ingredients camphor, ethanol, acrylate adhesive, aluminized polyester, cellulose paper, methacrylic acid copolymer

SIELKIN QUIT SMOKING NICOTINE PATCHES

nicotine patch

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83364-015
Route of Administration	TRANSDERMAL		

Active Ingredient/Active Moiety				
Ingredient Name			Basis of Strength	Strength
NICOTINE (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)			NICOTINE	7 mg in 1 g
Inactive Ingredients				
Ingredient Name				Strength
BORNEOL (UNII: M89NIB437X)				
ALCOHOL (UNII: 3K9958V90M)				
METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:1) (UNII: 74G4R6TH13)				
ACRYLIC ACID (UNII: J94PBK7X8S)				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83364-015-01	30 in 1 BOX	12/20/2024	
1		0.007 g in 1 PATCH; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category		Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other			12/20/2024	

Labeler - YITONGBADA (SHENZHEN) INTERNATIONAL TRADE CO., LTD (725220463)