

**BENZOIN TINCTURE- benzoin resin liquid
116956644**

Foster & Thrive Benzoin Tincture

Drug Facts

Active Ingredient

Benzoin Tincture

Purpose

Oral mucosal protectant

Use

Forms a coating over wound for protecting recurring canker sores

Warnings

For external use only. Do not swallow. Do not exceed recommended dosage.

When using this product

do not use in the eyes

do not use this product for longer than 1 week unless directed by a dentist or doctor

Stop use and consult a dentist or doctor if

- the condition persists or gets worse
- symptoms do not improve in 7 days.
- pain, irritation, swelling, redness, rash or fever develop

Keep out of the reach of children.

In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately. (1-800-222-1222)

Directions

Adult and children 6 months of age and older: Dry the affected area. Saturate a cotton applicator with medication and apply undiluted to the affected area not more than every 2 hours.

Children under 6 months of age: Consult a dentist or doctor.

Inactive Ingredients

Alcohol 79%, Aloe, Stoax, Tolu Balsam.

Questions?

Call 833-358-6431

Principal Display Panel

NDC 70677-1200-1

TAMPER EVIDENT: DO NOT USE IF SHRINK BAND AROUND CAP IS BROKEN OR MISSING.

Foster  Thrive™

Benzoin Tincture

USP XV
ORAL MUCOSAL PROTECTANT
Alcohol 79%

FLAMMABLE: Keep away from spark, heat & flame

© 2023 McKesson Corporation
Distributed by: McKesson Corp.
via SSSL, Memphis, TN 38141
Money Back Guarantee
www.fosterandthrive.com
Rev 07/24

2 FL oz (59 mL)

10939 95960 117

Drug Facts	Purpose Oral Mucosal Protectant
Active ingredient Benzoin Tincture	Forms a coating over wound for protecting recurring canker sores
Uses	For External Use Only. Do not swallow. Do not exceed recommended dosage. • FLAMMABLE: Keep away from spark, heat & flame.
Warnings	Stop use and consult a dentist or doctor promptly if • the condition persists or gets worse • symptoms do not improve within 7 days • pain, irritation, swelling, redness, rash or fever develop.
Directions	When using this product do not use in the eyes • do not use this product for longer than 1 week unless directed by a dentist or doctor. Keep out of reach of children. In case of accidental ingestion, seek professional assistance or contact a Poison Control Center immediately, (1-800-222-1222)
Other information	Adults and children 6 months of age and older: Dry the affected area. Saturate a cotton applicator with medication and apply undiluted to the affected area not more often than every 2 hours. Children under 6 months of age: Consult a dentist or doctor. Store at room temperature 15°-30°C (59°-86°F) in tight, light-resistant containers. • Avoid exposure to direct sunlight and excessive heat.
Inactive ingredients	Alcohol 79%, Aloe, Storax, and Tolu Balsam
Questions?	Call 833-358-6431

BENZOIN TINCTURE

benzoin resin liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70677-1200
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZOIN RESIN (UNII: GK21SBA74R) (BENZOIN RESIN - UNII:GK21SBA74R)	BENZOIN RESIN	1000 mg in 1 mL

Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
ALOE (UNII: V5VD430YW9)	
TOLU BALSAM (UNII: TD2LE91MBE)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70677-1200-1	59 mL in 1 BOTTLE, GLASS; Type 0: Not a Combination Product	12/17/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M022	12/17/2024	

Labeler - 116956644 (116956644)

Registrant - Pharma Nobis, LLC (118564114)

Establishment

Name	Address	ID/FEI	Business Operations
Pharma Nobis, LLC		118564114	manufacture(70677-1200) , analysis(70677-1200) , pack(70677-1200) , label(70677-1200)

Revised: 12/2024

116956644