

**FOSTER AND THRIVE NIGHTTIME SEVERE COLD AND FLU RELIEF-
acetaminophen, dextromethorphan hydrobromide, doxylamine succinate,
phenylephrine hydrochloride solution
Strategic Sourcing Services LLC**

McKesson Nighttime Severe Cold & Flu Relief Drug Facts

Active ingredients (in each 15 mL)

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Doxylamine Succinate 6.25 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Nasal decongestant

Uses

temporarily relieves common cold/flu symptoms:

- nasal congestion
- sinus congestion and pressure
- cough due to minor throat and bronchial irritation
- cough to help you sleep
- minor aches and pains
- sore throat
- headache
- fever
- runny nose and sneezing
- reduces swelling of nasal passages
- temporarily restores freer breathing through the nose
- promotes nasal and/or sinus drainage

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours

- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- to make a child sleepy

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- glaucoma
- thyroid disease
- diabetes
- cough that occurs with too much phlegm (mucus)
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **do not use more than directed**
- excitability may occur, especially in children

- marked drowsiness may occur
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

Stop use and ask a doctor if

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

Other information

- **each 15 mL contains:** sodium 15 mg
- store at 20-25°C (68-77°F)

Inactive ingredients

anhydrous citric acid, D&C yellow #10, edetate disodium, FD&C green #3, FD&C red #40, FD&C yellow #6, flavor, glycerin, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose, xanthan gum

Questions?

1-800-719-9260

Package/Label Principal Display Panel

COMPARE TO VICKS® NYQUIL® SEVERE HONEY FLAVOR ACTIVE INGREDIENTS

Foster & Thrive™

NIGHTTIME SEVERE Cold & Flu Relief

Acetaminophen

Phenylephrine HCl

Dextromethorphan HBr

Doxylamine Succinate

Pain Reliever

Fever Reducer

Cough Suppressant

Antihistamine

Nasal Decongestant

Gluten Free

Honey Flavor

12 FL OZ 355 mL

NDC70677-1030-1

COMPARE TO VICKS® NYQUIL® SEVERE HONEY FLAVOR ACTIVE INGREDIENTS*

Foster & Thrive™ NIGHTTIME SEVERE Cold & Flu Relief

Acetaminophen
Phenylephrine HCl
Dextromethorphan HBr
Doxylamine Succinate

Pain Reliever
Fever Reducer
Cough Suppressant
Antihistamine
Nasal Decongestant
Gluten Free

HONEY FLAVOR

12 FLOZ 355 mL

: 8LS4D KM F1

REV. 03/23

©2023 McKesson Corporation
Distributed by: McKesson Corp.,
via SSSL, Memphis, TN 38141
Money Back Guarantee
www.fosterandthrive.com

REV. 03/23

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

DO NOT USE IF PRINTED
NECKBAND IS BROKEN
OR MISSING

Drug Facts

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg.....	Pain reliever/ fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Doxylamine Succinate 6.25 mg.....	Antihistamine
Phenylephrine HCl 5 mg.....	Nasal decongestant

Uses temporarily relieves common cold/flu symptoms: ■ nasal congestion ■ sinus congestion and pressure ■ cough due to minor throat and bronchial irritation ■ cough to help you sleep ■ minor aches and pains ■ headache ■ sore throat ■ fever ■ runny nose and sneezing ■ reduces swelling of nasal passages ■ temporarily restores freer breathing through the nose ■ promotes nasal and/or sinus drainage

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if ■ adult takes more than 4,000 mg of acetaminophen in 24 hours ■ child takes more than 5 doses in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product
Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash If a skin reaction occurs, stop use and seek medical help right away.
Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

PEEL BACK AT
CORNER FOR MORE
INFORMATION

: 8LS4D KM B1



Drug Facts (continued)


Do not use ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients ■ to make a child sleepy
Ask a doctor before use if you have ■ liver disease ■ heart disease ■ high blood pressure ■ glaucoma ■ thyroid disease ■ diabetes ■ cough that occurs with too much mucus (trouble breathing)

YESIVE AREA
NO VARNISH • NO TYPE

Drug Facts (continued)

■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness
Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion, or cough gets worse or lasts more than 5 days (children) or 7 days (adults) ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
If pregnant or breast-feeding, ask a health professional before use.
Keep out of reach of children. **Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.
Directions ■ take only as directed - see Overdose warning ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs

ADHESIVE AREA
NG • NO VARNISH • NO TYPE



FOSTER AND THRIVE NIGHTTIME SEVERE COLD AND FLU RELIEF

acetaminophen, dextromethorphan hydrobromide, doxylamine succinate, phenylephrine hydrochloride solution

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70677-1030
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
DOXYLAMINE SUCCINATE (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	6.25 mg in 15 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
FD&C GREEN NO. 3 (UNII: 3P3ONR601S)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	

WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM CITRATE, UNSPECIFIED FORM (UNII: 1Q73Q2JULR)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
XANTHAN GUM (UNII: TTV12P4NEE)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70677-1030-1	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	06/30/2023	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	06/30/2023	

Labeler - Strategic Sourcing Services LLC (116956644)

Revised: 11/2024

Strategic Sourcing Services LLC