

GILTUSS D- giltuss d syrup
Gil Pharmaceutical Corp

Giltuss-D
ALLERGY & CONGESTION
ANTIHISTAMINIC AND NASAL DECONGESTANT

Drug Facts

Active ingredients (in each 10 mL) Purpose

Chlorpheniramine maleate 3.5 mg Antihistamine

Phenylephrine HCl 10 mg Nasal decongestant

Uses

Temporarily relieves these Symptoms due to hay fever (allergic rhinitis) or other upper-respiratory allergies:

- runny nose
- Sneezing
- Itchy, watery eyes
- nasal congestion
- itching of the nose or throat
- sinus congestion and pressure

Warnings

Do not use

- to sedate a child or to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug.
If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- glaucoma
- a breathing problem or persistent or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema

Ask a doctor or pharmacist before use if you are

- taking sedatives or tranquilizers

When using this product

- **do not use more than directed**
- may cause drowsiness
- avoid alcoholic beverages
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- You get nervous, dizzy, or sleepless
- symptoms do not get better within 7 days or are accompanied by fever

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- **not take more than 6 doses in any 24-hour period**
- **use enclosed dosing cup**
- **keep dosage cup for use with this product**
- **mL = milliliter**

Adults and children 12 years and over	10 mL every 4 hours
Children 6 to under 12 years	5 mL every 4 hours
Children under 6 years	Consult a doctor

Other information

- store at room temperature 15°-30°C (59°-86°F)
- close cap tightly
- **SAVE CARTON FOR COMPLETE DRUG FACTS**

Inactive ingredients

Citric Acid, Flavor, Glycerin, Methylparaben, Propylene Glycol, Propylparaben, Purified Water, Sodium Carboxymethylcellulose, Sucralose.

Questions or comments?



Call 1-787-848-9114, Monday-Friday 9AM-5PM EST. Call your doctor for medical advice in the event of side effects.

Label revised: 04/24

**MANUFACTURED FOR
GIL PHARMACEUTICAL CORP.
PONCE, PUERTO RICO 00717-1565**

PRINCIPAL DISPLAY PANEL

Giltuss-D ALLERGY & CONGESTION - NDC 58552-147-04 - Bottle Label

Drug Facts	<p>NO PSEUDOEPHEDRINE PSEUDOEPHEDRINA</p>  <p>Giltuss-D ALLERGY & CONGESTION ALERGIA Y CONGESTIÓN</p> <p>DYE FREE, PRESERVATIVE FREE, SODIUM FREE, SUGAR FREE SIN COLORANTES, SIN PRESERVATIVOS, LIBRE DE SODIO, SIN AZÚCAR</p> <p>SNEEZING, RUNNY NOSE, ITCHY & WATERY EYES, ITCHY THROAT OR NOSE, SINUS PRESSURE, STUFFY NOSE ESTORNIDOS, GOTEOS NASAL, PICOR Y OJOS LLOROSOS, PICOR DE GARGANTA Y/O NARIZ, PRESIÓN POR SINUSITIS, NARIZ TAPADA</p> <p>TUTTI FRUITI FLAVOR SABOR A TUTTI FRUITI</p> <p>4FL OZ (118 mL)</p>	<p>When using this product • do not use more than directed • may cause drowsiness • avoid alcoholic beverages • alcohol, sedatives, and tranquilizers may increase drowsiness • avoid alcoholic drinks • be careful when driving a motor vehicle or operating machinery • excitability may occur, especially in children</p>						
<p>Active Ingredients (in each 10 mL)</p> <p>Chlorpheniramine maleate, 3.5 mg Antihistamine Pseudoephedrine HCl, 10 mg Nasal decongestant</p>		<p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).</p>	<p>MANUFACTURED FOR GIL PHARMACEUTICAL CORP. PONCE, PUERTO RICO 00717-1586</p>					
<p>Uses Temporarily relieves these symptoms due to hay fever (allergic rhinitis) or other upper-respiratory allergies:</p> <ul style="list-style-type: none"> runny nose sneezing itchy, watery eyes nasal congestion itching of the nose or throat sinus congestion and pressure 		<p>Sip use and ask a doctor if • if you get nervous, dizzy, or sleepless • symptoms do not get better within 7 days or are accompanied by fever</p>						
<p>Warnings</p> <p>Do not use • to sedate a child or to make a child sleepy • if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.</p>		<p>Directions • not take more than 6 doses in any 24-hour period • use enclosed dosing cup • keep dosage cup for use with this product • mL = milliliter</p> <table border="1"> <tr> <td>Adults and children 12 years and over</td> <td>10 mL every 4 hours</td> </tr> <tr> <td>Children 6 to under 12 years</td> <td>5 mL every 4 hours</td> </tr> <tr> <td>Children under 6 years</td> <td>Consulta doctor</td> </tr> </table>		Adults and children 12 years and over	10 mL every 4 hours	Children 6 to under 12 years	5 mL every 4 hours	Children under 6 years
Adults and children 12 years and over	10 mL every 4 hours							
Children 6 to under 12 years	5 mL every 4 hours							
Children under 6 years	Consulta doctor							
<p>Ask a doctor before use if you have:</p> <ul style="list-style-type: none"> heart disease high blood pressure thyroid disease diabetes trouble urinating due to an enlarged prostate gland glaucoma a breathing problem or persistent or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> taking sedatives or tranquilizers 	<p>Other information • store at room temperature 15°-30°C (59°-86°F). • close cap tightly • SEE CAUTION FOR COMPLETE DRUG FACTS</p> <p>Inactive ingredients Citric Acid, Flavor, Glycerin, Methylparaben, Propylene Glycol, Propylparaben, Purified Water, Sodium Carboxymethylcellulose, Sucrose.</p> <p>Questions or comments? Call 1-787-848-9114, Monday-Friday 9AM-5PM EST. Call your doctor for medical advice in the event of side effects.</p>							

Giltuss-D ALLERGY & CONGESTION - NDC 58552-147-04 - Carton Label



GILTUSS D

giltuss d syrup

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:58552-147
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

CHLORPHENIRAMINE MALEATE (UNII: V1Q0O9OJ9Z) (CHLORPHENIRAMINE - UNII:3U6IO1965U)	CHLORPHENIRAMINE MALEATE	3.5 mg
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE	10 mg

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID ACETATE (UNII: DSO12WL7AU)	
GLYCERIN (UNII: PDC6A3C0OX)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
CARBOXYMETHYLCELLULOSE SODIUM (UNII: K679OBS311)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	TUTTI FRUTTI	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:58552-147-04	1 in 1 CARTON	04/11/2024	
1		1 in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	04/11/2024	

Labeler - Gil Pharmaceutical Corp (176826592)