

**PEPTO NAUSEA MOTION SICKNESS- meclizine hcl tablet
THE PROCTER & GAMBLE MANUFACTURING COMPANY**

Pepto Nausea MOTION SICKNESS

Pepto Nausea MOTION SICKNESS

Drug Facts

Active ingredient (in each tablet)

Meclizine HCl 25 mg

Purpose

Antiemetic

Uses

for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness

Warnings

Do not use

in children under 12 years of age unless directed by a doctor.

Ask a doctor before use if you have

- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- difficulty in urination due to enlargement of the prostate gland

Ask a doctor or pharmacist before use if you are

taking sedatives or tranquilizers.

When using this product

- drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

If pregnant or breast feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- take ½ to 1 hour before traveling
- adults and children 12 years and over: 1 to 2 tablets once daily or as directed by a doctor

Other information

- Store at room temperature 20°-25°C (68°-77°F).

Inactive ingredients

anhydrous lactose, corn starch, FD&C Red No. 40 Aluminum Lake, magnesium stearate, microcrystalline cellulose, silicon dioxide

Questions?

1-800-717-3786

TAMPER EVIDENT: CARTON SEALED FOR YOUR PROTECTION. DO NOT USE IF CARTON IS OPENED OR TORN. IMPORTANT: KEEP CARTON FOR FUTURE REFERENCE.

DIST. BY: PROCTER &

GAMBLE, CINCINNATI, OH

45202

PRINCIPAL DISPLAY PANEL

Meclizine HCl -
Antiemetic

Pepto

Nausea

MOTION

SICKNESS

TARGETED RELIEF

TREAT + PREVENT

NAUSEA

LESS DROWSY

10 TABLETS

Meclizine HCl -
Antiemetic

Pepto[®] Nausea

MOTION SICKNESS

**TARGETED
RELIEF**
TREAT + PREVENT
NAUSEA



LOT
EXP



21237744
R53224



PATENTS: www.pg.com/patents
DIST. BY: PROCTER & GAMBLE, CINCINNATI, OH 45202



| |
|--|
| • take ½ to 1 hour before traveling |
| • adults and children 12 years and over: 1 to 2 tablets once daily or as directed by a doctor |
| Other information |
| • Store at room temperature 20°-25°C (68°-77°F). |
| Inactive ingredients anhydrous lactose, corn starch, FD&C Red No. 40 Aluminum Lake, magnesium stearate, microcrystalline cellulose, silicon dioxide |
| Questions? 1-800-717-3786 |

TAMPER EVIDENT: CARTON SEALED FOR YOUR PROTECTION. DO NOT USE IF CARTON IS OPENED OR TORN. IMPORTANT: KEEP CARTON FOR FUTURE REFERENCE.

| |
|--|
| Drug Facts |
| Active ingredient (in each tablet) Meclizine HCl 25 mg.....Antiemetic |
| Uses for the prevention and treatment of nausea, vomiting, or dizziness associated with motion sickness |
| Warnings Do not use in children under 12 years of age unless directed by a doctor. Ask a doctor before use if you have • a breathing problem such as emphysema or chronic bronchitis • glaucoma • difficulty in urination due to enlargement of the prostate gland |
| Ask a doctor or pharmacist before use if you are taking sedatives or tranquilizers. |
| When using this product • drowsiness may occur • avoid alcoholic drinks • alcohol, sedatives, and tranquilizers may increase drowsiness • use caution when driving a motor vehicle or operating machinery |
| If pregnant or breast feeding , ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. |
| Directions |

PEPTO NAUSEA MOTION SICKNESS

meclizine hcl tablet

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:84126-340 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------------|----------|
| MECLIZINE HYDROCHLORIDE (UNII: HDP7W44CIO) (MECLIZINE - UNII:3L5TQ84570) | MECLIZINE HYDROCHLORIDE | 25 mg |

Inactive Ingredients

| Ingredient Name | Strength |
|---|----------|
| ANHYDROUS LACTOSE (UNII: 3S5Y5LH9PMK) | |
| STARCH, CORN (UNII: O8232NY3SJ) | |
| FD&C RED NO. 40 ALUMINUM LAKE (UNII: 6T47AS764T) | |
| MAGNESIUM STEARATE (UNII: 70097M6I30) | |
| MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U) | |
| SILICON DIOXIDE (UNII: ETJ7Z6XBU4) | |

Product Characteristics

| | | | |
|-----------------|-------|---------------------|----------|
| Color | pink | Score | no score |
| Shape | ROUND | Size | 9mm |
| Flavor | | Imprint Code | PN |
| Contains | | | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:84126-340-01 | 1 in 1 CARTON | 03/03/2025 | |
| 1 | | 10 in 1 VIAL, PLASTIC; Type 0: Not a Combination Product | | |
| 2 | NDC:84126-340-03 | 3 in 1 CELLO PACK | 03/03/2026 | |
| 2 | | 10 in 1 VIAL, PLASTIC; Type 0: Not a Combination Product | | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M009 | 03/03/2025 | |

Labeler - THE PROCTER & GAMBLE MANUFACTURING COMPANY (004238200)

Revised: 2/2026

THE PROCTER & GAMBLE MANUFACTURING COMPANY