

**SIELKIN QUIT SMOKING NICOTINE PATCHES 14MG- nicotine patch  
YITONGBADA (SHENZHEN) INTERNATIONAL TRADE CO., LTD**

*Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.*

-----

**Sielkin QUIT SMOKING NICOTINE PATCHES 14mg**

Drug Facts

# Drug Facts

## Active ingredients ( in each patch )

Nicotine 14mg delivered over 24 hours ..... Purpose Stop smoking aid

**Use** reduces withdrawal symptoms, including nicotine cravings associated with quitting smoking.

## Warnings

For external use only

### Ask a doctor before use if you have

■ heart disease ■ high blood pressure ■ skin allergies ■ diabetes ■ history of seizures

Ask a doctor or pharmacist before use if you are ■ using a non-nicotine stop smoking drug ■ pregnant or breast-feeding

When using this product if you have vivid dreams or other sleep disturbances remove this patch at bedtime

Stop use and ask a doctor if irritation or allergic reactions occur

Keep out of reach of children and pets. In case of accidental ingestion, seek medical advice immediately or contact a Poison Control Center (1-800-222-1222)

## Directions

■ if you are under 18 years of age, ask a doctor before use.

■ Before use, carefully read the insert for complete instructions.

■ If you smoke more than 10 cigarettes per day, follow this 10-week schedule:

	Step 1		Step 2	Step 3
Cigarettes per Day	Week 1 (Key Adjustment Phase)	Weeks 2-6	Weeks 7-8	Weeks 9-10
10-20 cigarettes	two 21mg patch/day	one 21mg patch/day	one 14mg patch/day	one 7mg patch/day
20+ cigarettes	three 21mg patch/day	two 21mg patch/day	one 14mg patch/day	one 7mg patch/day

■ If you smoke 10 or fewer cigarettes per day, skip **STEP 1 (21 mg)**. Start with **STEP 2 (14 mg)** for 6 weeks—using two 14 mg patches per day for the first week, then one 14 mg patch per day for weeks 2 to 6. Then move to **STEP 3 (7 mg)** for 2 weeks, and stop.

**Other information** Store at 20-25°C (68-77°F)

**Inactive ingredients** camphor, ethanol, acrylate adhesive, aluminized polyester, cellulose paper, methacrylic acid copolymer

**Questions or comments?** Contact us at: [cs@sielkin.com](mailto:cs@sielkin.com)

Active ingredients (in each patch)  
Nicotine 14mg delivered over 24 hours

Stop smoking aid

reduces withdrawal symptoms, including nicotine cravings associated with quitting smoking.

Ask a doctor before use if you have

■heart disease ■high blood pressure ■skin allergies ■diabetes ■history of seizures

Ask a doctor or pharmacist before use if you are■using a non-nicotine stop smoking drug■pregnant or breast-feeding

When using this product

if you have vivid dreams or other sleep disturbances remove this patch at bedtime.

Stop use and ask a doctor if irritation or allergic reactions occur

Keep out of reach of children and pets. In case of accidental ingestion, seek medical advice immediately or contact a Poison Control Center(1-800-222-1222)

### Warnings

For external use only

### Directions

■if you are under 18 years of age, ask a doctor before use.

■Before use, carefully read the insert for complete instructions.

■If you smoke more than 10 cigarettes per day, follow this 10-week schedule:

	Step 1		Step 2	Step 3
Cigarettes per Day	Week 1(Key Adjustment Phase)	Weeks 2-6	Weeks 7-8	Weeks 9-10
10-20 cigarettes	two 21mg patch/day	one 21mg patch/day	one 14mg patch/day	one 7 mg patch/day
20+ cigarettes	three 21mg patch/day	two 21mg patch/day	one 14mg patch/day	one 7 mg patch/day

■If you smoke 10 or fewer cigarettes per day, skip STEP 1(21mg). Start with STEP 2(14mg) for 6 weeks-using two14mg patches per day for the first week, then one 14mg patch per day for weeks 2 to 6. Then move to STEP 3(7mg) for 2weeks, and stop.

Other information Store at 20-25°C (68-77°F)

Inactive ingredients camphor, ethanol, acrylate adhesive, aluminized polyester, cellulose paper, methacrylic acid copolymer

<b>SIELKIN QUIT SMOKING NICOTINE PATCHES 14MG</b>			
nicotine patch			
<b>Product Information</b>			
<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:83364-016
<b>Route of Administration</b>	TRANSDERMAL		
<b>Active Ingredient/Active Moiety</b>			
	<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
	<b>NICOTINE</b> (UNII: 6M3C89ZY6R) (NICOTINE - UNII:6M3C89ZY6R)	NICOTINE	14 mg in 1 g

**Inactive Ingredients**

Ingredient Name	Strength
<b>ACRYLIC ACID</b> (UNII: J94PBK7X8S)	
<b>METHACRYLIC ACID - METHYL METHACRYLATE COPOLYMER (1:1)</b> (UNII: 74G4R6TH13)	
<b>BORNEOL</b> (UNII: M89NIB437X)	
<b>ALCOHOL</b> (UNII: 3K9958V90M)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83364-016-01	30 in 1 BOX	01/01/2026	
1		0.014 g in 1 PATCH; Type 0: Not a Combination Product		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
unapproved drug other		01/01/2026	

**Labeler** - YITONGBADA (SHENZHEN) INTERNATIONAL TRADE CO., LTD (725220463)

Revised: 1/2026

YITONGBADA (SHENZHEN) INTERNATIONAL TRADE CO., LTD