ACETAMINOPHEN CONGESTION AND PAIN- acetaminophen, phenylephrine hcl tablet, film coated

Major Pharmaceuticals

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Major Pharmaceuticals Acetaminophen Congestion and Pain Drug Facts

Active ingredients (in each caplet)

Acetaminophen 325 mg

Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer

Nasal decongestant

Uses

- temporarily relieves these symptoms associated with hay fever or other upper respiratory allergies, and the common cold:
- headache
- sinus congestion and pressure
- nasal congestion
- minor aches and pains
- helps decongest sinus openings and passages
- promotes sinus drainage
- helps clear nasal passages
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland

Ask a doctor or pharmacist before use if you are

taking the blood thinning drug warfarin

When using this product

do not exceed recommended dosage

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur

These could be signs of a serious condition.

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

Overdose warning: In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

do not take more than directed (see overdose warning)

adults and children 12 years and over

- take 2 caplets every 4 hours
- swallow whole do not crush, chew or

	dissolvedo not take more than 10 caplets in 24 hours
children under 12	ask a doctor
years	

Inactive ingredients

acesulfame potassium, carnauba wax, crospovidone, D&C yellow #10 aluminum lake, FD&C blue #1 aluminum lake, FD&C red #40 aluminum lake, flavor, lecithin, microcrystalline cellulose, polyethylene glycol, polyvinyl alcohol, povidone, pregelatinized starch, stearic acid, talc, titanium dioxide

Questions or comments?

1-800-616-2471

Principal Display Panel

Compare to the active ingredients in Tylenol® SINUS + HEADACHE

Acetaminophen

CONGESTION AND PAIN

CAPLETS

Acetaminophen – Pain reliever / Fever reducer

Phenylephrine HCl – Nasal decongestant

Relieves: Sinus Headache, Sinus Pressure & Nasal Congestion

Actual Size

For Adults

24 CAPLETS

with cool Blast Flavor

NON-DROWSY

DAYTIME

Drug Facts (confinued)

Directions ■do not takemore than drected (see overdosewarring)

adults and children
■ swalow whole – do not cural, drew or dissolve

availar not take more than 10 capers in 24 hours

children under 12 years | sake a doctor

If pregnant or breast-feeding, sek a health professional before use Keep out of reach of children. Overdose war ning in case of overdose, get medical help or contact a Poison Combol Center right away (1-800-222-1222). Ouck medical attention is critical for adults as well as for children even if you do not not can signs or symptoms.

 new symptoms occur These could be signs of a serious condrion.
 If pregnant or breast-feeding, sek a health r Keep out of reach of children. Overdose way

- redness or swelling is present ■ new symptoms occur
- fever gets worse or lasts more than 3 days ■ pain or nasal congestion gets worse or lasts more than 7 days ■ nervousness, dezziness, or sleeplessness occur

Stop use and ask a doctor if

When using this product do not exceed recommended dosage

Ask a doctor or pharmacist before use if you are taking the blood

bring eurinating due or en larged prostate gland

Mark a doctor beforeuse if you have Marcisease markent dsease marigh blood pressure Marcia doctor beforeuse if you have

atre bengniat

If you are now taking a prescription monoamine oxidase inhibitor (MAM).) (certain drugs for depression, psychiatric, on emotional conditions, or Parkinson's desease), or for 2 weeks after stopping conditions, or Parkinson's desease), or for 2 weeks after stopping the MAOI drug. It you do not know it your prescription drug contains at MAOI drug. It you do not have an it you have ever had an allengic reaction to this product or any of it you have ever had an allengic reaction to this product or any of

Drug Facts (continued)

 with any other drug containing scelaminophen (prescription or nonprescription). If you are not sure whether a drug contains scelaminophen, ask a doctor or pharmacist.

Allergy alert Acetaminophen may cause sevene skin neactions. Symptoms may include: ■ skin reddening ■ blistens ■ rash If askin reaction occurs, stopuse and seek medical helpright away.

- 3 or more atocholic drinks every day while using this product
 - aruch 42 ninertonimateus to gm 000,4 north erom
 - damage may occur if you take

L**iver wamin g.** This product contains ace tamin opher. Severe liver Warnings

- temporarily reduces fever
- pelbe cjest usesj baseades
 bromotes sinne drainage
- helps decongest sinus openings and passages
 - anisq bns serbs ronim
 - nassal congestion
 - asinus congestion and pressure
 - headache
- The period of the set of the second of the common cold:

Pheny lephrine HCI 5 mg......

Dbhen 325 mg

(18

Pant na ieie ver\fever ne duc er Pain relie ver\fever ne duc er

Massal decongestant

Active ingredients (in each caplet)

Drug Facts

Important Read all product information before using Keep this box for important information.

*This productis no transmutac tured or d stributed by McNeil Consumer Healthc are, distributor of Tylerol ® SINUS + HEADACHE

DO NOTUSE IF BLISTER UNIT IS BROKEN ORTORN



MAJOR[®]

NDC 0904-6760-24 Compare to the active ingredients In Tylenol® SINUS + HEADACHE*

Acetaminophen congestion and pain

Acetaminophen - Pain reliever / Fever reducer Phenylephrine HCI - Nasal decongestant

Relieves: Sinus Headache, Sinus Pressure & Nasal Congestion

For Adults



CAPLETS

NON-DROWSY

DAYTIME





Store at 20-25°C (68-77°F)

Questions or comments?

-800-616-2471

NAJOR® PHARMAGEUTICALS
17177 N Laurel Park Drive, Suite 233
Livonia, MI 48152
M-05 REV, 04/19
Re-Order No. 700960

ACETAMINOPHEN CONGESTION AND PAIN

acetaminophen, phenylephrine hcl tablet, film coated

Product Information			
Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0904-6760
Route of Administration	ORAL		

Active Ingredient/Active Moiety		
Ingredient Name	Basis of Strength	Strength
ACETAMINO PHEN (UNII: 36209 ITL9 D) (ACETAMINO PHEN - UNII: 36209 ITL9 D)	ACETAMINOPHEN	325 mg
PHENYLEPHRINE HYDRO CHLO RIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII: 1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg

Inactive Ingredients		
Ingredient Name	Strength	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)		
CARNAUBA WAX (UNII: R12CBM0EIZ)		
CROSPOVIDONE (15 MPA.S AT 5%) (UNII: 68401960MK)		
MICRO CRYSTALLINE CELLULO SE (UNII: OP1R32D61U)		
POLYETHYLENE GLYCOL (UNII: 3WJQ0SDW1A)		
POLYVINYL ALCOHOL (UNII: 532B59J990)		
PO VIDO NE (UNII: FZ989 GH94E)		
STEARIC ACID (UNII: 4ELV7Z65AP)		
TALC (UNII: 7SEV7J4R1U)		
TITANIUM DIO XIDE (UNII: 15FIX9 V2JP)		
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)		
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)		
FD&C RED NO. 40 (UNII: WZB9127XOA)		

Product Characteristics			
Color	GREEN	Score	no score
Shape	OVAL	Size	16 mm
Flavor	VANILLA (menthol)	Imprint Code	L272
Contains			

	Packaging				
Ш	# Item Code	Package Description	Marketing Start Date	Marketing End Date	
	NDC:0904-6760-24	12 in 1 CARTON	01/15/2007		
	1	2 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Marketing Information			
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date

OTC monograph final part341 01/15/2007

Labeler - Major Pharmaceuticals (191427277)

Revised: 12/2019 Major Pharmaceuticals