

**QUALITY CHOICE FIBER CAPLETS- calcium polycarbophil tablet, film coated
QUALITY CHOICE (CHAIN DRUG MARKETING ASSOCIATION)**

Quality Choice Fiber Caplets Calcium Polycarbophil

Drug Facts

Active Ingredient (in each caplet)

Calcium polycarbophil 625 mg equivalent to 500 mg polycarbophil

Purpose

Bulk-forming laxative

U ses

- relieves occasional constipation to help restore and maintain regularity
- this product generally produces bowel movement in 12 to 72 hours

Choking Warning: Taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have difficulty in swallowing. If you experience chest pain, vomiting, or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.

Ask your doctor before use if you have

- abdominal pain, nausea, or vomiting
- a sudden change in bowel habits that persists over a period of 2 weeks

Ask your doctor or pharmacist before use if you are

taking any other drug. Take this product 2 or more hours before or after other drugs. All laxatives may affect how other drugs work.

When using this product

- do not use for more than 7 days unless directed by a doctor
- do not take more than 8 caplets in a 24 hour period unless directed by a doctor

Stop use and ask a doctor if

rectal bleeding occurs or if you fail to have a bowel movement after use of this or any other laxative. These could be signs of a serious condition.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away at 1-800-222-1222.

Directions

- Take each dose of this product with at least 8 ounces (a full glass) of water or other

fluid. Taking this product without enough liquid may cause choking. (See choking warning).

- This product works naturally so continued use for one to three days is normally required to provide full benefit. Dosage may vary according to diet, exercise, previous laxative use or severity of constipation.

age	Recommended	Daily maximum
adults & children 12 years of age and over	2 caplets once a day	up to 4 times a day
children under 12 years	consult a physician	

Other information

- **each caplet contains:** calcium 125 mg
- store at room temperature
- protect contents from moisture

calcium carbonate, caramel, crospovidone, hypromellose, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium lauryl sulfate.

Questions or Comments?

1-866-467-2748

PRINCIPAL DISPLAY PANEL

NDC# **83324-276-90**

QUALITY CHOICE ®

***Compare to the Active ingredients in Fibercon ®**

Fiber Caplets

Calcium Polycarbophil 625 mg

Bulk-Forming

Fiber Laxative

Won't Cause Gas or Bloating As Effective as Fiber Powders

90 Caplets

100% QC SATISFACTION GURANTEED

Distributed by C.D.M.A., Inc.

Novi, MI 48375

www.qualitychoice.com

Questions: 800-935-2362

DO NOT USE IF PRINTED INNER SEAL UNDER CAP IS TORN OR MISSING

*This product is not manufactured or distributed by Pfizer Inc., owner of the registered trademark Fibercon® Caplets.

Package label for 90 Caplets

NDC 83324-276-90

QC
QUALITY CHOICE

*Compare to the Active Ingredient in FiberCon®

Fiber Caplets

Calcium Polycarbophil 625 mg

Bulk-Forming Fiber Laxative
Gentle Relief of Constipation
Won't Cause Gas or Bloating
As Effective as Fiber Powders

90 Caplets

DO NOT USE IF PRINTED INNER SEAL UNDER CAP IS TORN OR MISSING

Drug Facts

Active ingredient (in each caplet) Purpose
Calcium polycarbophil 625 mg Bulk-forming laxative equivalent to 500 mg polycarbophil.....

Uses ■ relieves occasional constipation to help restore and maintain regularity ■ this product generally produces bowel movement in 12 to 72 hours.

Warnings
Choking: Taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have difficulty in swallowing, if you experience chest pain, vomiting or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.
Ask your doctor before use if you have ■ abdominal pain, nausea or vomiting ■ a sudden change in bowel habits that persists over a period of 2 weeks.

Ask your doctor or pharmacist before use if you are taking any other drug. Take this product 2 or more hours before or after other drugs. All laxatives may affect how other drugs work.

When using this product ■ do not use for more than 7 days unless directed by a doctor ■ do not take more than 8 caplets in a 24 hour period unless directed by a doctor

Stop use and ask a doctor if rectal bleeding occurs or if you fail to have a bowel movement after use of this or any other laxative. These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away, 1-800-222-1222.

*This product is not manufactured or distributed by Pfizer Inc., owner of the registered trademark FiberCon® Caplets.

Distributed by CDMA, Inc.
Novi, MI 48275
www.cdma.com
Customer Care: 1-800-955-2942

LOT: 120900CLR

EXP:

35515190023

PEEL BACK HERE

Drug Facts (continued)

Directions ■ take each dose of this product with at least 8 ounces (a full glass) of water or other fluid. Taking this product without enough liquid may cause choking. See choking warning. ■ this product works naturally so continued use for one to three days is normally required to provide full benefit. Dosage may vary according to diet, exercise, previous laxative use or severity of constipation

age	recommended dose	daily maximum
adults & children 12 years of age and over	2 caplets once a day	up to 4 times a day
children under 12 years	consult a physician	

Other information ■ each caplet contains: calcium 125 mg ■ store at room temperature ■ protect contents from moisture

Inactive ingredients calcium carbonate, caramel color, croscovidone, hypromellose, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium lauryl sulfate

Questions or comments? 1-866-467-2748

STOP PEELING HERE

Package Label for 140 Caplets

NDC 83324-276-14

QC
QUALITY CHOICE

*Compare to the Active Ingredient in FiberCon® Caplets

Fiber Caplets

Calcium Polycarbophil 625 mg

Bulk-Forming Fiber Laxative
Gentle Relief of Constipation
Won't Cause Gas or Bloating
As Effective as Fiber Powders

140 Caplets

Drug Facts		Drug Facts (continued)										
Active ingredient (in each caplet) Calcium polycarbophil 625 mg equivalent to 500 mg polycarbophil.....Bulk-forming laxative	PurposeBulk-forming laxative	These could be signs of a serious condition. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away 1-800-222-1222										
Uses ■ relieves occasional constipation to help restore and maintain regularity ■ this product generally produces bowel movement in 12 to 72 hours.		Directions ■ take each dose of this product with at least 8 ounces (a full glass) of water or other fluid. Taking this product without enough liquid may cause choking. See choking warning. ■ this product works naturally so continued use for one to three days is normally required to provide full benefit. Dosage may vary according to diet, exercise, previous laxative use or severity of constipation										
Warnings Choking: Taking this product without adequate fluid may cause it to swell and block your throat or esophagus and may cause choking. Do not take this product if you have difficulty in swallowing. If you experience chest pain, vomiting or difficulty in swallowing or breathing after taking this product, seek immediate medical attention.		<table border="1"> <thead> <tr> <th>age</th> <th>recommended dose</th> <th>daily maximum</th> </tr> </thead> <tbody> <tr> <td>adults & children 12 years of age and over</td> <td>2 caplets once a day</td> <td>up to 4 times a day</td> </tr> <tr> <td>children under 12 years</td> <td colspan="2">consult a physician</td> </tr> </tbody> </table>		age	recommended dose	daily maximum	adults & children 12 years of age and over	2 caplets once a day	up to 4 times a day	children under 12 years	consult a physician	
age	recommended dose	daily maximum										
adults & children 12 years of age and over	2 caplets once a day	up to 4 times a day										
children under 12 years	consult a physician											
Ask your doctor before use if you have ■ abdominal pain, nausea or vomiting ■ a sudden change in bowel habits that persists over a period of 2 weeks.		Other information ■ each caplet contains: calcium 125 mg ■ store at room temperature ■ protect contents from moisture										
Ask your doctor or pharmacist before use if you are taking any other drug. Take this product 2 or more hours before or after other drugs. All laxatives may affect how other drugs work.		Inactive ingredients calcium carbonate, caramel color, crospovidone, hypromellose, magnesium stearate, maltodextrin, microcrystalline cellulose, polyethylene glycol, silicon dioxide, sodium lauryl sulfate										
When using this product ■ do not use for more than 7 days unless directed by a doctor ■ do not take more than 8 caplets in a 24 hour period unless directed by a doctor		Questions or comments? 1-866-467-2748										
Stop use and ask a doctor if rectal bleeding occurs or if you fail to have a bowel movement after use of this or any other laxative.												

This product is not manufactured or distributed by Pfizer Inc., owner of the registered trademark FiberCon® Caplets.
Distributed by CDMA, Inc.
www.qualitychoice.com
Questions: 800-935-2882

LOT: 1201400CLR
EXP: 3551519998 5

QUALITY CHOICE FIBER CAPLETS

calcium polycarbophil tablet, film coated

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83324-276
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
CALCIUM POLYCARBOPHIL (UNII: 8F049NKY49) (POLYCARBOPHIL - UNII: W25LM17A4W)	CALCIUM POLYCARBOPHIL	625 mg

Inactive Ingredients

Ingredient Name	Strength
CALCIUM CARBONATE (UNII: H0G9379FGK)	
CARAMEL (UNII: T9D99G2B1R)	
CROSPVIDONE (120 .MU.M) (UNII: 68401960MK)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
MALTODEXTRIN (UNII: 7CVR7L4A2D)	
MAGNESIUM STEARATE (UNII: 70097M6130)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	

Product Characteristics

Color	brown	Score	no score
Shape	OVAL	Size	19mm
Flavor		Imprint Code	RP120

Contains**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83324-276-90	90 in 1 BOTTLE; Type 0: Not a Combination Product	06/02/2025	
2	NDC:83324-276-14	140 in 1 BOTTLE; Type 0: Not a Combination Product	06/02/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M007	06/02/2025	

Labeler - QUALITY CHOICE (CHAIN DRUG MARKETING ASSOCIATION) (011920774)

Revised: 7/2025

QUALITY CHOICE (CHAIN DRUG MARKETING ASSOCIATION)