

**ELIZABETH ARDEN PREVAGE ANTI-AGING FOUNDATION BROAD SPECTRUM
SUNSCREEN SPF 30- octinoxate, titanium dioxide lotion
ELIZABETH ARDEN, INC**

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

Drug Facts

Active Ingredients

OCTINOXATE 6.0%

TITANIUM DIOXIDE 6.0%

Purpose

Sunscreen

Sunscreen

Uses

- Helps prevent sunburn
- If used as directed with other sun protection measures (see **Directions**), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

- **For external use only**
- **Do not use** on damaged or broken skin
- **When using this product** keep out of eyes. Rinse with water to remove.
- **Stop use and ask a doctor** if rash occurs
- **Keep out of reach of children.** If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions For Sunscreen Use:

- Apply liberally 15 minutes before sun exposure
- Reapply:
 - at least every 2 hours
 - use a water resistant sunscreen if swimming or sweating
- **Sun Protection Measures.**
Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:
 - Limit time in the sun, especially from 10 a.m. – 2 p.m.
 - Wear long-sleeve shirts, pants, hats, and sunglasses
- Children under 6 months: Ask a doctor.

Inactive Ingredients

WATER/AQUA/EAU, CYCLOPENTASILOXANE, DICAPRYLYL CARBONATE, MYRISTYL TRISILOXANE, POLYMETHYLSILSESQUIOXANE, CYCLOHEXASILOXANE, SORBITAN ISOSTEARATE, ALUMINA, ALUMINUM/MAGNESIUM HYDROXIDE STEARATE, GLYCERIN, TETRAHEXYLDECYL ASCORBATE, TRIMETHYLSILOXYSILICATE, MICA, CAPRYLYL DIMETHICONE ETHOXY GLUCOSIDE, CAMELLIA SINENSIS LEAF EXTRACT, CARNOSINE, DIMETHICONE, DIMETHICONOL, DISODIUM COCOYL GLUTAMATE, ERGOTHIONEINE, FERULIC ACID, HYDROXYDECYL UBIQUINOYL DIPALMITOYL GLYCERATE, PARFUM/FRAGRANCE, PCA DIMETHICONE, POLYHYDROXYSTEARIC ACID, SILICA, SILICA Silylate, SODIUM CHLORIDE, SODIUM COCOYL GLUTAMATE, STEARIC ACID, TOCOPHEROL, CHLORPHENESIN, PHENOXYETHANOL, POTASSIUM SORBATE, SODIUM DEHYDROACETATE, IRON OXIDES (CI 77491, CI 77492, CI 77499), TITANIUM DIOXIDE (CI 77891).

Other Information

- Protect this product from excessive heat and direct sun

Questions or comments?

- Call toll free 1-800 326-7337

Principal Display Panel - PreVage Anti-Aging Foundation Carton Label

prevage[®]

anti-aging foundation

broad spectrum sunscreen

SPF 30

1 FL. OZ. 30 ml

- For normal to dry skin
- Light to medium coverage.
- Naturally luminous finish.

The power of PREVAGE® now in an anti-aging foundation. Our unique Idebenone complex protects skin from environmental assaults to intercept visible aging signs. Vitamin C and key anti-aging ingredients correct skin's appearance by supporting natural collagen and elastin for a firmer, clearer look. Moisture-rich hydra-pigments flawlessly even skin tone for luminous skin perfection that lasts all day. Oil free.

To use: Smooth onto face.
See insert or elizabetharden.com for complete directions prior to first use.

Drug Facts

Active Ingredients	Purpose
OCTINOXATE 6.0%	Sunscreen
TITANIUM DIOXIDE 6.0%	Sunscreen

Uses

- Helps prevent sunburn
- If used as directed with other sun protection measures (see **Directions**), decreases the risk of skin cancer and early skin aging caused by the sun

Warnings

- For external use only
- Do not use on damaged or broken skin
- When using this product keep out of eyes. Rinse with water to remove.
- Stop use and ask a doctor if rash occurs
- Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.

Directions For Sunscreen Use:

- Apply liberally 15 minutes before sun exposure
- Reapply:
 - at least every 2 hours
 - use a water resistant sunscreen if swimming or sweating
- Sun Protection Measures. ▶

Drug Facts (continued)

Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF of 15 or higher and other sun protection measures including:

- Limit time in the sun, especially from 10 a.m. – 2 p.m.
- Wear long-sleeve shirts, pants, hats, and sunglasses
- Children under 6 months: Ask a doctor.

Inactive Ingredients

WATER/AQUA/EAU, CYCLOPENTASILOXANE, DICAPRYLYL CARBONATE, MYRISTYL TRISILOXANE, POLYMETHYLSILSESQUOXANE, CYCLOHEXASILOXANE, SORBITAN ISOSTEARATE, ALUMINA, ALUMINUM/MAGNESIUM HYDROXIDE STEARATE, GLYCERIN, TETRAHEXYLDECYL ASCORBATE, TRIMETHYLSILOXYSILICATE, MICA, CAPRYLYL DIMETHICONE ETHOXY GLUCOSIDE, CAMELLIA SINENSIS LEAF EXTRACT, CARNOSINE, DIMETHICONE, DIMETHICONOL, DISODIUM COCOYL GLUTAMATE, ERGOTHIONEINE, FERULIC ACID, ▶

prevage®

Drug Facts (continued)

HYDROXYDECYL UBIQUINOYL DIPALMITOYL GLYCERATE, PARFUM/FRAGRANCE, PCA DIMETHICONE, POLYHYDROXYSTEARIC ACID, SILICA, SILICA Silylate, SODIUM CHLORIDE, SODIUM COCOYL GLUTAMATE, STEARIC ACID, TOCOPHEROL, CHLORPHENESIN, PHENOXYETHANOL, POTASSIUM SORBATE, SODIUM DEHYDROACETATE, IRON OXIDES (CI 77491, CI 77492, CI 77499), TITANIUM DIOXIDE (CI 77891).

Other Information

- Protect this product from excessive heat and direct sun

Questions or comments?

- Call toll free 1-800 326-7337


anti-aging foundation
broad spectrum sunscreen
SPF 30

1 FL. OZ. 30 ml

Distributed in U.S.A. by
© Elizabeth Arden, Inc.,
New York, NY 10003
Made in U.S.A.
elizabetharden.com
U.S. Patent 6,756,045 and other pending/
issued U.S. and international patents.

266320





ELIZABETH ARDEN PREVAGE ANTI-AGING FOUNDATION BROAD SPECTRUM SUNSCREEN SPF 30

octinoxate, titanium dioxide lotion

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:67938-3051
Route of Administration	TOPICAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
OCTINOXATE (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	.06 g in 1 mL
TITANIUM DIOXIDE (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	.06 g in 1 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	
CYCLOMETHICONE 5 (UNII: 0TH5PC10R)	
DICAPRYLYL CARBONATE (UNII: 609A3V1SUA)	
MYRISTYL TRISILOXANE (UNII: J7960S4R1T)	
CYCLOMETHICONE 6 (UNII: XHK3U310BA)	
SORBITAN ISOSTEARATE (UNII: 01S2G2C1E4)	
ALUMINUM OXIDE (UNII: LM26O6933)	
GLYCERIN (UNII: PDC6A3C0OX)	
TETRAHEXYLDECYL ASCORBATE (UNII: 9LBV3F07AZ)	
MICA (UNII: V8A1AW0880)	
GREEN TEA LEAF (UNII: W2ZU1RY8B0)	
CARNOSINE (UNII: 8HO6PVN24W)	
DIMETHICONE (UNII: 92RU3N3Y1O)	
DISODIUM COCOYL GLUTAMATE (UNII: MBK0CP8F5A)	
ERGOTHIONEINE (UNII: BDZ3DQM98W)	
FERULIC ACID (UNII: AVM951ZWST)	
HYDROXYDECYL UBIQUINOYL DIPALMITOYL GLYCERATE (UNII: OV1BT2N8RC)	
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)	
SODIUM CHLORIDE (UNII: 451W47IQ8X)	
SODIUM COCOYL GLUTAMATE (UNII: BMT4RCZ3HG)	
STEARIC ACID (UNII: 4ELV7Z65AP)	

TOCOPHEROL (UNII: R0ZB2556P8)	
CHLORPHENESIN (UNII: I670DAL4SZ)	
PHENOXYETHANOL (UNII: HE492ZZ3T)	
POTASSIUM SORBATE (UNII: 1VPU26JZZ4)	
SODIUM DEHYDRO ACETATE (UNII: 8W46YN971G)	
FERRIC OXIDE RED (UNII: 1K09F3G675)	
FERRIC OXIDE YELLOW (UNII: EX438O2MRT)	
FERROSO FERRIC OXIDE (UNII: XM0M87F357)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:67938-3051-1	1 in 1 CARTON	08/01/2017	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part352	08/01/2017	

Labeler - ELIZABETH ARDEN, INC (849222187)

Revised: 8/2017

ELIZABETH ARDEN, INC