

**SUN PROTECTION SPF 50- avobenzene 2.9%, homosalate 4.0%, octisalate 4.9%, octocrylene 9.5% spray**  
**Skinresource.md, L.L.C.**

-----

## ***Drug Facts***

### ***Active Ingredient***

Avobenzene – 2.9% - Homosalate – 4.0% - Octisalate – 4.9% - Octocrylene – 9.5%

Active Ingredient

Avobenzene - 2.9% - Homosalate - 4.0% - Octisalate - 4.9% - Octocrylene - 9.5%

***Inactive Ingredients:*** Acrylates/Octylacrylamide Copolymer, Alcohol Denat, Aloe Barbadensis Leaf Extract, Butyloctyl Salicylate, Carthamus Tinctorius (Safflower) Seed Oil, Ethyl Ferulate, Helianthus Annuus (Sunflower) Seed Oil\*, Polyester-8, Tocopherol. \* Indicates Certified Organic Ingredient.

Inactive Ingredients:

Acrylates/Octylacrylamide Copolymer, Alcohol Denat, Aloe Barbadensis Leaf Extract, Butyloctyl Salicylate, Carthamus Tinctorius (Safflower) Seed Oil, Ethyl Ferulate, Helianthus Annuus (Sunflower)

Seed Oil\*, Polyester-8, Tocopherol.

\* Indicates Certified Organic Ingredient.

**Keep out of reach of children**

■ If product is swallowed, get medical help or contact a Poison Control Center right away

Keep out of reach of children:

if product is swallowed, get medical help or contact a Poison Control Center right away

## Warnings

■ For external use only ■ Flammable Do not spray near heat, sparks, sources of ignition, or flames. Protect this product from excessive heat and direct sun

*Do not use* ■ on damaged or broken skin

*When using this product* ■ keep out of eyes ■ Rinse with water to remove.

Stop use and ask a doctor if skin irritation or redness occurs for more than 72 hours.

### Warnings

For external use only Flammable Do not spray near heat, sparks, sources of ignition, or flames.

Protect this product from excessive heat and direct sun

Do not use on damaged or broken skin

When using this product keep out of eyes Rinse with water to remove.

Stop use and ask a doctor if skin irritation or redness occurs for more than 72 hours.

AVOBENZONE - 2.9% - HOMOSALATE - 4.0% - OCTISALATE - 4.9% - OCTOXYLENE - 9.5%

## Purpose Sunscreen

**Directions** ■ Apply liberally 15 minutes before sun exposure. **Reapply:** Every 2 hours during the day or after swimming or sweating. • Children under 6 months: ask a doctor.

**Sun Protection Measure:** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF value of 15 or higher and other sun protection measures including: Limit time in the sun, especially from 10 a.m. to 2 p.m. • Wear long-sleeved shirts, pants, hats, and sun glasses.

Directions • Apply liberally 15 minutes before sun exposure. Reapply: Every 2 hours during the day or after swimming or sweating. • Children under 6 months: ask a doctor.

Sun Protection Measure: Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF value of 15 or higher and other sun protection measures including: Limit time in the sun, especially from 10 a.m.

to 2 p.m. • Wear long-sleeved shirts, pants, hats, and sun glasses.

**Uses** • Helps prevent sunburn • If used as directed with other sun protection measures, decreases the risk of skin cancer and early skin aging caused by the sun.

## **Warnings**

■ **For external use only** ■ **Flammable** Do not spray near heat, sparks, sources of ignition, or flames. Protect this product from excessive heat and direct sun

**Do not use** ■ on damaged or broken skin

**When using this product** ■ keep out of eyes ■ Rinse with water to remove.

**Stop use and ask a doctor** if skin irritation or redness occurs for more than 72 hours.

## **Keep out of reach of children**

■ If product is swallowed, get medical help or contact a Poison Control Center right away

**Directions** ■ Apply liberally 15 minutes before sun exposure. **Reapply:** Every 2 hours during the day or after swimming or sweating. • Children under 6 months: ask a doctor.

**Sun Protection Measure:** Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF value of 15 or higher and other sun protection measures including: Limit time in the sun, especially from 10 a.m. to 2 p.m. • Wear long-sleeved shirts, pants, hats, and sun glasses.

Uses • Helps prevent sunburn • If used as directed with other sun protection measures, decreases the risk of skin cancer and early skin aging caused by the sun.

## Warnings

• For external use only Flammable Do not spray near heat, sparks, sources of ignition, or flames.

Protect this product from excessive heat and direct sun

Do not use on damaged or broken skin

When using this product keep out of eyes • Rinse with water to remove.

Stop use and ask a doctor if skin irritation or redness occurs for more than 72 hours.

Keep out of reach of children

If product is swallowed, get medical help or contact a Poison Control Center right away

Directions • Apply liberally 15 minutes before sun exposure. Reapply: Every 2 hours during the day after swimming or sweating. • Children under 6 months: ask a doctor.

Sun Protection Measure: Spending time in the sun increases your risk of skin cancer and

early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF value of 15 or higher and other sun protection measures including: Limit time in the sun, especially from 10 a.m.

to 2 p.m. • Wear long-sleeved shirts, pants, hats, and sun glasses.



## Sun Protection Spray SPF 50

Prescribe it for Yourself.®

 MD



6 Fl. Oz. (177 mL)

<b>Drug Facts</b>
<b>Active Ingredient</b> Avobenzone – 2.9% - Homosalate – 4.0% - Octisalate – 4.9% - Octocrylene – 9.5%
<b>Purpose</b> Sunscreen
<b>Uses</b> • Helps prevent sunburn • If used as directed with other sun protection measures, decreases the risk of skin cancer and early skin aging caused by the sun.
<b>Warnings</b> ■ <b>For external use only</b> ■ <b>Flammable</b> Do not spray near heat, sparks, sources of ignition, or flames. Protect this product from excessive heat and direct sun
<b>Do not use</b> ■ on damaged or broken skin
<b>When using this product</b> ■ keep out of eyes ■ Rinse with water to remove.
<b>Stop use and ask a doctor</b> if skin irritation or redness occurs for more than 72 hours.
<b>Keep out of reach of children</b> ■ If product is swallowed, get medical help or contact a Poison Control Center right away
<b>Directions</b> ■ Apply liberally 15 minutes before sun exposure. <b>Reapply:</b> Every 2 hours during the day or after swimming or sweating. • Children under 6 months: ask a doctor.
<b>Sun Protection Measure:</b> Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a broad spectrum SPF value of 15 or higher and other sun protection measures including: Limit time in the sun, especially from 10 a.m. to 2 p.m. • Wear long-sleeved shirts, pants, hats, and sun glasses.
<b>Inactive Ingredients:</b> Acrylates/Octylacrylamide Copolymer, Alcohol Denat, Aloe Barbadensis Leaf Extract, Butyloctyl Salicylate, Carthamus Tinctorius (Safflower) Seed Oil, Ethyl Ferulate, Helianthus Annuus (Sunflower) Seed Oil*, Polyester-8, Tocopherol. * Indicates Certified Organic Ingredient.
<b>Questions?</b> call (619) 776-3464 Option 3
Skin Resource.MD • 4250 Pacific Hwy, Suite 128, San Diego CA 92110 © Skin Resource.MD All rights reserved • Made in United States

 **NO ANIMAL TESTING** Distributed by Skin Resource.MD LLC, San Diego CA 92110 • SkinResourceMD.com

## SUN PROTECTION SPF 50

avobenzone 2.9%, homosalate 4.0%, octisalate 4.9%, octocrylene 9.5% spray

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:83694-011
<b>Route of Administration</b>	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>HOMOSALATE</b> (UNII: V06SV4M95S) (HOMOSALATE - UNII:V06SV4M95S)	HOMOSALATE	7.08 g in 177 g
<b>OCTOCRYLENE</b> (UNII: 5A68WGF6WM) (OCTOCRYLENE - UNII:5A68WGF6WM)	OCTOCRYLENE	16.815 g in 177 g
<b>OCTISALATE</b> (UNII: 4X49Y0596W) (OCTISALATE - UNII:4X49Y0596W)	OCTISALATE	8.673 g in 177 g
<b>AVOBENZONE</b> (UNII: G63QQF2NOX) (AVOBENZONE - UNII:G63QQF2NOX)	AVOBENZONE	5.133 g in 177 g

## Inactive Ingredients

Ingredient Name	Strength
<b>ETHYL FERULATE</b> (UNII: 5B8915UELW)	
<b>ALOE BARBADENSIS LEAF</b> (UNII: ZY81Z83H0X)	
<b>DEHYDRATED ALCOHOL</b> (UNII: 3K9958V90M)	
<b>POLYESTER-8 (1400 MW, CYANODIPHENYLPROPENOYL CAPPED)</b> (UNII: T9296U138P)	
<b>TOCOPHEROL</b> (UNII: R0ZB2556P8)	
<b>ACRYLATES/OCTYLACRYLAMIDE COPOLYMER (75000 MW)</b> (UNII: JU3XHR8VWK)	
<b>BUTYLOCTYL SALICYLATE</b> (UNII: 2EH13UN8D3)	
<b>CARTHAMUS TINCTORIUS (SAFFLOWER) SEED OIL</b> (UNII: 65UEH262IS)	
<b>HELIANTHUS ANNUUS (SUNFLOWER) SEED OIL</b> (UNII: 3W1JG795YI)	

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83694-011-01	177 g in 1 BOTTLE, SPRAY; Type 0: Not a Combination Product	03/01/2025	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M020	03/01/2025	

**Labeler** - Skinresource.md, L.L.C. (022050636)

Revised: 3/2025

Skinresource.md, L.L.C.