

**OHUI POWDERY METAL CUSHION COMPACT 02 NATURAL BEIGE- titanium dioxide powder**

**LG Household and Health Care Inc**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**ACTIVE INGREDIENTS**

ENSULIZOLE 2.7%, OCTINOXATE 7%, TITANIUM DIOXIDE 7.18%, ZINC OXIDE 2.91%

**PURPOSE**

Sunscreen

**USES**

- Helps prevent sunburn.
- If used as directed with other sun protection measures (see Directions),

decreases the risk of skin cancer and early skin aging caused by the sun.

**WARNINGS**

**For external use only**

**Do not use** on damaged or broken skin.

**When using this product:** Kepp out of the eyes. Rinse with water to remove.

**Stop use and ask doctor**, if rash occurs.

**Keep out of reach of children.** If product is swallowed, get medical help or contact a Posion Control Center right away.

**DIRECTIONS**

- Apply liberally 15 minutes before sun exposure.
- Use a water resistant sunscreen if swimming or sweating.
- Reapply at least every 2 hours.
- **Sun Protection Measures.** Spending time in the sun increases your risk

Of skin cancer and early skin aging. To decrease this risk, regularly use a Sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:

- Limit time in the sun, especially from 10 a.m. to 2 p.m.
- Wear longsleeved shirts, pants, hats, and sunglasses.
- Children under 6 months: Ask a doctor.

**OTHER INFORMATION**


- Protect this product from excessive heat and direct sun.

**INACTIVE INGREDIENTS**

CYCLOPENTASILOXANE, WATER, METHYL TRIMETHICONE, DIPROPYLENE GLYCOL, PEG-10 DIMETHICONE, POLYMETHYL METHACRYLATE, NIACINAMIDE, TROMETHAMINE, 1,2-HEXANEDIOL, DISTEARDIMONIUM HECTORITE, STEARIC ACID, ALUMINUM HYDROXIDE, DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER, SODIUM CHLORIDE, SORBITAN SESQUIOLEATE, SORBITAN SESQUIISOSTEARATE, ZINC STEARATE, DISTEARYLDIMONIUM CHLORIDE, TOCOPHERYL ACETATE, PALMITOYL PROLINE, DIMETHICONE/METHICONE COPOLYMER, ADENOSINE, MAGNESIUM PALMITOYL GLUTAMATE, SODIUM PALMITOYL SARCO SINATE, PHENOXYETHANOL, TRIETHOXYCAPRYLYLSILANE, TRIETHOXSILYLETHYL POLYDIMETHYLSILOXYETHYL HEXYL DIMETHICONE, BIOSACCHARIDE GUM-1, PALMITIC ACID, BUTYLENE GLYCOL, CETYL PEG/PPG-10/1 DIMETHICONE, HEXYL LAURATE, POLYGLYCERYL-4 ISOSTEARATE, ALBIZIA JULIBRISSIN BARK EXTRACT, FRAGRANCE, TITANIUM DIOXIDE(CI 77891), MICA, IRON OXIDES(CI 77492), IRON OXIDES(CI 77491), IRON OXIDES(CI 77499)

**PRINCIPAL DISPLAY PANEL**

OHUI Powdery Metal Cushion Compact  
 Sunscreen Broad Spectrum SPF 50  
 02 Natural Beige  
 Net Wt. 1.05 oz / 30 g (15gx2)

		<p><b>POWDERY METAL CUSHION COMPACT</b>                  COUSSIN POUR LE TEINT MÉTAL                  Broad Spectrum SPF 50</p> <p><b>02</b>                  Natural Beige                  Beige Naturel</p>																																															
<p>O HUI  <b>POWDERY METAL CUSHION COMPACT</b>                  Broad Spectrum SPF 50</p> <p>This cushion foundation helps to keep a long-lasting coverage with liquid-to-powder formula, leaving skin flawless and silky smooth.  <b>How to Use:</b> Place the puff on the metal plate and press down to pump once or twice. Tap gently to spread evenly over the face.</p> <p>Ce fond de teint en coussin promet une couverture de longue durée. Avec sa poudre liquide il laisse la peau douce et soyeuse.                  Utilisation: placez la houppette sur la plaque de métal et pompez une ou deux fois. Tapotez délicatement pour répandre sur le visage.</p> <p>Imported by LG Household &amp; Health Care America, Inc. Santa Fe Springs, CA 90670 USA                  Questions: 1-888-498-9004                  MADE IN KOREA</p>	<p><b>Drug Fact (Continued)</b></p> <p><b>Other Information</b>                  ■ Protect this product from excessive heat and direct sun.</p> <p><b>Inactive Ingredients</b>                  Cyclopentasiloxane, Water, Methyl Trimethicone, Dipropylene Glycol, PEG-10 Dimethicone, Polymethyl Methacrylate, Niacinamide, Tromethamine, 1,2-Hexanediol, Distearidimonium Hectorite, Stearic Acid, Aluminum Hydroxide, Dimethiconal/Vinyl Dimethicone Crosspolymer, Sodium Chloride, Sorbitan Sesquioxide, Sorbitan Sesquiossearate, Zinc Stearate, Distearidimonium Chloride, Tocopheryl Acetate, Palmitoyl Proline, Dimethiconal/Methicone Copolymer, Adenosine, Magnesium Palmityl Glutamate, Sodium Palmitoyl Sarcosinate, Phenoxyethanol, Triethoxycaprylylsilane, Triethoxymethyl Polydimethylsilyloxyethyl Hexyl Dimethicone, Biosaccharide Gum-1, Palmitic Acid, Butylene Glycol, Cetyl PEG/PPG-10/1 Dimethicone, Hexyl Laurate, Polyglyceryl-4 Isostearate, Albizia Julibrissin Bark Extract, Fragrance, Titanium dioxide(CI 77891), Mica, Iron Oxides(CI 77491, CI 77492, CI 77499)</p> <p><b>Questions?</b>                  Call 1-888-498-9004 (Monday-Friday 9 a.m. - 6 p.m.)</p>	<p>O HUI</p> <p><b>POWDERY METAL CUSHION COMPACT</b>                  COUSSIN POUR LE TEINT MÉTAL</p> <p>Sunscreen                  Broad Spectrum SPF 50</p> <p>Net Wt. 1.05 oz / 30 g (15gx2)</p>	<table border="1"> <tr> <th colspan="2">Drug Facts</th> </tr> <tr> <th>Active Ingredients</th> <th>Purpose</th> </tr> <tr> <td>Erythrolol 2.7%</td> <td>Sunscreen</td> </tr> <tr> <td>Octinoxide 7%</td> <td>Sunscreen</td> </tr> <tr> <td>Titanium dioxide 7.16%</td> <td>Sunscreen</td> </tr> <tr> <td>Zinc Oxide 2.91%</td> <td>Sunscreen</td> </tr> <tr> <td colspan="2"><b>Uses</b></td> </tr> <tr> <td colspan="2">■ Helps prevent sunburn.</td> </tr> <tr> <td colspan="2">■ If used as directed with other sun protection measures (see Directions), may decrease the risk of skin cancer and early skin aging caused by the sun.</td> </tr> <tr> <td colspan="2"><b>Warnings</b></td> </tr> <tr> <td colspan="2">For external use only</td> </tr> <tr> <td colspan="2">Do not use on damaged or broken skin.</td> </tr> <tr> <td colspan="2">When using this product: Keep out of the eyes. Rinse with water to remove.</td> </tr> <tr> <td colspan="2">■ Stop use and ask a doctor, if rash occurs.</td> </tr> <tr> <td colspan="2">Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.</td> </tr> <tr> <td colspan="2"><b>Directions</b></td> </tr> <tr> <td colspan="2">■ Apply liberally 15 minutes before sun exposure.</td> </tr> <tr> <td colspan="2">■ Use a water resistant sunscreen if swimming or sweating.</td> </tr> <tr> <td colspan="2">■ Reapply at least every 2 hours.</td> </tr> <tr> <td colspan="2">■ Sun Protection Measures: Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:</td> </tr> <tr> <td colspan="2">• Limit time in the sun, especially from 10 a.m. to 2 p.m.</td> </tr> <tr> <td colspan="2">• Wear long-sleeved shirts, pants, hats, and sunglasses.</td> </tr> <tr> <td colspan="2">• Children under 6 months: Ask a doctor.</td> </tr> </table>	Drug Facts		Active Ingredients	Purpose	Erythrolol 2.7%	Sunscreen	Octinoxide 7%	Sunscreen	Titanium dioxide 7.16%	Sunscreen	Zinc Oxide 2.91%	Sunscreen	<b>Uses</b>		■ Helps prevent sunburn.		■ If used as directed with other sun protection measures (see Directions), may decrease the risk of skin cancer and early skin aging caused by the sun.		<b>Warnings</b>		For external use only		Do not use on damaged or broken skin.		When using this product: Keep out of the eyes. Rinse with water to remove.		■ Stop use and ask a doctor, if rash occurs.		Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.		<b>Directions</b>		■ Apply liberally 15 minutes before sun exposure.		■ Use a water resistant sunscreen if swimming or sweating.		■ Reapply at least every 2 hours.		■ Sun Protection Measures: Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:		• Limit time in the sun, especially from 10 a.m. to 2 p.m.		• Wear long-sleeved shirts, pants, hats, and sunglasses.		• Children under 6 months: Ask a doctor.	
Drug Facts																																																	
Active Ingredients	Purpose																																																
Erythrolol 2.7%	Sunscreen																																																
Octinoxide 7%	Sunscreen																																																
Titanium dioxide 7.16%	Sunscreen																																																
Zinc Oxide 2.91%	Sunscreen																																																
<b>Uses</b>																																																	
■ Helps prevent sunburn.																																																	
■ If used as directed with other sun protection measures (see Directions), may decrease the risk of skin cancer and early skin aging caused by the sun.																																																	
<b>Warnings</b>																																																	
For external use only																																																	
Do not use on damaged or broken skin.																																																	
When using this product: Keep out of the eyes. Rinse with water to remove.																																																	
■ Stop use and ask a doctor, if rash occurs.																																																	
Keep out of reach of children. If product is swallowed, get medical help or contact a Poison Control Center right away.																																																	
<b>Directions</b>																																																	
■ Apply liberally 15 minutes before sun exposure.																																																	
■ Use a water resistant sunscreen if swimming or sweating.																																																	
■ Reapply at least every 2 hours.																																																	
■ Sun Protection Measures: Spending time in the sun increases your risk of skin cancer and early skin aging. To decrease this risk, regularly use a sunscreen with a Broad Spectrum SPF value of 15 or higher and other sun protection measures including:																																																	
• Limit time in the sun, especially from 10 a.m. to 2 p.m.																																																	
• Wear long-sleeved shirts, pants, hats, and sunglasses.																																																	
• Children under 6 months: Ask a doctor.																																																	
																																																	

**OHUI POWDERY METAL CUSHION COMPACT 02 NATURAL BEIGE**  
 titanium dioxide powder

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:53208-603
<b>Route of Administration</b>	TOPICAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP) (TITANIUM DIOXIDE - UNII:15FIX9V2JP)	TITANIUM DIOXIDE	1.077 g in 15 g
<b>ENSULIZOLE</b> (UNII: 9YQ9D11W42) (ENSULIZOLE - UNII:9YQ9D11W42)	ENSULIZOLE	0.405 g in 15 g
<b>OCTINOXATE</b> (UNII: 4Y5P7MUD51) (OCTINOXATE - UNII:4Y5P7MUD51)	OCTINOXATE	1.05 g in 15 g
<b>ZINC OXIDE</b> (UNII: SOI2LOH54Z) (ZINC OXIDE - UNII:SOI2LOH54Z)	ZINC OXIDE	0.436 g in 15 g

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>DIPROPYLENE GLYCOL</b> (UNII: E107L85C40)	
<b>MAGNESIUM PALMITOYL GLUTAMATE</b> (UNII: DH37YM1F48)	
<b>PHENOXYETHANOL</b> (UNII: HIE492ZZ3T)	
<b>ALUMINUM HYDROXIDE</b> (UNII: 5QB0T2IUN0)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>ADENOSINE</b> (UNII: K72T3FS567)	
<b>SODIUM PALMITOYL SARCO SINATE</b> (UNII: 7297LY09YF)	
<b>TRIETHOXYCAPRYL YLSILANE</b> (UNII: LDC331P08E)	
<b>PALMITIC ACID</b> (UNII: 2V16EO95H1)	
<b>HEXYL LAURATE</b> (UNII: 4CG9F9W01Q)	
<b>POLYGLYCERYL-4 ISOSTEARATE</b> (UNII: 820DPX33S7)	
<b>ALBIZIA JULIBRISSIN BARK</b> (UNII: 0J9G6W44DV)	
<b>NIACINAMIDE</b> (UNII: 25X51I8RD4)	
<b>BIO SACCHARIDE GUM-1</b> (UNII: BB4PU4V09H)	
<b>BUTYLENE GLYCOL</b> (UNII: 3XUS85K0RA)	
<b>CETYL PEG/PPG-10/1 DIMETHICONE (HLB 2)</b> (UNII: V2W71V8T0X)	
<b>MICA</b> (UNII: V8A1AW0880)	
<b>PEG-10 DIMETHICONE (600 CST)</b> (UNII: 8PR7V1SVM0)	
<b>POLY(METHYL METHACRYLATE; 450000 MW)</b> (UNII: Z47NNT4J11)	
<b>TROMETHAMINE</b> (UNII: 023C2WHX2V)	
<b>1,2-HEXANEDIOL</b> (UNII: TR046Y3K1G)	
<b>PALMITOYL PROLINE</b> (UNII: I49727TDYF)	
<b>CYCLOMETHICONE 5</b> (UNII: 0THT5PC10R)	
<b>METHYL TRIMETHICONE</b> (UNII: S73ZQ10GXM)	
<b>DISTEARDIMONIUM HECTORITE</b> (UNII: X687XDK09L)	
<b>STEARIC ACID</b> (UNII: 4ELV7Z65AP)	
<b>SORBITAN SESQUIOLEATE</b> (UNII: 0W8RRI5W5A)	
<b>ZINC STEARATE</b> (UNII: H92E6QA4FV)	
<b>DISTEARYLDIMONIUM CHLORIDE</b> (UNII: OM9573ZX3X)	
<b>ALPHA-TOCOPHEROL ACETATE</b> (UNII: 9E8X80D2L0)	
<b>DIMETHICONE/VINYL DIMETHICONE CROSSPOLYMER (SOFT PARTICLE)</b> (UNII: 9E4CO0W6C5)	
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:53208-603-01	2 in 1 CARTON	01/25/2016	
1		15 g in 1 CONTAINER; Type 1: Convenience Kit of Co-Package		

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part352	01/25/2016	

**Labeler** - LG Household and Health Care Inc (688276187)**Establishment**

Name	Address	ID/FEI	Business Operations
LG Household and Health Care Inc		688276187	manufacture(53208-603)

**Establishment**

Name	Address	ID/FEI	Business Operations
ZENISCE Co.Ltd.		690188305	manufacture(53208-603)

Revised: 2/2016

LG Household and Health Care Inc