

CONEX- dexbrompheniramine maleate, pseudoephedrine hcl liquid
Llorens Pharmaceutical International Division

Llorens-Conex 812

Active Ingredients (in each 5 mL tsp.)

Dexbrompheniramine Maleate, USP 2 mg Antihistamine

Pseudoephedrine HCL, USP 60 mg Nasal Decongestant

Antihistamine

Nasal Decongestant

Uses

- Temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- helps decongest sinus opening and sinus passages
- reduces swelling of nasal passages, shrinks swollen membranes, and temporarily restores freer breathing through the nose
- temporarily alleviates the following symptoms due to hay fever (allergic rhinitis)
- runny nose
- sneezing
- itching of the nose or throat
- itchy and watery eyes

Warnings:

- **Do not use if** you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if you are taking a prescription drug that contains an MAOI, ask your doctor or pharmacist before taking this product.

Ask a doctor before use if you are taking sedatives or tranquilizers.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- difficulty in urination due to the enlargement of the prostate gland.

When using this product

- do not exceed the recommended dosage
- excitability may occur, especially in children
- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- symptoms do not improve within 7 days or occur with a fever.

Keep out of the reach of children.In case of overdose, get medical help or contact a Poison Control Center right away.

If pregnant or breast-feeding,ask a Health Professional before use.

Directions:Do not exceed more than 4 doses in any 24-hour period or as directed by a doctor.

| | |
|--|--|
| Age | Dose |
| Adults and children 12 years of age and over | take two teaspoonfuls (10 mL) every 4 to 6 hours |
| Children under 12 years of age | Ask a doctor |

Inactive Ingredients: citric acid, flavor, glycerine, menthol, methylparaben, propylene glycol, propylparaben, purified water, sodium citrate and sucralose

Questions or Comments? 1-866-595-5598

Drug Facts

Active ingredients (in each 5 mL tsp.) **Purposes**
Dexbrompheniramine Maleate, USP.....2 mg Antihistamine
Pseudoephedrine HCl, USP.....60 mg Nasal Decongestant

Uses: ■ Temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies ■ helps decongest sinus openings and sinus passages ■ reduces swelling of nasal passages, shrinks swollen membranes, and temporarily restores freer breathing through the nose ■ temporarily alleviates the following symptoms due to hay fever (allergic rhinitis) ■ runny nose ■ sneezing ■ itching of the nose or throat ■ itchy and watery eyes

Warnings
Do not use if you are taking a monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before using this product.
Ask a doctor before use if you are taking sedatives or tranquilizers.
Ask a doctor before use if you have ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ difficulty in urination due to enlargement of prostate gland.
When using this product ■ do not exceed the recommended dose ■ excitability may occur, especially in children ■ drowsiness may occur ■ avoid alcoholic beverages ■ alcohol, sedatives and tranquilizers may increase drowsiness ■ use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if
■ nervousness, dizziness, or sleeplessness occurs
■ symptoms do not improve within 7 days or occur with a fever. ➡



NDC 54859-812-16

CONEX

Antihistamine
Nasal Decongestant

Cherry Flavor

Drug Facts (continued)

If pregnant or breast-feeding, ask a Health Professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions: Do not exceed more than 4 doses in any 24-hour period or as directed by a doctor.

| Age | Dose |
|--|--|
| Adults and children 12 years of age and over | Take one teaspoonful (5 mL) every 4 to 6 hours |
| Children under 12 years of age | Ask a doctor |

Other information ■ Store at room temperature 20°-25°C (68°- 77°F). Permitted excursions between 15°-30°C (59°-86°F). ■ Tamper evident: Do not use if foil seal over bottle opening is torn, broken or missing.
■ **Pharmacists** - Preserve and dispense in tight, light- resistant container with a child resistant cap as defined in the USP.

Inactive ingredients: Citric Acid, Flavor, Glycerin, Menthol, Methylparaben, Propylene Glycol, Propylparaben, Purified Water, Sodium Citrate and Sucralose.

Questions or Comments? 1-866-595-5598

THIS IS A BULK CONTAINER NOT INTENDED FOR RETAIL

Code #: L-254 Rev. 12/24



3 54859 81216 2

16 fl.oz. (473 ml)

Manufactured by:
Llorens Pharmaceutical International Division, Inc.
Miami, FL 33147
MADE IN USA 
Visit us at www.llorensparm.com

Lot:
Exp:

CONEX

dexbrompheniramine maleate, pseudoephedrine hcl liquid

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:54859-812 |
| Route of Administration | ORAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|----------------------------------|------------------|
| PSEUDOEPHEDRINE HYDROCHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F) | PSEUDOEPHEDRINE HYDROCHLORIDE | 60 mg in 5 mL |
| DEXBROMPHENIRAMINE MALEATE (UNII: BPA9UT29BS) (DEXBROMPHENIRAMINE - UNII:75T64B71RP) | DEXBROMPHENIRAMINE MALEATE | 2 mg in 5 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|---|-----------------|
| CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| MENTHOL (UNII: L7T10EIP3A) | |
| METHYLPARABEN (UNII: A2I8C7HI9T) | |
| WATER (UNII: 059QF0K00R) | |
| PROPYLENE GLYCOL (UNII: 6DC9Q167V3) | |
| PROPYLPARABEN (UNII: Z8IX2SC1OH) | |
| SUCRALOSE (UNII: 96K6UQ3ZD4) | |
| SODIUM CITRATE (UNII: 1Q73Q2JULR) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|----------|------------------|---|-----------------------------|---------------------------|
| 1 | NDC:54859-812-16 | 473 mL in 1 BOTTLE; Type 0: Not a Combination Product | 12/01/2024 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|---------------------------|---|-----------------------------|---------------------------|
| OTC Monograph Drug | M012 | 12/01/2024 | |

Labeler - Llorens Pharmaceutical International Division (037342305)

Revised: 12/2025

Llorens Pharmaceutical International Division