

**CONEX- dexbrompheniramine maleate, pseudoephedrine hcl liquid**  
**Llorens Pharmaceutical International Division**

-----  
**Llorens-Conex 812**

**Active Ingredients (in each 5 mL tsp.)**

Dexbrompheniramine Maleate, USP ..... 2 mg ..... Antihistamine

Pseudoephedrine HCL, USP ..... 60 mg ..... Nasal Decongestant

**Antihistamine**

**Nasal Decongestant**

**Uses**

- Temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- helps decongest sinus opening and sinus passages
- reduces swelling of nasal passages, shrinks swollen membranes, and temporarily restores freer breathing through the nose
- temporarily alleviates the following symptoms due to hay fever (allergic rhinitis)
- runny nose
- sneezing
- itching of the nose or throat
- itchy and watery eyes

**Warnings:**

- **Do not use if** you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if you are taking a prescription drug that contains an MAOI, ask your doctor or pharmacist before taking this product.

**Ask a doctor before use if you are** taking sedatives or tranquilizers.

**Ask a doctor before use if you have**

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- difficulty in urination due to the enlargement of the prostate gland.

**When using this product**

- do not exceed the recommended dosage
- excitability may occur, especially in children
- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**

- nervousness, dizziness or sleeplessness occur
- symptoms do not improve within 7 days or occur with a fever.

**Keep out of the reach of children.**In case of overdose, get medical help or contact a Poison Control Center right away.

**If pregnant or breast-feeding,**ask a Health Professional before use.

**Directions:**Do not exceed more than 4 doses in any 24-hour period or as directed by a doctor.

Age	Dose
Adults and children 12 years of age and over	take two teaspoonfuls (10 mL) every 4 to 6 hours
Children under 12 years of age	Ask a doctor

**Inactive Ingredients:** citric acid, flavor, glycerine, menthol, methylparaben, propylene glycol, propylparaben, purified water, sodium citrate and sucralose

**Questions or Comments? 1-866-595-5598**

**Drug Facts**

**Active ingredients (in each 5 mL tsp.)**      **Purposes**  
Dexbrompheniramine Maleate, USP.....2 mg ..... Antihistamine  
Pseudoephedrine HCl, USP.....60 mg ..... Nasal Decongestant

**Uses:** ■ Temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies ■ helps decongest sinus openings and sinus passages ■ reduces swelling of nasal passages, shrinks swollen membranes, and temporarily restores freer breathing through the nose ■ temporarily alleviates the following symptoms due to hay fever (allergic rhinitis) ■ runny nose ■ sneezing ■ itching of the nose or throat ■ itchy and watery eyes

**Warnings**  
Do not use if you are taking a monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before using this product.  
Ask a doctor before use if you are taking sedatives or tranquilizers.  
Ask a doctor before use if you have ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ difficulty in urination due to enlargement of prostate gland.  
When using this product ■ do not exceed the recommended dose ■ excitability may occur, especially in children ■ drowsiness may occur ■ avoid alcoholic beverages ■ alcohol, sedatives and tranquilizers may increase drowsiness ■ use caution when driving a motor vehicle or operating machinery

**Stop use and ask a doctor if**  
■ nervousness, dizziness, or sleeplessness occurs  
■ symptoms do not improve within 7 days or occur with a fever. ➡



NDC 54859-812-16

# CONEX

Antihistamine  
Nasal Decongestant

Cherry Flavor

**Drug Facts (continued)**

If pregnant or breast-feeding, ask a Health Professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

**Directions:** Do not exceed more than 4 doses in any 24-hour period or as directed by a doctor.

Age	Dose
Adults and children 12 years of age and over	Take one teaspoonful (5 mL) every 4 to 6 hours
Children under 12 years of age	Ask a doctor

**Other information** ■ Store at room temperature 20°-25°C (68°- 77°F). Permitted excursions between 15°-30°C (59°-86°F). ■ Tamper evident: Do not use if foil seal over bottle opening is torn, broken or missing.  
■ **Pharmacists** - Preserve and dispense in tight, light- resistant container with a child resistant cap as defined in the USP.

**Inactive ingredients:** Citric Acid, Flavor, Glycerin, Menthol, Methylparaben, Propylene Glycol, Propylparaben, Purified Water, Sodium Citrate and Sucralose.

**Questions or Comments?** 1-866-595-5598

THIS IS A BULK CONTAINER NOT INTENDED FOR RETAIL

Code #: L-254      Rev. 12/24



3 54859 81216 2

16 fl.oz. (473 ml)

Manufactured by:  
Llorens Pharmaceutical International Division, Inc.  
Miami, FL 33147  
MADE IN USA   
Visit us at [www.llorensparm.com](http://www.llorensparm.com)

Lot:  
Exp:

**CONEX**

dexbrompheniramine maleate, pseudoephedrine hcl liquid

**Product Information**

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:54859-812
<b>Route of Administration</b>	ORAL		

**Active Ingredient/Active Moiety**

<b>Ingredient Name</b>	<b>Basis of Strength</b>	<b>Strength</b>
<b>DEXBROMPHENIRAMINE MALEATE</b> (UNII: BPA9UT29BS) (DEXBROMPHENIRAMINE - UNII:75T64B71RP)	DEXBROMPHENIRAMINE MALEATE	2 mg in 5 mL
<b>PSEUDOEPHEDRINE HYDROCHLORIDE</b> (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	60 mg in 5 mL

**Inactive Ingredients**

<b>Ingredient Name</b>	<b>Strength</b>
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>MENTHOL</b> (UNII: L7T10EIP3A)	
<b>METHYLPARABEN</b> (UNII: A2I8C7HI9T)	
<b>WATER</b> (UNII: 059QF0K00R)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	

**Packaging**

<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:54859-812-16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	12/01/2024	

**Marketing Information**

<b>Marketing Category</b>	<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug	M012	12/01/2024	

**Labeler** - Llorens Pharmaceutical International Division (037342305)

Revised: 12/2024

Llorens Pharmaceutical International Division