

SDA LABORATORIES TUSSIN- guaifenesin liquid
SDA Laboratories, Inc.

SDA-Tussin 563

In each 5 mL

Guaifenesin - 100 mg

Purpose

Expectorant

Uses

- helps loosen phlegm (mucus)
- helps thin bronchial secretions to make coughs more productive

Warnings

Ask a doctor before use if you have

- a cough with too much phlegm (mucus)
- a cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema

Stop use and ask a doctor if cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding,ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away

Directions

- Do not take more than 6 doses in any 24 hour period
- Measure only with dosing cup provided
- mL- milliliters
- Adults and children over 12 years of age: take 2 teaspoonfuls (10 mL) every 4 hours as needed or as directed by a doctor
- Children under 12 years of age: Do not use

Inactive ingredients: cherry flavor, citric acid, FD&C red #40, menthol, methylparaben, propylene glycol, propylparaben, purified water, sodium citrate, sucralose

Questions or comments? 1-203-861-0005



Compare to the active ingredient in Robitussin®

Drug Facts

Active ingredient (in each 5 mL teaspoonful) Purpose
Guaifenesin, USP 100 mg.....Expectorant

Uses

■ helps loosen phlegm (mucus) ■ helps thin bronchial secretions to make coughs more productive

Warnings

Do not use ■ if you ever had an allergic reaction to any of the ingredients in this product

Ask a doctor before use if you have

■ a cough with too much phlegm (mucus) ■ a cough that lasts or is chronic such as occurs with smoking, asthma, chronic bronchitis or emphysema.

Stop use and ask a doctor if cough lasts more than 7 days, comes back or is accompanied by fever, rash, or headache that lasts. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. ▶

Code: L-232 Rev. 06/24

Manufactured for:
SDA Laboratories, Inc.
280 Railroad Ave.,
Greenwich, CT 06830

Lot # :

Exp Date :

NDC 66424-563-16

TUSSIN

Guaifenesin

- Loosens and Relieves Chest Congestion
- Alcohol Free
- Sugar Free

Cherry Mint

EXPECTORANT
ORAL SOLUTION

16 FL.OZ. (473 mL)

Drug Facts (continued)

Directions

- Do not take more than 6 doses in any 24 hour period
- Do not exceed recommended dose
- mL-milliliters
- Adults and children 12 years and over: take 2 teaspoonfuls (10 mL) every 4 hours as needed or as directed by a doctor.
- Children under 12 years of age: Do not use

Other Information

- each teaspoonful (5 mL) contains: sodium 2 mg
- store at room temperature 20°-25°C (68°-77°F)
- protect from freezing ■ do not refrigerate ■ protect from light
- Pharmacists-Preserve and dispense in a light, light resistant container with a child resistant cap as defined in the USP
- **TAMPER-EVIDENT:** Do not use if foil seal over bottle opening is torn, broken or missing

Inactive ingredients: cherry flavor, citric acid, FD&C Red #40, menthol, methylparaben, propylene glycol, propylparaben, purified water, sodium citrate and sucralose.

Questions or comments? 1-203-861-0005

*This product is not manufactured or distributed by the owner of the registered trademark Robitussin®.

THIS IS A BULK CONTAINER NOT INTENDED FOR RETAIL



SDA LABORATORIES TUSSIN

guaifenesin liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:66424-563
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	100 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
MENTHOL (UNII: L7T10EIP3A)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:66424-563-16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	12/01/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/01/2024	

Labeler - SDA Laboratories, Inc. (948067889)

Revised: 12/2024

SDA Laboratories, Inc.