

CONEX PEDIATRIC- dextbrompheniramine maleate, pseudoephedrine hcl liquid
Llorens Pharmaceutical International Division

Llorens-ConexPed 803

Active Ingredients (in each 5 mL tsp.)

Dextbrompheniramine Maleate, USP 1 mg Antihistamine

Pseudoephedrine HCL, USP 30 mg Nasal Decongestant

Antihistamine

Nasal Decongestant

Uses

- Temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- helps decongest sinus opening and sinus passages
- reduces swelling of nasal passages, shrinks swollen membranes, and temporarily restores freer breathing through the nose
- temporarily alleviates the following symptoms due to hay fever (allergic rhinitis)
- runny nose
- sneezing
- itching of the nose or throat
- itchy and watery eyes

Warnings:

- **Do not use if** you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if you are taking a prescription drug that contains an MAOI, ask your doctor or pharmacist before taking this product.

Ask a doctor before use if you are taking sedatives or tranquilizers.

Ask a doctor before use if you have

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- difficulty in urination due to the enlargement of the prostate gland.

When using this product

- do not exceed the recommended dosage
- excitability may occur, especially in children
- drowsiness may occur
- avoid alcoholic beverages
- alcohol, sedatives and tranquilizers may increase drowsiness
- use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- symptoms do not improve within 7 days or occur with a fever.

Keep out of the reach of children.In case of overdose, get medical help or contact a Poison Control Center right away.

If pregnant or breast-feeding,ask a Health Professional before use.

Directions:Do not exceed more than 4 doses in any 24-hour period or as directed by a doctor.

Age	Dose
Adults and children 12 years of age and over	take two teaspoonfuls (10 mL) every 4 to 6 hours
Children under 12 years of age	take 1 teaspoonful (5 mL) every 4 to 6 hours
Children under 6 years of age	ask a doctor

Inactive Ingredients: citric acid, flavor, glycerine, menthol, methylparaben, propylene glycol, propylparaben, purified water, sodium citrate and sucralose

Questions or Comments? 1-866-595-5598

Drug Facts

Active Ingredients (in each 5 mL tsp.) Purpose
Dexbrompheniramine Maleate, USP..... 1 mgAntihistamine
Pseudoephedrine HCl, USP..... 30 mgNasal Decongestant

Uses ■ Temporarily relieves nasal congestion due to the common cold, hay fever or other upper respiratory allergies ■ helps decongest sinus openings and sinus passages ■ reduces swelling of nasal passages, shrinks swollen membranes, and temporarily restores freer breathing through the nose ■ temporarily alleviates the following symptoms due to hay fever (allergic rhinitis) ■ runny nose ■ sneezing ■ itching of the nose or throat ■ itchy and watery eyes

Warnings
■ Do not use if you are taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if you are taking a prescription drug that contains an MAOI, ask your doctor or pharmacist before taking this product.

Ask a doctor before use if you are taking sedatives or tranquilizers.

Ask a doctor before use if you have ■ heart disease ■ high blood pressure ■ thyroid disease ■ diabetes ■ glaucoma ■ difficulty in urination due to the enlargement of the prostate gland.

When using this product ■ do not exceed the recommended dosage ■ excitability may occur, especially in children ■ drowsiness may occur ■ avoid alcoholic beverages ■ alcohol, sedatives and tranquilizers may increase drowsiness ■ use caution when driving a motor vehicle or operating machinery

Stop use and ask a doctor if ■ nervousness, dizziness or sleeplessness occur ■ symptoms do not improve within 7 days or occur with a fever.

NDC 54859-803-16

Conex[®]

PEDIATRIC

Oral Solution

- Nasal & Sinus Congestion
- Runny Nose, Sneezing
- Itchy, Watery Eyes

• ANTIHISTAMINE
• NASAL DECONGESTANT

Cherry

16 FL OZ (473 mL)

Manufactured by:
LLORENS
Llorens Pharmaceutical
International Division
Miami, FL 33147
www.llorensparm.com



Drug Facts (continued)

If pregnant or breast-feeding, ask a Health Professional before use. **Keep out of reach of children.** In case of accidental overdose, get medical help or contact a Poison Control Center right away.

Directions Do not exceed more than 4 doses in any 24-hour period or as directed by a doctor.

Age	Dose
Adults and children 12 years of age and over	take two teaspoonfuls (10 mL) every 4 to 6 hours
Children 6 to under 12 years of age	take 1 teaspoonful (5 mL) every 4 to 6 hours
Children under 6 years of age	ask a doctor

Other information ■ Store at room temperature 15-30°C (59-86°F).
■ Tamper-Evident: Do not use if foil seal over bottle opening is torn, broken, or missing.
■ Pharmacist - Preserve and dispense in light, light-resistant container with a child resistant cap as defined in the USP.

Inactive Ingredients: citric acid, flavors, glycerin, menthol, methylparaben, propylene glycol, propylparaben, purified water, sodium citrate, sucralose

Questions or Comments? 1-866-595-5598

THIS IS A BULK CONTAINER NOT INTENDED FOR RETAIL.

CODE# L-86 REV:07/22



Lot. #
Exp. Date:

CONEX PEDIATRIC

dexbrompheniramine maleate, pseudoephedrine hcl liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:54859-803
---------------------	----------------	---------------------------	---------------

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DEXBROMPHENIRAMINE MALEATE (UNII: BPA9UT29BS) (DEXBROMPHENIRAMINE - UNII:75T64B71RP)	DEXBROMPHENIRAMINE MALEATE	1 mg in 5 mL
PSEUDOEPHEDRINE HYDROCHLORIDE (UNII: 6V9V2RYJ8N) (PSEUDOEPHEDRINE - UNII:7CUC9DDI9F)	PSEUDOEPHEDRINE HYDROCHLORIDE	30 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
GLYCERIN (UNII: PDC6A3C0OX)	
MENTHOL (UNII: L7T10EIP3A)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
WATER (UNII: 059QF0KO0R)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54859-803-16	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	12/01/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	12/01/2024	

Labeler - Llorens Pharmaceutical International Division (037342305)

Revised: 12/2024

Llorens Pharmaceutical International Division