

**LLORENS CARE CHILDREN ACETAMINOPHEN- acetaminophen suspension**  
**Llorens Pharmaceutical International Division, Inc.**

-----  
**Llorens-APAPSusChild 850**

**Active ingredient (in each 5mL = 1 teaspoon)**

**Acetaminophen 160 mg**

**Purpose**

**Pain Reliever / Fever Reducer**

Uses

- temporarily reduces fever
- temporarily relieves minor aches and pains due to:
- headache
- sore throat
- the common cold
- toothache

**Warnings**

**Liver warning**

This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen.

Allergy Alert:

acetaminophen may cause severe skin reactions. Symptoms may include

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning**

if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use**

- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this

product.

When using this product: Do not exceed recommended dose.

**Ask a doctor before use if your child has** liver disease.

**Ask a doctor or pharmacist before use if your child is** taking the blood thinning drug warfarin.

**Stop use and ask a doctor if**

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present. These could be signs of a serious condition.

**Keep out of reach of children.**

**Overdose warning**

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

**Directions**

- do not give more than directed
- shake well before using
- this product does not contain directions or complete warnings for adult use
- find dose in chart below; if possible, use weight to dose, otherwise use age
- dosage may be repeated every 4 hours, or as directed by a doctor
- do not give more than 5 times in 24 hours

Weight (lbs)	under 24	24-35	36-47	48-59	60-71	72-95	over 95
Age (years)	under 2	2 to 3	4 to 5	6 to 8	9 to 10	11	12 to adult
Dosage Teaspoonful (tsp)	consult physician	1 tsp (5 mL)	1 1/2 tsp (7.5 mL)	2 tsp (10 mL)	2 1/2 tsp (12.5 mL)	3 tsp (15 mL)	4 tsp (20 mL)

**Inactive ingredients**

artificial and natural grape flavor, citric acid, glycerin, microcrystalline cellulose and carboxymethylcellulose, sodium, potassium sorbate, purified water, sorbitol solution, sucralose, and xanthan gum

**QUESTIONS? COMMENTS?**

Call 1-866-595-5598

NDC: 54859-850-16

**Drug Facts**

**Active Ingredient (per 5 mL teaspoonful)** Acetaminophen 160 mg .....Pain Reliever/Fever Reducer **Purpose**

**Uses**  
 • temporarily reduces fever • temporarily relieves minor aches and pains due to: • headache • sore throat • the common cold  
 • toothache

**Warnings**  
**Liver Warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes • more than 5 doses in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen

**Allergy Alert:** acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash

**Sore Throat Warning:** if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor

**Do not use** • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if your child is allergic to acetaminophen or any of the inactive ingredients in this product

**When using this product:** Do not exceed recommended dose.

**Ask a doctor before use if your child has liver disease.**

**Ask a doctor before use if your child is taking the blood thinning drug warfarin.**

**Stop use and ask a doctor if** • pain gets worse or lasts more than 5 days • fever gets worse or lasts more than 3 days • new symptoms occur • redness or swelling is present. These could be signs of a serious condition.

**Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical even if you do not notice any signs or symptoms.



# CHILDREN'S ACETAMINOPHEN

## Oral Suspension

160 mg / 5 mL

Dye Free  
 Alcohol Free  
 Aspirin Free

**Drug Facts (continued)**

**Directions** • do not give more than directed  
 • shake well before using  
 • this product does not contain directions or complete warnings for adult use  
 • find dose in chart below; if possible, use weight to dose; otherwise use age  
 • dosage may be repeated every 4 hours, or as directed by a doctor  
 • do not give more than 5 times in 24 hours

Weight (kg)	under 24	24 - 35	36 - 47	48 - 59	60 - 71	72 - 85	over 85
Age (years)	under 2	2 to 3	4 to 5	6 to 8	9 to 10	11	12 to adult
Dosage Teaspoonful (tsp)	consult physician	1 tsp (5 mL)	1 1/2 tsp (7.5 mL)	2 tsp (10 mL)	2 1/2 tsp (12.5 mL)	3 tsp (15 mL)	4 tsp (20 mL)

**Other Information**

• store at room temperature 20° - 25°C (68°-77°F)  
 • protect from freezing • protect from light  
**Tamper-Evident:** Do not use if foil seal over bottle opening is torn, broken or missing.

**Inactive Ingredients:** artificial and natural grape flavor, citric acid, glycerin, microcrystalline cellulose and carboxymethylcellulose sodium, potassium sorbate, purified water, sorbitol solution, sucralose, sucrose and xanthan gum.

**Questions and comments? 1-866-595-5598**

\*This product is not manufactured or distributed by the owner of the registered trademark Tylenol®.

**Manufactured by:**

Llorens Pharmaceutical International Division, Inc.  
 7080 NW 37th Court  
 Miami, FL 33147  
 www.LlorensPharm.com

Rev.: 08/24

Code #: L-241



Compare to Children's Tylenol® Oral Suspension Active Ingredients\*  
 Pain Reliever / Fever Reducer

Lot #:

Exp Date:



Grape Flavor  
 16 oz (473 mL)



# LLORENS CARE CHILDREN ACETAMINOPHEN

acetaminophen suspension

## Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:54859-850
<b>Route of Administration</b>	ORAL		

## Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

## Inactive Ingredients

Ingredient Name	Strength
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>POTASSIUM SORBATE</b> (UNII: 1VPU26JZZ4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>SODIUM CITRATE</b> (UNII: 1Q73Q2JULR)	
<b>CITRIC ACID MONOHYDRATE</b> (UNII: 2968PHW8QP)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED</b> (UNII: K679OBS311)	

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54859-850-16	473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/01/2024	
2	NDC:54859-850-04	118 mL in 1 BOTTLE; Type 0: Not a Combination Product	12/01/2024	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	12/01/2024	

**Labeler** - Llorens Pharmaceutical International Division, Inc. (037342305)

Revised: 12/2024

Llorens Pharmaceutical International Division, Inc.