

LLORENS CARE CHILDREN ACETAMINOPHEN- acetaminophen suspension
Llorens Pharmaceutical International Division, Inc.

Llorens-APAPSusChild 850

Active ingredient (in each 5mL = 1 teaspoon)

Acetaminophen 160 mg

Purpose

Pain Reliever / Fever Reducer

Uses

- temporarily reduces fever
- temporarily relieves minor aches and pains due to:
- headache
- sore throat
- the common cold
- toothache

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen.

Allergy Alert:

acetaminophen may cause severe skin reactions. Symptoms may include

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if your child is allergic to acetaminophen or any of the inactive ingredients in this

product.

When using this product: Do not exceed recommended dose.

Ask a doctor before use if your child has liver disease.

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

Stop use and ask a doctor if

- pain gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- new symptoms occur
- redness or swelling is present. These could be signs of a serious condition.

Keep out of reach of children.

Overdose warning

Taking more than the recommended dose (overdose) may cause liver damage. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical even if you do not notice any signs or symptoms.

Directions

- do not give more than directed
- shake well before using
- this product does not contain directions or complete warnings for adult use
- find dose in chart below; if possible, use weight to dose, otherwise use age
- dosage may be repeated every 4 hours, or as directed by a doctor
- do not give more than 5 times in 24 hours

Weight (lbs)	under 24	24-35	36-47	48-59	60-71	72-95	over 95
Age (years)	under 2	2 to 3	4 to 5	6 to 8	9 to 10	11	12 to adult
Dosage Teaspoonful (tsp)	consult physician	1 tsp (5 mL)	1 1/2 tsp (7.5 mL)	2 tsp (10 mL)	2 1/2 tsp (12.5 mL)	3 tsp (15 mL)	4 tsp (20 mL)

Inactive ingredients

artificial and natural grape flavor, citric acid, glycerin, microcrystalline cellulose and carboxymethylcellulose, sodium, potassium sorbate, purified water, sorbitol solution, sucralose, and xanthan gum

QUESTIONS? COMMENTS?

Call 1-866-595-5598

NDC: 54859-850-16

Drug Facts

Active Ingredient (per 5 mL teaspoonful) Acetaminophen 160 mgPain Reliever/Fever Reducer **Purpose**

Uses
 • temporarily reduces fever • temporarily relieves minor aches and pains due to: • headache • sore throat • the common cold
 • toothache

Warnings
Liver Warning: This product contains acetaminophen. Severe liver damage may occur if your child takes • more than 5 doses in 24 hours, which is the maximum daily amount • with other drugs containing acetaminophen

Allergy Alert: acetaminophen may cause severe skin reactions. Symptoms may include: • skin reddening • blisters • rash

Sore Throat Warning: if sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea or vomiting, consult a doctor

Do not use • with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. • if your child is allergic to acetaminophen or any of the inactive ingredients in this product

When using this product: Do not exceed recommended dose.

Ask a doctor before use if your child has liver disease.

Ask a doctor before use if your child is taking the blood thinning drug warfarin.

Stop use and ask a doctor if • pain gets worse or lasts more than 5 days • fever gets worse or lasts more than 3 days • new symptoms occur • redness or swelling is present. These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical even if you do not notice any signs or symptoms.



CHILDREN'S ACETAMINOPHEN

Oral Suspension

160 mg / 5 mL

Dye Free
 Alcohol Free
 Aspirin Free

Drug Facts (continued)

Directions • do not give more than directed
 • shake well before using
 • this product does not contain directions or complete warnings for adult use
 • find dose in chart below; if possible, use weight to dose; otherwise use age
 • dosage may be repeated every 4 hours, or as directed by a doctor
 • do not give more than 5 times in 24 hours

Weight (kg)	under 24	24 - 35	36 - 47	48 - 59	60 - 71	72 - 85	over 85
Age (years)	under 2	2 to 3	4 to 5	6 to 8	9 to 10	11	12 to adult
Dosage Teaspoonful (tsp)	consult physician	1 tsp (5 mL)	1 1/2 tsp (7.5 mL)	2 tsp (10 mL)	2 1/2 tsp (12.5 mL)	3 tsp (15 mL)	4 tsp (20 mL)

Other Information

• store at room temperature 20° - 25°C (68°-77°F)
 • protect from freezing • protect from light
 • **Tamper-Evident:** Do not use if foil seal over bottle opening is torn, broken or missing.

Inactive Ingredients: artificial and natural grape flavor, citric acid, glycerin, microcrystalline cellulose and carboxymethylcellulose sodium, potassium sorbate, purified water, sorbitol solution, sucralose, sucrose and xanthan gum.

Questions and comments? 1-866-595-5598

*This product is not manufactured or distributed by the owner of the registered trademark Tylenol®.

Manufactured by:

Llorens Pharmaceutical International Division, Inc.
 7080 NW 37th Court
 Miami, FL 33147
 www.LlorensPharm.com

Rev.: 08/24

Code #: L-241



Compare to Children's Tylenol® Oral Suspension Active Ingredients*
 Pain Reliever / Fever Reducer

Lot #:

Exp Date:



Grape Flavor
 16 oz (473 mL)



LLORENS CARE CHILDREN ACETAMINOPHEN

acetaminophen suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:54859-850
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	160 mg in 5 mL

Inactive Ingredients

Ingredient Name	Strength
SUCROSE (UNII: C151H8M554)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
POTASSIUM SORBATE (UNII: 1VPU26JZZ4)	
XANTHAN GUM (UNII: TTV12P4NEE)	
GLYCERIN (UNII: PDC6A3C0OX)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	
CITRIC ACID MONOHYDRATE (UNII: 2968PHW8QP)	
SORBITOL (UNII: 506T60A25R)	
WATER (UNII: 059QF0KO0R)	
MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)	
CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED (UNII: K679OBS311)	

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:54859-850-16	473 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	12/01/2024	
2	NDC:54859-850-04	118 mL in 1 BOTTLE; Type 0: Not a Combination Product	12/01/2024	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M013	12/01/2024	

Labeler - Llorens Pharmaceutical International Division, Inc. (037342305)

Revised: 12/2025

Llorens Pharmaceutical International Division, Inc.