

**MULTI-SYMPTOM COLD AND FEVER CHILDRENS- acetaminophen,
dextromethorphan hbr, guaifenesin, phenylephrine hcl solution
CVS PHARMACY**

CVS 44-075

Active ingredients (in each 10 mL)

Acetaminophen 325 mg
Dextromethorphan HBr 10 mg
Guaifenesin 200 mg
Phenylephrine HCl 5 mg

Purpose

Pain reliever/fever reducer
Cough suppressant
Expectorant
Nasal decongestant

Uses

- temporarily relieves common cold and flu symptoms:
 - minor aches and pains
 - sore throat
 - headache
 - stuffy nose
 - the intensity of coughing
 - nasal congestion
 - cough due to minor throat and bronchial irritation
 - the impulse to cough to help your child get to sleep
- temporarily reduces fever
- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if your child takes

- more than 5 doses in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen

Allergy alert: Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- liver disease
- high blood pressure
- heart disease
- thyroid disease
- diabetes
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if your child is

taking the blood thinning drug warfarin.

When using this product

do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed**
- **this product does not contain directions or complete warnings for adult use**
- do not give more than 5 doses in any 24-hour period
- mL = milliliter
- only use the dose cup provided
- repeat dose every 4 hours while symptoms last
- dose as follows or as directed by a doctor
- children 6 to under 12 years: 10 mL in dosing cup provided
- children under 6 years: do not use

Other information

- **each 10 mL contains:** sodium 5 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

anhydrous citric acid, FD&C blue #1, FD&C red #40, flavors, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sodium metabisulfite, sorbitol, sucralose

Questions or comments?

1-800-426-9391

Principal display panel

♥**CVS™**

NDC 69842-475-36

Ages

6-11

years

Mixed Berry Flavor

Children's

MULTI-SYMPTOM

COLD & FEVER

Acetaminophen

Pain Reliever, Fever Reducer

Dextromethorphan HBr, Cough Suppressant

Guaifenesin, Expectorant

Phenylephrine HCl, Nasal Decongestant

Oral Solution

Relieves: Minor aches & pains,

Fever, Cough, Nasal congestion,
Chest congestion, Stuffy nose

Thins & Loosens Mucus | No Added Alcohol

4 FL OZ (118 mL)

dosing cup
included

**TAMPER EVIDENT: DO NOT USE IF PRINTED
NECK WRAP IS BROKEN OR MISSING**

Distributed by: CVS Pharmacy, Inc.

One CVS Drive, Woonsocket, RI 02895

© 2025 CVS/pharmacy

CVS.com® 1-800-SHOP CVS

V-39965

**♥ 100% money back
guaranteed.**

[CVS.com/returnpolicy](https://www.cvs.com/returnpolicy)


50844 ORG082407536

package contains one bottle
actual size

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org



Children's MULTI-SYMPTOM COLD & FEVER

4 FL OZ (118 mL)

NDC 69842-475-36



**Ages
6-11
years**

Mixed Berry Flavor

Children's MULTI-SYMPTOM COLD & FEVER

Acetaminophen

Pain Reliever, Fever Reducer
Dextromethorphan HBr, Cough Suppressant
Guaifenesin, Expectorant
Phenylephrine HCl, Nasal Decongestant
Oral Solution

Relieves: Minor aches & pains,
Fever, Cough, Nasal congestion,
Chest congestion, Stuffy nose

Thins & Loosens Mucus | No Added Alcohol

4 FL OZ (118 mL)



TAMPER EVIDENT: DO NOT USE IF PRINTED
NECK WRAP IS BROKEN OR MISSING



package contains one bottle
actual size

Drug Facts

KEEP OUTER PACKAGE FOR
COMPLETE PRODUCT INFORMATION

Active ingredients (in each 10 mL) Purpose
Acetaminophen 325 mg Pain reliever/fever reducer
Dextromethorphan HBr 10 mg Cough suppressant
Guaifenesin 200 mg Expectorant
Phenylephrine HCl 5 mg Nasal decongestant

Uses

- temporarily relieves common cold and flu symptoms:
 - minor aches and pains
 - sore throat
 - headache
 - stuffy nose
 - the intensity of coughing
 - nasal congestion
 - cough due to minor throat and bronchial irritation
 - the impulse to cough to help your child get to sleep
 - temporarily reduces fever
 - helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive

Warnings

- Liver warning:** This product contains acetaminophen. Severe liver damage may occur if your child takes
 - more than 5 doses in 24 hours, which is the maximum daily amount
 - with other drugs containing acetaminophen
- Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:
 - skin reddening
 - blisters
 - rash
 If a skin reaction occurs, stop use and seek medical help right away.
- Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- in a child who is taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your child's prescription drug contains an MAOI, ask a doctor or pharmacist before giving this product.

Ask a doctor before use if the child has

- liver disease
- high blood pressure
- heart disease
- thyroid disease
- diabetes
- persistent or chronic cough such as occurs with asthma
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if your child is taking the blood thinning drug warfarin.

Drug Facts (continued)

When using this product do not exceed recommended dosage.

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion, or cough gets worse or lasts more than 5 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Prompt medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than directed
- this product does not contain directions or complete warnings for adult use
- do not give more than 5 doses in any 24-hour period
- mL = milliliter
- only use the dose cup provided
- repeat dose every 4 hours while symptoms last
- dose as follows or as directed by a doctor
- children 6 to under 12 years: 10 mL in dosing cup provided
- children under 6 years: do not use

Other information

- each 10 mL contains: sodium 5 mg
- store at 25°C (77°F); excursions permitted between 15°-30°C (59°-86°F)
- see end flap for expiration date and lot number

Inactive ingredients

anhydrous citric acid, FD&C blue #1, FD&C red #40, flavors, glycerin, polyethylene glycol, propylene glycol, purified water, sodium benzoate, sodium citrate dihydrate, sodium metabisulfite, sorbitol, sucralose

Questions or comments? 1-800-426-9391

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2025 CVS/pharmacy
CVS.com® 1-800-SHOP CVS
V-39965

PARENTS:
Learn about teen medicine abuse
www.StopMedicineAbuse.org



How2Recycle.com

B-0231-075-36-J
ORG082407536

#895140

FPO 100%
UPC# 050428057049

X XXXXXX XXXXXX X

No Print/No Varnish
Lot & EXP

CVS 44-075

MULTI-SYMPTOM COLD AND FEVER CHILDRENS

acetaminophen, dextromethorphan hbr, guaifenesin, phenylephrine hcl solution

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:69842-475

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 10 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 10 mL
GUAIFENESIN (UNII: 495W7451VQ) (GUAIFENESIN - UNII:495W7451VQ)	GUAIFENESIN	200 mg in 10 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYETHYLENE GLYCOL, UNSPECIFIED (UNII: 3WJQ0SDW1A)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	
SODIUM METABISULFITE (UNII: 4VON5FNS3C)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	

Product Characteristics

Color	blue	Score	
Shape		Size	
Flavor	BERRY (Mixed)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69842-475-36	1 in 1 CARTON	06/10/2021	
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

OTC Monograph Drug	M012	06/10/2021	
--------------------	------	------------	--

Labeler - CVS PHARMACY (062312574)

Establishment

Name	Address	ID/FEI	Business Operations
LNK International, Inc.		967626305	manufacture(69842-475) , pack(69842-475)

Revised: 2/2026

CVS PHARMACY