

MAXIMUM STRENGTH WART REMOVER- maximum strength wart remover liquid

Shenzhen Yangan Technology Co., Ltd.

MAXIMUM STRENGTH WART REMOVER LIQUID

Salicylic Acid.....5.0%

Wart remover

USES

- To remove warts such as plantar wart, flat wart, common wart and corns, calluses.
- The plantar wart is recognized by its location only on the bottom of the foot its tenderness, and the interruption of the footprint pattern.

WARNINGS

- The liquid can get easily volatilized & crystallized, tighten the cap after use.
- Avoid long-term contact with air while using.
- For external use only, don't contact with eyes or swallow. Children should use it under the supervision of adults.

DO NOT USE

- On damaged skin(cuts, abrasions, eczema, sunburn).
- If you are allergic to any of the ingredients in this product.
- If you are pregnant or breastfeeding.

KEEP OUT OF REACH OF CHILDREN

- If swallowed, get medical help or contact a Poison Control Center right away.

DIRECTION

- Wash the affected area.
- May soak the wart in warm water for 5 minutes.
- Dry area thoroughly.
- Using the applicator(cotton swab), apply a layer of ointment to sufficiently cover each wart
- Allow it to fully absorb and cover it with a bandage as needed.
- Repeat this procedure once or twice daily as needed (until the wart is removed) for up to 12 weeks.

OTHER INFORMATION:

- Store between 68°F to 77°F (20°C to 25°C).
- Protect from excessive heat above 104°F (40°C).
- See a doctor as you may have a more serious skin condition.

Shard paraffin, lanolin, phenol.

DIRECTION
 •Wash the affected area.
 •Dry area thoroughly.
 •Using the applicator (cotton swab), apply a layer of ointment to sufficiently cover each wart.
 •Allow it to fully absorb and cover it with a bandage as needed.
 •Repeat this procedure once or twice daily as needed (until the wart is removed) for up to 12 weeks.

MAXIMUM STRENGTH
TAG & WART REMOVER
 FAST ACTING LIQUID
 15 ml/0.5 fl. oz.
 NATURAL FPRMULA

WARNINGS
 •This liquid can get easily oxidized & crystallized.
 •Repeat this cap after use. Avoid long-term contact with or after using.
 •For external use only, don't contact with eyes or mouth.
 •Children should use it under the supervision of adults.
DO NOT USE
 •On damaged skin (cuts, abrasions, eczema, sunburn, etc.) or on any of the ingredients in this product.
 •If you are pregnant or breastfeeding.

DIRECTION
 •Wash the affected area.
 •May soak the wart in warm water for 5 minutes.
 •Dry area thoroughly.
 •Using the applicator (cotton swab), apply a layer of ointment to sufficiently cover each wart.
 •Allow it to fully absorb and cover it with a bandage as needed.
 •Repeat this procedure once or twice daily as needed (until the wart is removed) for up to 12 weeks.

OTHER INFORMATION:
 •Store between 68°F to 77°F (20°C to 25°C).
 Protect from excessive heat above 104°F (40°C).
 See a doctor as you may have a more serious skin condition.

INACTIVE INGREDIENT
 Shard paraffin, lanolin, phenol.

MAXIMUM STRENGTH
TAG & WART REMOVER
 FAST ACTING LIQUID
 15 ml/0.5 fl. oz.
 NATURAL FPRMULA


DURG FACTS
 ACTIVE INGREDIENT
 Salicylic Acid..... 5.0%

USES
 •To remove warts such as planter wart, flat wart, common wart and corns, calluses.
 •The planter wart is recognized by its location only on the bottom of the foot its tenderness, and the interruption of the footprint pattern.

WARNINGS
 •The liquid can get easily volatilized & crystallized, tighten the cap after use.
 •Avoid long-term contact with air while using.
 •For external use only, don't contact with eyes or swallow. Children should use it under the supervision of adults.

DO NOT USE
 •On damaged skin (cuts, abrasions, eczema, sunburn, etc.) or on any of the ingredients in this product.
 •If you are pregnant or breastfeeding.
KEEP OUT OF REACH OF CHILDREN
 If swallowed, get medical help or contact a Poison Control Center right away.

MAXIMUM STRENGTH
TAG & WART REMOVER
 FAST ACTING LIQUID
 15 ml/0.5 fl. oz.
 NATURAL FPRMULA

MADE IN CHINA


MAXIMUM STRENGTH WART REMOVER

maximum strength wart remover liquid

Product Information

| | | | |
|--------------------------------|----------------|---------------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:84023-113 |
| Route of Administration | TOPICAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|---|-------------------|----------------|
| SALICYLIC ACID (UNII: O414PZ4LPZ) (SALICYLIC ACID - UNII:O414PZ4LPZ) | SALICYLIC ACID | 5 mg in 100 mL |

Inactive Ingredients

| Ingredient Name | Strength |
|-----------------------------------|----------|
| LANOLIN (UNII: 7EV65EAW6H) | |
| PHENOL (UNII: 339NCG44TV) | |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|--|----------------------|--------------------|
| 1 | NDC:84023-113-01 | 15 mL in 1 BOTTLE; Type 0: Not a Combination Product | 12/30/2024 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|--------------------|--|----------------------|--------------------|
| OTC Monograph Drug | M028 | 12/30/2024 | |

Labeler - Shenzhen Yangan Technology Co., Ltd. (419283765)

Establishment

| Name | Address | ID/FEI | Business Operations |
|--------------------------------------|---------|-----------|------------------------|
| Shenzhen Yangan Technology Co., Ltd. | | 419283765 | manufacture(84023-113) |

Revised: 12/2024

Shenzhen Yangan Technology Co., Ltd.