

ASPIRIN 325 MG EC- aspirin tablet
Reliable 1 Laboratories LLC

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

ASPIRIN 325 MG EC TABLET

Drug Facts

Aspirin USP, 325 mg (NSAID)*

*nonsteroidal anti-inflammatory drug

Pain Reliever/Fever Reducer

temporarily relieves

- headache
- menstrual pain
- minor pain of arthritis
- muscle pain
- toothache
- pain and fever of colds

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include

- hives
- facial swelling
- asthma (wheezing)
- shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (ibuprofen, naproxen, or others)
- take more or for a longer time than directed
- have 3 or more alcoholic drinks every day while using this product

Drug Facts continued on back of label

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

Ask a doctor or pharmacist before use if you are

- taking a prescription drug for diabetes, gout or arthritis
- under a doctor's care for any serious condition
- taking any other drug

Stop use and ask a doctor if

- you experience any of the following signs of stomach bleeding:
 - feel faint
 - vomit blood
 - have bloody or black stools
 - have stomach pain that does not get better
 - pain gets worse or lasts more than 10 days
 - fever gets worse or lasts more than 3 days
 - redness or swelling is present in the painful area
 - any new symptoms appear
 - ringing in the ears or loss of hearing occurs

These could be signs of a serious condition

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

DRINK A FULL GLASS OF WATER WITH EACH DOSE

- **adults and children 12 years and over:** 1-2 tablets every 4 hours while symptoms last, not more than 12 tablets in 24 hours
- **children under 12 years:** ask a doctor

Other information

- store at room temperature 15°-30°C (59°-86°F)

Inactive ingredients

croscarmellose sodium, D&C yellow #10 (Al-lake), FD&C yellow #6 (Al-lake), HPMC, methacrylic acid copolymer, polyethylene glycol, propylene glycol, povidone, sodium lauryl sulfate, starch, stearic acid, talc, titanium dioxide

Questions or comments?

Questions or comments ? call **516-341-0666**, 8:30 am - 4:30 pm ET, Monday - Friday

TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

*Reliable 1 Laboratories LLC is not affiliated with the owner of the trademark Ecotrin®.

Distributed by: **Reliable 1 Laboratories LLC, Valley Stream, NY 11580**

www.reliable1labs.com

NDC 69618-015-01



**Enteric Coated
Aspirin USP
325 mg (NSAID)**

**PAIN RELIEVER
FEVER REDUCER**

100 TABLETS

†Compare to Active Ingredient in Ecotrin®

Drug Facts

Active ingredient (in each tablet) **Purpose**
Aspirin USP, 325 mg (NSAID)* Pain Reliever/Fever Reducer
*nonsteroidal anti-inflammatory drug

Uses temporarily relieves ■ headache ■ menstrual pain
■ minor pain of arthritis ■ muscle pain ■ toothache
■ pain and fever of colds

Warnings
Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.
Allergy alert: Aspirin may cause a severe allergic reaction which may include ■ hives ■ asthma (wheezing) ■ facial swelling ■ shock

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding.
The chance is higher if you
■ are age 60 or older
■ have had stomach ulcers or bleeding problems
■ take a blood thinning (anticoagulant) or steroid drug
■ take other drugs containing prescription or nonprescription NSAIDs (ibuprofen, naproxen, or others)
■ take more or for a longer time than directed
■ have 3 or more alcoholic drinks every day while using this product

Do not use if you have ever had an allergic reaction to any other pain reliever/fever reducer

Drug Facts continued on back of label
TAMPER EVIDENT: DO NOT USE IF IMPRINTED SAFETY SEAL UNDER CAP IS BROKEN OR MISSING

Reliable 1 Laboratories LLC is not affiliated with the owner of the trademark Ecotrin®.
Distributed by: **Reliable 1 Laboratories LLC, Valley Stream, NY 11580**
www.reliable1labs.com



Rev. 0715
Lot No.:
Exp. Date:
PEEL HERE FOR MORE DRUG FACTS

Drug Facts (continued)

Ask a doctor before use if
■ stomach bleeding warning applies to you
■ you have a history of stomach problems, such as heartburn
■ you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
■ you are taking a diuretic ■ you have asthma

Ask a doctor or pharmacist before use if you are
■ taking a prescription drug for diabetes, gout or arthritis
■ under a doctor's care for any serious condition
■ taking any other drug

Stop use and ask a doctor if
■ you experience any of the following signs of stomach bleeding: ■ feel faint ■ vomit blood
■ have bloody or black stools
■ pain gets worse or lasts more than 10 days
■ fever gets worse or lasts more than 3 days
■ redness or swelling is present in the painful area
■ any new symptoms appear
■ ringing in the ears or loss of hearing occurs

These could be signs of a serious condition

If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions DRINK A FULL GLASS OF WATER WITH EACH DOSE ■ **adults and children 12 years and over:** 1-2 tablets every 4 hours while symptoms last; not more than 12 tablets in 24 hours
■ **children under 12 years:** ask a doctor

Other information ■ store at room temperature 15°-30°C (59°-86°F)

Inactive ingredients croscarmellose sodium, D&C yellow #10 (Al-lake), FD&C yellow #6 (Al-lake), HPMC, methacrylic acid copolymer, polyethylene glycol, propylene glycol, povidone, sodium lauryl sulfate, starch, stearic acid, talc, titanium dioxide

Questions or comments? call 516-341-0666, 8:30 am - 4:30 pm ET, Monday - Friday

NDC 69618-015-01

Enteric Coated

Aspirin USP 325 mg (NSAID)

PAIN RELIEVER

FEVER REDUCER

100 TABLETS

*Compare to Active Ingredients in Ecotrin®

ASPIRIN 325 MG EC

aspirin tablet

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:69618-015

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	325 mg

Inactive Ingredients

Ingredient Name	Strength
POVIDONE (UNII: FZ989GH94E)	
METHACRYLIC ACID (UNII: 1CS02G8656)	
STEARIC ACID (UNII: 4ELV7Z65AP)	
STARCH, CORN (UNII: O8232NY3SJ)	
CROSCARMELOSE SODIUM (UNII: M28OL1HH48)	
FD&C YELLOW NO. 6 (UNII: H77VEI93A8)	
D&C YELLOW NO. 10 (UNII: 35SW5USQ3G)	
HYPROMELLOSES (UNII: 3NXW29V3WO)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)	
TALC (UNII: 7SEV7J4R1U)	
SODIUM LAURYL SULFATE (UNII: 368GB5141J)	
POLYETHYLENE GLYCOL 1000 (UNII: U076Q6Q621)	

Product Characteristics

Color	orange	Score	no score
Shape	ROUND	Size	10mm
Flavor		Imprint Code	AP;110
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69618-015-10	1000 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/01/2015	
2	NDC:69618-015-01	100 in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product	11/01/2015	



ASPIRIN

325mg EC Tablet

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part343	11/01/2015	

Labeler - Reliable 1 Laboratories LLC (079718111)

Registrant - Reliable 1 Laboratories LLC (079718111)

Revised: 3/2023

Reliable 1 Laboratories LLC