

**EQUATE DAYTIME COLD AND FLU- acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride solution**  
**Wal-Mart Stores Inc**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Wal-Mart Daytime Cold & Flu Drug Facts**

**Active ingredients (in each 15 mL)**

Acetaminophen 325 mg

Dextromethorphan HBr 10 mg

Phenylephrine HCl 5 mg

**Purpose**

Pain reliever/fever reducer

Cough suppressant

Nasal decongestant

**Uses**

- temporarily relieves common cold/flu symptoms:
- nasal congestion
- cough due to minor throat and bronchial irritation
- sore throat
- headache
- minor aches and pains
- fever

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if

- adult takes more than 4,000 mg of acetaminophen in 24 hours
- child takes more than 5 doses in 24 hours
- taken with other drugs containing acetaminophen
- adult has 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening

- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

### **Ask a doctor or pharmacist before use if you are**

taking the blood thinning drug warfarin

### **When using this product**

**do not use more than directed**

### **Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults)
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

### **Directions**

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
children 6 to under 12 yrs	15 mL every 4 hrs
children 4 to under 6 yrs	ask a doctor
children under 4 yrs	do not use

### **Other information**

- **each 15 mL contains:** sodium 7 mg
- store at 20-25°C (68-77°F)

### **Inactive ingredients**

butylated hydroxyanisole, edetate disodium, FD&C yellow no. 6, flavor, glycerin, menthol, monobasic sodium phosphate, polyethylene glycol, propylene glycol, purified water, saccharin sodium, sucrose, xanthan gum

### **Questions?**

**1-888-287-1915**

### **Package/Label Principal Display Panel**

equate™

Compare to Vicks® DayQuil® Cold & Flu active ingredients

NON-DROWSY

DAYTIME

ALCOHOL-FREE

Cold & Flu

Pain Reliever, Fever Reducer, Cough Suppressant, Nasal Decongestant

## Multi-Symptom Relief

- Acetaminophen - Aches, fever
- Dextromethorphan HBr - Cough
- Phenylephrine HCl - Nasal congestion

Antihistamine-Free

12 FL OZ (355mL)

NDC 49335-725-40

**equate™**

Compare to  
Vicks® DayQuil®  
Cold & Flu active  
ingredients\*



**NON-DROWSY**  
**ALCOHOL FREE**

# Cold & Flu

Pain Reliever, Fever Reducer, Cough Suppressant, Nasal Decongestant

## Multi-Symptom Relief

- **Acetaminophen** - Aches, fever
- **Dextromethorphan HBr** - Cough
- **Phenylephrine HCl** - Nasal congestion

Antihistamine-Free

**12 FL OZ (355mL)**

: 65640 2E F5

**DO NOT USE IF PRINTED NECKBAND IS  
BROKEN OR MISSING**

164152

## PARENTS:

Learn about teen medicine abuse

[www.StopMedicineAbuse.org](http://www.StopMedicineAbuse.org)

### Drug Facts

Active ingredients (in each 15 mL)	Purpose
Acetaminophen 325 mg.....	Pain reliever/ fever reducer
Dextromethorphan HBr 10 mg.....	Cough suppressant
Phenylephrine HCl 5 mg.....	Nasal decongestant

**Uses** ■ temporarily relieves common cold/flu symptoms:  
 ■ cough due to minor throat and bronchial irritation  
 ■ nasal congestion ■ minor aches and pains  
 ■ sore throat ■ headache ■ fever

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if ■ adult takes more than 4,000 mg of acetaminophen in 24 hours ■ child takes more than 5 doses in 24 hours ■ taken with other drugs containing acetaminophen ■ adult has 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening  
 ■ blisters ■ rash  
 If a skin reaction occurs, stop use and seek medical help right away.

**PEEL BACK AT  
CORNER FOR MORE  
INFORMATION**

: 65640 2E B1

### Drug Facts (continued)

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.  
 ■ if you have ever had an allergic reaction to this product or any of its ingredients

ADHESIVE AREA  
NO VARNISH • NO TYPE

### Drug Facts (continued)

Stop use and ask a doctor if ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion or cough gets worse or lasts more than 5 days (children) or 7 days (adults) ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.



If pregnant or breast-feeding, ask a health professional before use.

**Keep out of reach of children. Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** ■ take only as directed - see Overdose warning ■ only use the dose cup provided ■ do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 6 hrs
children 6 to under 12 yrs	15 mL every 4 hrs

ADHESIVE AREA  
NO VARNISH • NO TYPE



## EQUATE DAYTIME COLD AND FLU

acetaminophen, dextromethorphan hydrobromide, phenylephrine hydrochloride solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:49035-726
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	325 mg in 15 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 15 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 15 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>BUTYLATED HYDROXYANISOLE</b> (UNII: REK4960K2U)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>FD&amp;C YELLOW NO. 6</b> (UNII: H77VEI93A8)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>MENTHOL, UNSPECIFIED FORM</b> (UNII: L7T10EIP3A)	
<b>SODIUM PHOSPHATE, MONOBASIC, UNSPECIFIED FORM</b> (UNII: 3980JIH2SW)	
<b>POLYETHYLENE GLYCOL, UNSPECIFIED</b> (UNII: 3WJQ0SDWL1A)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SUCROSE</b> (UNII: C151H8M554)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

## Product Characteristics

<b>Color</b>	ORANGE (clear)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	MENTHOL (with fruit)	<b>Imprint Code</b>	
<b>Contains</b>			

## Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:49035-726-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	09/24/2019	

## Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part341	09/24/2019	

**Labeler** - Wal-Mart Stores Inc (051957769)

Revised: 10/2022

Wal-Mart Stores Inc