

**PROJEL-20- benzocaine gel**  
**Septodont, Inc.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Projel-20 Bubblegum**

**Active Ingredient:**

Benzocaine 20%

**Inactive Ingredients:**

Flavor, polyethylene glycol, sodium saccharin.

**Purpose:**

Oral Anesthetic intended for adults and children 2 years of age and older.

**Use:**

Projel-20<sup>®</sup> is a 20% benzocaine topical anesthetic gel packaged in 60 gram jars for use in the operatory. The gel is indicated for temporary relief of pain associated with minor irritation or injury in the mouth and gums or of pain due to canker sores, denture sores or orthodontic appliances. ProJel-20<sup>®</sup> is available in four flavors: Cherry, Pina Colada, Bubblegum and Mint.

**WARNINGS**

**Methemoglobinemia warning:**

Use of this product may cause methemoglobinemia, a serious condition that must be treated promptly because it reduces the amount of oxygen carried in blood. This can occur even if you have used this product before. Stop use and seek immediate medical attention if you or a child in your care develops:

- pale, gray, or blue colored skin (cyanosis)
- headache
- rapid heart rate
- shortness of breath
- dizziness or lightheadedness
- fatigue or lack of energy

**Allergy alert: Do not use on patients with history of allergy due to local anesthetics such as procaine, butacaine, benzocaine or other “caine” anesthetics.**

**DO NOT USE**

- For more than 7 days unless directed by a dentist or physician
- If you have a history of allergy to local anesthetics such as procaine, butacaine, benzocaine or other ‘caine’ anesthetics
- **For teething**
- **In children under 2 years of age**

## **WHEN USING THIS PRODUCT**

- Do not exceed recommended dosage
- Patient should not ingest or swallow
- Avoid contact with eyes
- Localized allergic reactions may occur after prolonged or repeated use
- Discontinue use if irritation/inflammation or fever and infection develops

## **STOP USE AND ASK A DOCTOR IF**

- Sore mouth symptoms do not improve in 7 days
- Swelling, rash or fever develops
- Irritation, pain, or redness persists or worsens.

**Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.**

If more than used for pain relief is accidentally swallowed, get medical help or contact a Poison Control Center immediately.

## **Directions:**

Apply only amount needed to the oral mucosa to prevent or relieve pain.  
Children under 2 years of age: do not use.

## **OTHER INFORMATION**

For professional use only in the practice of dentistry.  
Store at 20°C - 25°C (68°F - 77°F). Do not permit to freeze.

## **HOW SUPPLIED**

Projel-20<sup>®</sup>, Benzocaine 20% is available in in 60 gram jars (NDC 0362-0124-05).

Manufactured by:  
SEPTODONT, Inc.  
416 S. Taylor Ave  
Louisville, CO 80027  
800-872-8305 |  
www.septodontusa.com

Rev. 10/2018

**Bubblegum Projel Label**

Bubblegum

septodont

Projel-20®

Oral Anesthetic  
Net wt: 2.1 oz (60 g)



Manufactured by:  
SEPTODONT, Inc.  
416 S. Taylor Ave | Louisville, CO 80027  
800-872-8305 | www.septodontusa.com

For professional use only  
in the practice of dentistry

**WARNING:** Patient should not ingest  
or swallow. Avoid contact with eyes.  
Keep out of each of children.

**Active ingredient: benzocaine 20%**  
**Inactive ingredients:** flavor,  
polyethylene glycol, sodium saccharin.  
**For oral mucosal use only**

NDC 0362-0124-05

Peel  
Back

**PEEL BACK** to review  
drug facts  
Store at 20°- 25°C  
(68°F - 77°F).

**Do Not Freeze.**  
**KEEP TIGHTLY CLOSED.**

Rev 10/2018  
C-LABP-1805-C

## PROJEL-20

benzocaine gel

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:0362-0124
Route of Administration	TOPICAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BENZOCAINE (UNII: U3RS Y48JW5) (BENZOCAINE - UNII:U3RS Y48JW5)	BENZOCAINE	20 g in 100 g

### Inactive Ingredients

Ingredient Name	Strength
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
POLYETHYLENE GLYCOL 3350 (UNII: G2M7P15E5P)	
POLYETHYLENE GLYCOL 300 (UNII: 5655G9Y8AQ)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0362-0124-05	60 g in 1 JAR; Type 0: Not a Combination Product	12/15/1989	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph final	part346	12/15/1989	08/30/2020

**Labeler** - Septodont, Inc. (627058738)

### Establishment

Name	Address	ID/FEI	Business Operations
Septodont Inc.		012925513	manufacture(0362-0124) , label(0362-0124)

