HAIR REGEN HAIR REGROWTH TREATMENT- womens liquid minoxidil 6pk liquid YIWU YIZHOU TRADING CO.,LTD

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

84232-004

Active Ingredient(s)

Minoxidil 2% w/v

Purpose

HAIR REGROWTH TREATMENT

Use

to regrow hair on top of scalp (vertex only, see pictures on side of box)

Warnings

For external use only. For use by women Flammable: Keep away from fire or flame

Do not use

- ●Your degree of hair loss is different than that shown on the side of this carton, because this product may not work for you
- ●You have no family history of hair loss.
- ●Your hair loss is sudden and/or patchy.
- ●You do not know the reason for your hair loss.
- ●You are under 18 years of age. Do not use on babies and children.
- ●Your scalp is red, inflamed, infected, irritated, or painful.
- •You use other medicines on the scalp.

When using this product.

- ●Do not apply on other parts of your body.
- Avoid contact with the eyes. In case of accidental contact, rinse eyes with cool tap water.
- •Some people have experienced changes in hair color and/or texture.
- ●It takes time to regrow hair, results may occur at 2 months with twice a day usage. For some women, you may need to use this product for at least 4 months before you see results.
- ●The amount of hair regrowth is different for each person. This product will not work on all women.

Stop use

- Chest pain, rapid heartbeat, faintness, or dizziness occurs
- •Sudden, unexplained weight gain occurs
- Your hands or feet swell
- Scalp rritation or redness occurs
- Unwanted facial hair growth occurs
- ●You do not see hair growth in 4 months
- •May be harmful if used when pregnant or breast-feeding.
- ●Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. .

Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away.

Directions

- ●Apply one mL. with dropper 2 times a day directly onto the scalp in the hair loss area.
- Using more or more often will not improve resufts.
- •continued use is necessary to increase and keep your hair regrowth, or hair loss will begin again.

Other information

- •See hair loss pictures on side of box.
- •Before use, read all information on box and enclosed instructions.
- ■Keep the box, it contains important information.
- Hair regrowth has not been shown to last longer than 48 weeks in large clinical trials with continuous treatment with minoxidil topical solution 2% for women.
- ●In clinical studies of mostly white women aged 18-45 with mild to moderate hair loss, 19% showed moderate hair regrowth after 8 months of using 2% minoxidil topical solution, compared to 7% with a placebo. Additionally, 40% had minimal regrowth with minoxidil, while 33% saw minimal regrowth with the placebo.
- ●Store at 20° to 25°C (68° to 77°F). Keep cap tightly closed.

Inactive ingredients

alcohol, propylene glycol, purified water.

Package Label - Principal Display Panel

Working Title: MINOXIDIL LIQUID FOR WOMAN - Bottle Label - Version 19C



Font: Helvetica Neue LT Com 77 Bold Condensed

Font: Helvetica Neue LT Std 57 Condensed



HA(R REGEN Extra Strength For Women MINOXIDIL

HAIR REGROWTH TREATMENT **TOPICAL SOLUTION USP. 2%**

MONTH / SUPPLY

Read and keep carton and leaflet for full product information

Active ingredient Minoxidil 2% w/v

Warnings For external use only Flammable: Keep away from fire or flame

- Do not use if

 you have no family history of hair loss
 your hair loss is sudden and/or patchy
 you hair loss is associated with childhirith
 you do not know the reason for your hair loss
 you are under 18 years of age. Do not use on babies and children.
 your scalp is red, inflamed, inflocted, inritated, or painful
 you use other medicines on the scalp

Ask a doctor before use if you have heart disea

- Ask a occur before use if you have near crisease

 When using this product

 do not apply to other parts of the body

 avoid contact with the eyes. In case of accidental contact, rinse
 eyes with large amounts of cool tap water

 some people have experienced changes in hair color and/or texture

- Stop use and ask a doctor if m chest pain, rapid heartbeat, faintness, or dizziness occurs

- sudden, unexplained weight gain occurs
 your hands or feet swell
 scalp irritation or redness occurs
 unwanted facial hair growth occurs

May be harmful if used when pregnant or breast-feeding. Keep out of reach of children. If swallowed, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

- apply one mL 2 times a day directly onto the scalp in the hair loss area see carton and enclosed leaflet for complete directions

- using more or more often will not improve results
 continued use is necessary or hair loss will begin again

Inactive ingredients alcohol, propylene glycol, purified water

Store at 20° to 25°C (68° to 77°F) Keep tightly closed

HAIR REGEN

Distributed by HA(R REGEN®

167 Madison Avenue Ste 205 #236 New York City, NY 10016

LOT NO. WHD972 EXP. 12/2028



- Font: Myriad Pro Semibold Live for changes
- · Font attached with file





HONG IN USED TO THE STATE OF TH

Al visition
In this absolute and made above summer aged
The fit such militim machines have been. Will
also explain machines have been such as of
any pill machines and applied above amountaines.
The such a facetor, belief standard, with the observed
approvision and standard and all the observed
approvision and standard and approvision and
approvision and standard and approvision and
approvision and standard and approvision and
approvision and approvision and
approvision and approvision and
approvision and
approvision and
approximately approximately approximately
approvision and
approximately
approx

4455 E. Nunneley Rd, #101 Gilbert, AZ, 85296

CONFIDENTIAL
Copyright Material

HAIR REGEN HAIR REGROWTH TREATMENT

womens liquid minoxidil 6pk liquid

Product Information

Product Type HUMAN OTC DRUG Item Code (Source) NDC:84232-004

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name
Basis of Strength
MINOXIDIL (UNII: 5965120SH1) (MINOXIDIL - UNII:5965120SH1)

MINOXIDIL
2 g in 100 mL

Inactive Ingredients

Ingredient Name

ALCOHOL (UNII: 3K9958V90M)

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)

WATER (UNII: 059QF0K00R)

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date		
1	NDC:84232-004- 01	60 mL in 1 BOTTLE; Type 0: Not a Combination Product	01/08/2025			
Marketing Information						
	Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date		
	approved drug ner		01/03/2025			

Labeler - YIWU YIZHOU TRADING CO.,LTD (840905001)

Establishment						
Name	Address	ID/FEI	Business Operations			
YIWU YIZHOU TRADING CO.,LTD		840905001	manufacture(84232-004)			

Revised: 1/2025 YIWU YIZHOU TRADING CO.,LTD