

**DENTEMP CANKER COVER- canker cover tablet, extended release
DOC Brands**

Dentemp Canker Cover 2ct

Menthol 2.5 mg

Oral Pain Reliever

For temporary relief of pain associated with Canker Sores

Do not use this product for more than 7 days unless directed by a dentist or doctor. Stop use and ask a dentist or doctor if • sore mouth symptoms do not improve in 7 days • irritation, pain or redness worsens • swelling, rash or fever develops.

Do not exceed recommended dosage.

If pregnant or breastfeeding or taking prescription drugs, ask a health care professional before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

adults and children 5 years of age and older, apply up to 3 tablets a day, as needed. Place the tablet on a clean, dry finger with the white side up. Place the white side on the sore and hold in place for 10 seconds. If sore is difficult to reach (in the fold between the cheek and gum or near the teeth or lip) break the tablet along the score and use half. Some discomfort may occur during the first few minutes, but will quickly subside, followed by hours of soothing relief. Within 30 minutes, the tablet forms a clear, gel-like bandage that seals and protects the sore for hours before dissolving. Do not remove the tablet before it dissolves. In case of discomfort, the tablet may be removed by gently peeling the tablet from the sides while washing with warm water. Do not use any instrument to remove the tablet. • children under 5 years, ask a doctor.

store in a cool, dry place.

annatto, carbomer homopolymer type a, citrus oil, hydroxypropyl cellulose, magnesium chloride, povidone K30, povidone K90, silicon dioxide, xylitol

833-362-2763 or consumeraffairs@doc-brands.com

**NEW
LOOK**



Dentemp®

**CLINICALLY PROVEN TO
HELP HEAL AS FAST AS 1 DAY**

**CANKER SORES
MOUTH SORES
ORAL ULCERS
IRRITATION
FROM BRACES**

canker cover®

MEDICATED ORAL PAIN RELIEVER

**HELPS HEAL AS FAST AS 1 DAY
RELIEVES PAIN & HEALS SORES
STAYS ON WHILE EATING
UP TO 12 HOURS OF SOOTHING RELIEF**



**PLACE WHITE
SIDE DIRECTLY
ON SORE**

**DUAL-LAYER
DESIGN**

**MEDICATED
LAYER
RELIEVES PAIN
& PROMOTES
HEALING**

**BIOADHESIVE
LAYER FORMS
PROTECTIVE
BARRIER &
STAYS ON
WHILE EATING**

2 COMPLETE TREATMENTS

*Clinical on File
U.S. Pat. No. 7,943,169
www.dentemp.com
©2025 Manuf. for DOC Brands, Inc.
Knoxville, TN 37932 USA
Made in USA | 213.007A



Drug Facts

Active ingredient (per each tablet)
Menthol 2.5 mg

Purpose
Oral pain reliever

Indications for the temporary relief of pain associated with canker sores.

Warnings
Do not use this product for more than 7 days unless directed by a dentist or doctor.
Stop use and ask a dentist or doctor if some mouth symptoms do not improve in 7 days • Irritation, pain or redness worsens • Swelling, rash or fever develops
Do not exceed recommended dosage.

If pregnant or breastfeeding or taking prescription drugs, ask a health care professional before use. Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions • adults and children 5 years of age and older, apply up to 3 tablets a day, as needed. Place the tablet on a clean, dry finger with the white side up. Place the white side on the sore and hold in place for 10 seconds. If sore is difficult to reach (in the fold) between the cheek and gum or near the teeth or lip) break the tablet along the score and use half. Some discomfort may occur during the first few minutes, but will quickly subside. Followed by hours of soothing relief. Within 30 minutes, the tablet forms a clear gel-like bandage that seals and protects the sore for hours before dissolving. Do not remove the tablet before it dissolves. In case of discomfort, the tablet may be removed by gently peeling the tablet from the sides while washing with warm water. Do not use any instrument to remove the tablet. • children under 5 years, ask a doctor.

Other information store in a cool dry place.

Inactive ingredients amato, carbomer homopolymer type a, citrus oil, hydroxypropyl cellulose, magnesium chloride, povidone K30, povidone K90, silicon dioxide, xylitol

Questions or comments? 833-362-2763 or consumeraffairs@doc-brands.com

DENTEMP CANKER COVER

canker cover tablet, extended release

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:73653-213
Route of Administration	TOPICAL, ORAL		

Active Ingredient/Active Moiety				
Ingredient Name		Basis of Strength	Strength	
MENTHOL (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)		MENTHOL	2.5 mg	
Inactive Ingredients				
Ingredient Name			Strength	
MAGNESIUM CHLORIDE (UNII: 02F3473H9O)				
XYLITOL (UNII: VCQ006KQ1E)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
POVIDONE K90 (UNII: RDH86HJV5Z)				
CARBOMER HOMOPOLYMER TYPE A (UNII: F68VH75CJC)				
CITRUS LIMON FRUIT OIL (UNII: 0HNC1J1YED)				
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)				
ANNATTO (UNII: 6PQP1V1B6O)				
POVIDONE K30 (UNII: U725QWY32X)				
Product Characteristics				
Color	pink	Score	no score	
Shape	ROUND	Size	10mm	
Flavor	MINT	Imprint Code		
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:73653-213-02	2 in 1 CARTON	02/01/2025	
1	NDC:73653-213-01	1 in 1 BLISTER PACK; Type 0: Not a Combination Product		
Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M022	01/09/2025		

Labeler - DOC Brands (081254601)