DENTEMP CANKER COVER- canker cover tablet, extended release DOC Brands

Dentemp Canker Cover 2ct

Menthol 2.5 mg

Oral Pain Reliever

For temporary relief of pain associated with Canker Sores

Do not use this product for more than 7 days unless directed by a dentist or doctor. Stop use and ask a dentist or doctor if • sore mouth symptoms do not improve in 7 days • irritation, pain or redness worsens • swelling, rash or fever develops.

Do not exceed recommended dosage.

If pregnant or breastfeeding or taking prescription drugs, ask a health care professional before use.

Keep out of reach of children. In case of accidental overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

adults and children 5 years of age and older, apply up to 3 tablets a day, as needed. Place the tablet on a clean, dry finger with the white side up. Place the white side on the sore and hold in place for 10 seconds. If sore is difficult to reach (in the fold between the cheek and gum or near the teeth or lip) break the tablet along the score and use half. Some discomfort may occur during the first few minutes, but will quickly subside, followed by hours of soothing relief. Within 30 minutes, the tablet forms a clear, gel-like bandage that seals and protects the sore for hours before dissolving. Do not remove the tablet before it dissolves. In case of discomfort, the tablet may be removed by gently peeling the tablet from the sides while washing with warm water. Do not use any instrument to remove the tablet. • children under 5 years, ask a doctor.

store in a cool, dry place.

annatto, carbomer homopolymer type a, citrus oil, hydroxypropyl cellulose, magnesium chloride, povidone K30, povidone K90, silicon dioxide, xylitol

833-362-2763 or consumeraffairs@doc-brands.com



NEW!

Healing Process:

Before

Application

Dual Layer Design: Medicated Layer Promotes Healing Bioadhesive Layer Forms Protective Barrier

Helps Heal as Fast as 1 day:

- **OCANKER Sores**
- Mouth Sores
- Oral Ulcers
- Mirritation from Braces



sucrose. glucose preservatives



Dentemp' Canker Cover

Oral Pain Reliever

CLINICALLY PROVEN*

- Helps Heal as Fast as 1 Day 🏈 Relieves Pain & Heals Sores Up to 12 Hours of Soothing Relief
- Stays on While Eating



*Clinical on File U.S. Pat. No. 7,943,169 www.dentemp.com ©2023 Manuf. for DOC Brands, Inc. Wayne, PA 19087 USA Made in USA | 213.007



TECHNO!



Questions or comments? 833-362-2763 or consumeraffairs@doc-brands.com

cellulose, magnesium chloride, povidone K3O, povidone K9O, silicon dioxide, xylitol Inactive ingredients annatto, carbomer homopolymer type a, citrus oil, hydroxypropyl

Other Information store in a cool, dry place.

any instrument to remove the tablet. • children under 5 years, ask a doctor. bre soon and gnotes takes the teeth of places, the tables alone and before and before and before and before and before the core and soon are alone as the second by the first two minutes, but will quickly subside, followed by hours of soothing soller, but alone alone alone and protects the sore for hours before alsolving. Do not remove the tablet before it alsolves in case of discomfort, the tablet may be removed the tablet and protects the sore of discomfort, the tablet may be removed by gnully peeling the tablet from the sides will be washing with warm water. Do not use any peeling the tablet from the sides will be washing with warm stablet and to come at the place. needed. Place the tablet on a clean, dry finger with the white side up. Place the white side on the reach (in the fold of the sore is difficult to reach (in the fold of the sore is difficult to reach (in the fold of the sore is difficult to reach (in the fold of the sore). Directions - adults and children 5 years of age and older, apply up to 3 tablets a day, as

get medical help or contact a Polson Control Center (1-800-222-1222) right away If pregnant or breastfeeding or taking prescription drugs, ask a health care professional before use. Keep out of reach of children, in case of accidental overdose,

Do not exceed recommended dosage.

days + initation, pain or redness worsens • swelling, rash or fever develops Warnings

Not use this product for more than 7 days unless directed by a dentist or doctor.

Stop use and ask a dentist or doctor if • sore mouth symptoms do not improve in 7

Indications for the temporary relief of pain associated with canker sores.

Oral pain reliever psoding

Menthol 2.5 mg Active ingredient (per each tablet)

Drug Facts

DENTEMP CANKER COVER

canker cover tablet, extended release

Product Information

Product Type

HUMAN OTC DRUG

Item Code (Source)

NDC:73653-212

Active	Ingredient/Active	Moiety
--------	-------------------	--------

Ingredient Name	Basis of Strength	Strength
MENTHOL (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)	MENTHOL	2.5 mg

Inactive Ingredients			
Ingredient Name	Strength		
MAGNESIUM CHLORIDE (UNII: 02F3473H9O)			
XYLITOL (UNII: VCQ006KQ1E)			
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)			
POVIDONE K90 (UNII: RDH86HJV5Z)			
CARBOMER HOMOPOLYMER TYPE A (UNII: F68VH75CJC)			
CITRUS LIMON FRUIT OIL (UNII: 0HNC1J1YED)			
HYDROXYPROPYL CELLULOSE, UNSPECIFIED (UNII: 9XZ8H6N6OH)			
ANNATTO (UNII: 6PQP1V1B6O)			
POVIDONE K30 (UNII: U725OW/32X)			

Product Characteristics				
Color	pink	Score	no score	
Shape	ROUND	Size	10mm	
Flavor	MINT	Imprint Code		
Contains				

P	Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date	
1	NDC:73653-212- 07	2 in 1 CARTON	02/01/2025		
1	NDC:73653-212- 01	1 in 1 BLISTER PACK; Type 0: Not a Combination Product			

Marketing Information				
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date	
OTC Monograph Drug	M022	01/09/2025		

Labeler - DOC Brands (081254601)

Revised: 1/2025 DOC Brands