

**LIDOPRO PATCH (LIDOCAINE 4%, MENTHOL 1%)- lidocaine and menthol patch patch**  
**Clinic Pharma**

-----

**83881-402 LidoPro Patch (Lidocaine 4%, Menthol 1%)**

**Active ingredient**

Lidocaine 4%

Menthol 1%

**Purpose**

External Anesthetic

External Analgesic

**Uses**

For the temporary relief of pain.

**Warnings**

For external use only.

**Do not use**

- on the face or rashes, on wounds or damaged skin
- in the eyes, mouth, or other mucous membranes
- on genitals
- with a heating pad
- right before or after heart surgery
- any patch from a pouch that has been opened for 7 or more days
- in large quantities, particularly over raw surfaces or blistered areas
- if tamper-evident seal is torn, broken, or missing
- more than 2 patches per day unless directed by a doctor
- children under 18 years of age

**Ask a doctor before use if you have**

- allergies to topical products
- high blood pressure, heart disease, or kidney disease

**When using this product**

- Avoid contact with eyes. If eye contact occurs, rinse thoroughly with water.
- The risk of heart attack or stroke may increase if you use more than directed or for longer than directed.

**Stop use and ask doctor if**

condition worsens, or if symptoms persist for more than 7 days or clear up and occur again within a few days.

**If pregnant or breast-feeding, ask a health professional before use**

Do not use during last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

If put in mouth, seek medical help or contact a Poison Control Center right away. Dispose of the used patches by folding sticky ends together.

**Directions**

Adults 18 years and older:

- clean and dry affected area
- open pouch and remove one patch
- remove protective film from patch
- apply one patch to the affected area of pain and leave in place for 8 to 12 hours

**Other information**

- Some individuals may not experience pain relief until several minutes or hours after applying the patch
- avoid storing product in direct sunlight
- protect product from excessive moisture
- store at 67-77°F (19-25°C)

**Inactive ingredients**

Carboxymethylcellulose sodium, Dihydroxyaluminum aminoacetate, Edetate sodium, Glycerin, Kaolin, Methyl parahydroxybenzoate, Partially neutralized Sodium polyacrylate, Polyacrylic acid, Polysorbate 80, Propyl parahydroxybenzoate, Propylene glycol, Sodium Polyacrylate, Tartaric acid, Titanium dioxide, water

**Questions?**

Call toll-free (800) 224-2048



15  
patches

NDC 83881-402-15



Pain Relieving Patch

- ✓ Deep Penetrating
- ✓ Long-lasting
- ✓ Soothing

15 PATCHES  
10 cm x 12 cm  
(3.9" x 4.7")



<b>Drug Facts</b>	
<b>Active Ingredients (in each patch)</b>	<b>Purpose</b>
Lidocaine 4%	External Anesthetic
Menthol 1%	External Analgesic
<b>Uses</b>	
For the temporary relief of pain.	
<b>Warnings</b>	
For external use only.	
<b>Do not use</b> - on the face or rashes, on wounds or damaged skin - in the eyes, mouth, or other mucous membranes - on genitals - with a heating pad - right before or after heart surgery - any patch from a pouch that has been opened for 7 or more days - in large quantities, particularly over raw surfaces or blistered areas - if tamper-evident seal is torn, broken, or missing - more than 2 patches per day unless directed by a doctor - children under 18 years of age	
<b>Ask a doctor before use if you have</b> - allergies to topical products - high blood pressure, heart disease, or kidney disease.	
<b>When using this product</b> - avoid contact with eyes. If eye contact occurs, rinse thoroughly with water - the risk of heart attack or stroke may increase if you use more than directed or for longer than directed.	
<b>Stop use and ask doctor</b> if condition worsens, or if symptoms persist for more than 7 days or clear up and occur again within a few days.	
<b>If pregnant or breast-feeding, ask a health professional before use.</b> Do not use during last 3 months of pregnancy because it may cause problems in the unborn child or complications during delivery.	
<b>Keep out of reach of children.</b> If put in mouth, seek medical help or contact a Poison Control Center right away. Dispose of the used patches by folding sticky ends together.	
<b>Directions</b>	
<b>Adults 18 years and older:</b>	
- Clean and dry affected area	- If pain lasts after using the first patch, a second patch may be applied for up to another 8 to 12 hours
- open pouch and remove one patch	- only use one patch at a time
- remove protective film from patch	- wash hands with soap and water after applying or removing patch
- apply one patch to the affected area of pain and leave in place for 8 to 12 hours	- re-seal pouch containing unused patches after each use
<b>Other information</b>	
- some individuals may not experience pain relief until several minutes or hours after applying the patch	
- avoid storing product in direct sunlight - protect product from excessive moisture - store at 67-77°F (19-25°C)	
<b>Inactive ingredients</b>	
Carbonated polyethylene glycol, Dihydroxyaluminum aminoacetate, Edetate sodium, Glycerin, Kaolin, Methyl parahydroxybenzoate, Partially neutralized Sodium Polycrylate, Polycrylic acid, Polysorbate 80, Propyl parahydroxybenzoate, Propylene glycol, Sodium Polycrylate, Tartaric acid, Titanium dioxide, Water	
<b>Questions?</b>	
Call toll-free (800) 224-2048	

Formulated and Designed in Nevada

Made in EU



Patent Pending

Manufactured For:




Las Vegas, NV 89121

Lot:

Exp.:



3 83881 40215 7



## LIDOPRO PATCH (LIDOCAINE 4%, MENTHOL 1%)

lidocaine and menthol patch patch

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:83881-402
Route of Administration	TRANSDERMAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MENTHOL (UNII: L7T10EIP3A) (MENTHOL - UNII:L7T10EIP3A)	MENTHOL	84 mg in 8400 mg
LIDOCAINE (UNII: 98PI200987) (LIDOCAINE - UNII:98PI200987)	LIDOCAINE	336 mg in 8400 mg

### Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C0OX)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
SODIUM POLYACRYLATE (8000 MW) (UNII: 285CYO341L)	
TARTARIC ACID (UNII: W4888I119H)	

<b>WATER</b> (UNII: 059QF0KO0R)	
<b>DIHYDROXYALUMINUM AMINOACETATE</b> (UNII: DO250MG0W6)	
<b>POLYACRYLIC ACID (8000 MW)</b> (UNII: 73861X4K5F)	
<b>PROPYLPARABEN</b> (UNII: Z8IX2SC1OH)	
<b>SODIUM POLYACRYLATE (2500000 MW)</b> (UNII: 05I15JN12J)	
<b>KAOLIN</b> (UNII: 24H4NWX5CO)	
<b>EDETATE SODIUM</b> (UNII: MP1J8420LU)	
<b>TITANIUM DIOXIDE</b> (UNII: 15FIX9V2JP)	
<b>PROPANEDIOL</b> (UNII: 5965N8W85T)	
<b>CARBOXYMETHYLCELLULOSE SODIUM</b> (UNII: K679OBS311)	

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:83881-402-05	5 mg in 1 POUCH; Type 0: Not a Combination Product	01/09/2025	
2	NDC:83881-402-15	15 mg in 1 POUCH; Type 0: Not a Combination Product	01/09/2025	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M017	01/09/2025	

**Labeler** - Clinic Pharma (119158469)