

**CHILDRENS COLD AND COUGH - brompheniramine maleate,
dextromethorphan, phenylephrine hcl liquid
CVS**

Childrens Cold and Cough 577

ACTIVE INGREDIENTS (in each 10 mL)

Brompheniramine maleate, USP 2 mg

Dextromethorphan HBr, USP 10 mg

Phenylephrine HCl, USP 5 mg

PURPOSES

Antihistamine

Cough Suppressant

Nasal decongestant

USES

- temporarily relieves cough due to minor bronchial irritation as may occur with a cold, and nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily relieves these symptoms due to hay fever (allergic rhinitis):
 - sneezing
 - Itching of the nose or throat
 - runny nose
 - itchy, watery eyes
- temporarily restores freer breathing through the nose

WARNINGS

.

DO NOT USE

- to sedate a child or to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, or psychiatric, for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or

pharmacist before taking this product.

ASK A DOCTOR BEFORE USE IF YOU HAVE

- heart disease
- glaucoma
- diabetes
- thyroid disease
- high blood pressure
- difficulty in urination due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- a breathing problem or persistent or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis or emphysema

ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE

- taking any other oral nasal decongestant or stimulant
- taking sedative or tranquilizers

WHEN USING THIS PRODUCT

- **do not take more than directed**
- may cause drowsiness
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic beverages
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

STOP USE AND ASK A DOCTOR IF

- nervousness, dizziness or sleeplessness occur
- symptoms do not improve within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

IF PREGNANT OR BREAST-FEEDING,

ask a health professional before use.

KEEP OUT OF REACH OF CHILDREN.

In case of overdose, get medical help or contact a Poison Control Center right away.

DIRECTIONS

- do not take more than 6 doses in any 24-hour period
- measure only with dosing cup provided
- do not exceed recommended dosage
- mL = milliliter

Age	Dose
adults & children 12 years & over	20 mL every 4 hours
children 6 to under 12 years	10 mL every 4 hours
children under 6 years	do not use

OTHER INFORMATION

- **each 10 mL contains:** sodium 10 mg
- store at 20-25°C (68-77°F)
- measuring cup provided

INACTIVE INGREDIENTS

anhydrous citric acid, artificial flavor, FD&C blue #1, FD&C Red #40, glycerin, propylene glycol, propyl gallate, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose

PRINCIPAL DISPLAY PANEL

CVS

Health ®

Compare to the active ingredients in Children's Dimetapp® Cold & Cough*

NDC 51316-578-03

Children's

Cold & Cough

BROMPHENIRAMIINE MALEATE

Antihistamine

DEXTROMETHORPHAN HYDROBROMIDE

Cough Suppressant

PHENYLEPHRINE HYDROCHLORIDE

Nasal Decongestant

DM Liquid

- Relieves nasal congestion;
Stuffy nose; Runny nose;
Sneezes
- Soothes Cough
- Calm allergy symptoms

Alcohol free

6+ Years

Grape Flavor

Dosage Cup Included

4 FL OZ (118 mL)

Drug Facts

Active ingredients (in each 10 mL) Purposes
Brompheniramine maleate, USP 2 mg.....Antihistamine
Dextromethorphan HBr, USP 10 mg.....Cough suppressant
Phenylephrine HCl, USP 5 mg.....Nasal decongestant

Uses

- temporarily relieves cough due to minor bronchial irritation as may occur with a cold, and nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily relieves these symptoms due to hay fever (allergic rhinitis): sneezing itching of the nose or throat runny nose itchy, watery eyes
- temporarily restores freer breathing through the nose

Warnings

Do not use

- to sedate a child or to make a child sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

- heart disease ■ glaucoma ■ diabetes
- thyroid disease ■ high blood pressure
- difficulty in urination due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- a breathing problem or persistent or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are

- taking any other oral nasal decongestant or stimulant
- taking sedatives or tranquilizers

When using this product

- do not take more than directed
- may cause drowsiness
- alcohol, sedatives, and tranquilizers may increase drowsiness
- avoid alcoholic beverages
- be careful when driving a motor vehicle or operating machinery

Drug Facts (continued)

- excitability may occur, especially in children

Stop use and ask a doctor if

- nervousness, dizziness or sleeplessness occur
- symptoms do not improve within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.

Directions

- do not take more than 6 doses in any 24-hour period
- do not exceed recommended dosage
- mL = milliliter

Age	Dose
adults & children 12 years & over	20 mL every 4 hours
children 6 to under 12 years	10 mL every 4 hours
children under 6 years	do not use

Other information

- each 10 mL contains: sodium 10 mg
- store at 20-25°C (68-77°F)
- measuring cup provided

Inactive ingredients anhydrous citric acid, artificial flavor, FD&C blue #1, FD&C red #40, glycerin, propylene glycol, propyl gallate, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose

*This product is not manufactured or distributed by Pfizer Consumer Healthcare, the distributor of Children's Dimetapp® Cold & Cough.

KEEP THIS BOX FOR COMPLETE INSTRUCTIONS

DO NOT USE IF SEAL UNDER CAP IMPRINTED WITH "SEALED FOR YOUR PROTECTION" IS BROKEN OR MISSING.

CVS Health.

Children's Cold & Cough

DM Liquid

CVS Health.

Compare to the active ingredients in Children's Dimetapp® Cold & Cough*

CVS Health.

Children's Cold & Cough

BROMPHENIRAMINE MALEATE

Antihistamine

DEXTROMETHORPHAN HBr
Cough suppressant

PHENYLEPHRINE HCl
Nasal decongestant

DM Liquid

- Relieves nasal congestion; Stuffy nose; Runny nose; Sneezes
- Soothes cough
- Calms allergy symptoms

Alcohol free
6+ Years



Grape Flavor
Dosage Cup Included



4 FL OZ (118 mL)

Package Contains One Bottle

MFR# 53041 REV 0623

#477080



0 50428 36399 7

LOT:

EXP:

CVS Health® Children's Cold & Cough offers relief from nasal congestion, runny nose, coughing, and sneezing to help your child get through the day.

PARENTS:

Learn about teen medicine abuse
www.StopMedicineAbuse.org

Distributed by: CVS Pharmacy, Inc.
One CVS Drive, Woonsocket, RI 02895
© 2023 CVS/pharmacy
CVS.com® 1-800-SHOP CVS

V-15002

100% money back guaranteed.
CVS.com/returnpolicy



10
WestRock 1400735-1
12/24

Drug FactsDO NOT USE IF SEAL UNDER CAP
IMPRINTED WITH "SEALED FOR YOUR
PROTECTION" IS BROKEN OR MISSING.**Active ingredients (in each 10 mL)** **Purposes**
Brompheniramine maleate, USP 2 mg...Antihistamine
Dextromethorphan HBr, USP 10 mg...Cough suppressant
Phenylephrine HCl, USP 5 mg...Nasal decongestant**Uses** ■ temporarily relieves cough due to
minor bronchial irritation as may occur with
a cold, and nasal congestion due to the
common cold, hay fever or other upper
respiratory allergies ■ temporarily relieves
these symptoms due to hay fever (allergic
rhinitis): • sneezing • itching of the nose or
throat • runny nose • itchy, watery eyes
■ temporarily restores freer breathing
through the nose**Warnings****Do not use** ■ to sedate a child or to make
a child sleepy ■ if you are now taking a
prescription monoamine oxidase inhibitor
(MAOI) (certain drugs for depression,
psychiatric, or emotional conditions, or
Parkinson's◀ **Drug Facts** continued on other panel**Children's
Cold &
Cough****BROMPHENIRAMINE
MALEATE**
Antihistamine**DEXTROMETHORPHAN HBr**
Cough suppressant**PHENYLEPHRINE HCl**
Nasal decongestant
DM Liquid**Alcohol free**
6+ Years**4 FL OZ (118 mL)****Grape
Flavor****Drug Facts** (continued)disease), or for 2 weeks after stopping the MAOI drug. If you
do not know if your prescription drug contains an MAOI,
ask a doctor or pharmacist before taking this product.**Directions** ■ do not take more than 6 doses in any
24-hour period ■ do not exceed recommended dosage
■ mL = milliliter

Age	Dose
adults & children 12 years & over	20 mL every 4 hours
children 6 to under 12 years	10 mL every 4 hours
children under 6 years	do not use

Other information ■ each 10 mL contains: sodium 10
mg ■ store at 20-25°C (68-77°F) ■ measuring cup provided**Keep carton for complete information.**

MFR# 53041 REV 0623

Distributed by:
CVS Pharmacy, Inc.
One CVS Drive
Woonsocket, RI 02895
© 2023 CVS/pharmacy
CVS.com®
1-800-SHOP CVS
V-15002

LOT:

EXP:

CHILDRENS COLD AND COUGH

brompheniramine maleate, dextromethorphan, phenylephrine hcl liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:51316-578
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BROMPHENIRAMINE MALEATE (UNII: IXA7C9ZN03) (BROMPHENIRAMINE - UNII:H57G17P2FN)	BROMPHENIRAMINE MALEATE	2 mg in 10 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 10 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE	5 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
GLYCERIN (UNII: PDC6A3C0OX)	

PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SORBITOL (UNII: 506T60A25R)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
PROPYL GALLATE (UNII: 8D4SNN7V92)	
SODIUM CITRATE (UNII: 1Q73Q2JULR)	

Product Characteristics

Color	PURPLE	Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51316-578-03	1 in 1 CARTON	01/17/2025	
1		118 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	01/17/2025	

Labeler - CVS (062312574)

Establishment

Name	Address	ID/FEI	Business Operations
Guardian Drug Company		119210276	MANUFACTURE(51316-578)

Revised: 1/2025

CVS