

**DAYTIME COLD AND COUGH AND NIGHTTIME COLD AND CONGESTION CHILDRENS- brompheniramine maleate, dextromethorphan hbr, phenylephrine hcl, diphenhydramine hcl, phenylephrine hcl
Walgreens**

DRUG FACTS

Active ingredients for Nighttime (in each 10 mL)

Diphenhydramine HCl 12.5 mg

Phenylephrine HCl 5 mg

Active ingredients for Daytime (in each 10 mL)

Brompheniramine Maleate 2 mg

Dextromethorphan HBr 10mg

Phenylephrine HCl 5 mg

Purpose for Nighttime

Antihistamine / Cough suppressant

Nasal Decongestant

Purpose for Daytime

Antihistamine

Cough suppressant

Nasal decongestant

Uses

Nighttime

- temporarily relieves these symptoms occurring with a cold, hay fever, or other upper respiratory allergies
- nasal congestion
- cough
- runny nose
- sneezing
- itchy, watery eyes
- itching of the nose or throat

Daytime

- temporarily relieves cough due to minor throat and bronchial irritation occurring with a cold, and nasal congestion due to the common cold, hay fever or other upper respiratory allergies
- temporarily relieves these symptoms due to hay fever (allergic rhinitis)
 - runny nose
 - itchy, watery eyes
 - sneezing
 - itching of the nose or throat
- temporarily restores freer breathing through the nose

Warnings

Do not use

Nighttime

- to sedate a child or to make a child sleepy.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product
- with any other product containing diphenhydramine, even one used on skin.

Daytime

- to sedate a child or to make sleepy
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI)(certain drugs for depression,psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

Ask a doctor before use if you have

Nighttime

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged prostate gland
- cough that occurs with too much phlegm (mucus)
- a breathing problem or chronic cough that last as occurs with smoking,asthma, chronic bronchitis or emphysema

Daytime

- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- trouble urinating due to an enlarged gland

- cough that occurs with too much phlegm (mucus)
- a breathing problem or persistent or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis or emphysema

Ask a doctor or pharmacist before use if you are

Nighttime

- taking any other oral nasal decongestant or stimulant
- taking sedative or tranquilizers

Daytime

- taking any other oral nasal decongestant or stimulant
- taking sedative or tranquilizers

When using these products

Nighttime

- **do not use more than directed**
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedatives and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Daytime

- **do not use more than directed**
- marked drowsiness may occur
- avoid alcoholic drinks
- alcohol, sedative and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

Nighttime

- you get nervous, dizzy or sleepless
- symptoms do not get better within 7 days or occur with a fever
- cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition

Daytime

- nervousness, dizziness or sleeplessness occur
- symptoms do not improve within 7 days or are accompanied by fever
- cough lasts more than 7 days, comes back or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition

If pregnant or breast-feeding,

Nighttime

ask a health professional before use.

Daytime

ask a health professional before use

Keep out of reach of children.

Nighttime

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Daytime

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.

Directions

Nighttime

- do not take more than 6 doses in any 24 hours period
- do not exceed recommended dosage
- measure only with dosing cup provided. Do not use any other dosing device
- keep dosing cup with product
- mL = milliliter

Age	Dose
adults and children 12 years and over	20 mL every 4 hours
children 6 to under 12 years	10 mL every 4 hours
children under 6 years	do not use

Daytime

- do not take more than 6 doses in any 24 hours period
- measure only with dosing cup provided. Do not use any other dosing device.
- keep dosing cup with product
- mL = milliliter

Age	Dose
adults and children 12 years and over	20 mL every 4 hours
children 6 to under 11 years	10 mL every 4 hours
Children under 6 years	do not use

Other information

Nighttime

- **each 10 mL contains:** sodium 6 mg

- store at 20-25°C (68-77°F)
- do not refrigerate
- protect from light

Daytime

- **each 10 mL contains:** 5 mg
- store between 20-25°C (68-77°F). Do not refrigerate.

Inactive ingredients

Nighttime

acesulfame potassium, anhydrous citric acid, EDTA disodium, FD&C Blue #1, FD&C red #40, flavor, maltitol, propylene glycol, purified water, sodium benzoate, trisodium citrate dihydrate

Daytime

citric acid, FD&C blue #1, FD&C red #40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucralose

Questions or comments?

Call 1-877-753-3935 Monday-Friday 9AM-5PM EST

Principal Display Panel

Compare to the active ingredients in Children's Dimetapp® Cold & Cough & Nighttime Cold & Congestion††

Children's Cold & Congestion Relief

DIPHENHYDRAMINE HCl 12.5 mg /

ANRIHISTAMINE / COUGH SUPPRESSANT

PHENYLEPHRINE HCl 5 mg /

NASAL DECONGESTANT

Nighttime

Alcohol Free

- Relieves stuffy nose, runny nose, sneezing, itchy & watery eyes & coughing

AGES 6 YEARS & OLDER

Grape flavor

DOSING CUP INCLUDED

AGES 6 + YEARS

TAMPER EVIDENT: DO NOT USE IF CARTON IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING.

††This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Children's Dimetapp® Nighttime Cold & Congestion.

DAYTIME

Children's Cold & Cough Relief

BROMPHENIRAMINE MALEATE 2 mg

ANTIHISTAMINE

DEXTROMETHORPHAN HBr 10 mg /

COUGH SUPPRESSANT

PHENYLEPHRINE HCl 5 mg /

NASAL DECONGESTANT

Daytime

Alcohol Free

- Relieves nasal congestion, runny nose, itchy & watery eyes, sneezing & coughing

AGES 6 YEARS & OLDER

Grape flavor

FL OZ (mL)

††This product is not manufactured or distributed by Pfizer Consumer Healthcare, Distributor of Children's Dimetapp® Cold & Cough.

DISTRIBUTED BY: WALGREEN CO.

200 WILMOT RD., DEERFIELD, IL 60015

Product Label

<p>Walgreens Cold & Congestion Relief Drug Facts</p> <p>Drug Facts (continued)</p> <p>Directions</p> <ul style="list-style-type: none"> do not take more than 6 doses in any 24 hour period do not exceed recommended dosage measure only with dosing cup provided. Do not use any other dosing device. keep dosing cup with product mL = milliliter <table border="1"> <tr> <th>Age</th> <th>Dose</th> </tr> <tr> <td>adults and children 12 years and over</td> <td>20 mL every 4 hours</td> </tr> <tr> <td>children 6 to under 12 years</td> <td>10 mL every 4 hours</td> </tr> <tr> <td>children under 6 years</td> <td>do not use</td> </tr> </table> <p>Other information</p> <ul style="list-style-type: none"> each 10 mL contains: sodium 6 mg store at 20-25°C (68-77°F) do not refrigerate protect from light <p>Inactive ingredients aceaflame potassium, anhydrous citric acid, EDTA disodium, FD&C blue #1, FD&C red #40, flavor, methyl, propylene glycol, purified water, sodium benzoate, titanium dioxide, xanthan gum</p> <p>Questions or comments? Call 1-877-753-8636 Monday-Friday 9AM-5PM EST</p> <p><small>***This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Children's Dimetapp® Cold & Cough.</small></p>	Age	Dose	adults and children 12 years and over	20 mL every 4 hours	children 6 to under 12 years	10 mL every 4 hours	children under 6 years	do not use	<p>Walgreens Cold & Cough Relief Drug Facts</p> <p>Drug Facts</p> <p>Active ingredients (in each 10 mL)</p> <table border="1"> <tr> <th>Active ingredients</th> <th>Purposes</th> </tr> <tr> <td>Brompheniramine maleate 2 mg</td> <td>Antihistamine</td> </tr> <tr> <td>Dextromethorphan HBr 10 mg</td> <td>Cough suppressant</td> </tr> <tr> <td>Phenylephrine HCl 5 mg</td> <td>Nasal decongestant</td> </tr> </table> <p>Uses</p> <ul style="list-style-type: none"> temporarily relieves cough due to minor throat and bronchial irritation occurring with a cold, and nasal congestion due to the common cold, hay fever or other upper respiratory allergies temporarily relieves these symptoms due to hay fever (allergic rhinitis) runny nose itchy, watery eyes sneezing itching of the nose or throat temporarily restores free breathing through the nose <p>Warnings</p> <p>Do not use</p> <ul style="list-style-type: none"> to sedate a child or to make a child sleepy if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> heart disease high blood pressure thyroid disease diabetes glaucoma trouble urinating due to an enlarged prostate gland cough that occurs with too much phlegm (mucus) a breathing problem or persistent or chronic cough that lasts such as occurs with smoking, asthma, chronic bronchitis, or emphysema <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> taking any other oral nasal decongestant or stimulant taking sedatives or tranquilizers <p>When using this product</p> <ul style="list-style-type: none"> do not use more than directed marked drowsiness may occur avoid alcoholic beverages alcohol, sedatives, and tranquilizers may increase drowsiness be careful when driving a motor vehicle or operating machinery excitability may occur, especially in children <p>Stop use and ask a doctor if</p> <ul style="list-style-type: none"> neurovision, dizziness or sleeplessness occur symptoms do not improve within 7 days or are accompanied by fever cough lasts more than 7 days, comes back, or is accompanied by fever, rash, or persistent headache. These could be signs of a serious condition. 	Active ingredients	Purposes	Brompheniramine maleate 2 mg	Antihistamine	Dextromethorphan HBr 10 mg	Cough suppressant	Phenylephrine HCl 5 mg	Nasal decongestant	<p>Walgreens Cold & Cough Relief Drug Facts</p> <p>Drug Facts (continued)</p> <p>If pregnant or breast-feeding, ask a health professional before use.</p> <p>Keep out of reach of children.</p> <p>In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.</p> <p>Directions</p> <ul style="list-style-type: none"> do not take more than 6 doses in any 24-hour period measure only with dosing cup provided. Do not use any other dosing device. keep dosing cup with product mL = milliliter <table border="1"> <tr> <th>Age</th> <th>Dose</th> </tr> <tr> <td>adults and children 12 years and over</td> <td>20 mL every 4 hours</td> </tr> <tr> <td>children 6 to 11 years</td> <td>10 mL every 4 hours</td> </tr> <tr> <td>children under 6 years</td> <td>do not use</td> </tr> </table> <p>Other information</p> <ul style="list-style-type: none"> each 10 mL contains: sodium 5 mg store between 20-25°C (68-77°F). Do not refrigerate. <p>Inactive ingredients citric acid, FD&C blue #1, FD&C red #40, flavor, glycerin, propylene glycol, purified water, sodium benzoate, sodium citrate, sorbitol, sucrose</p> <p>Questions or comments? Call 1-877-753-8636 Monday-Friday 9AM-5PM EST</p> <p>CAUTION: DO NOT USE IF CAPTION IS OPENED OR IF PRINTED SAFETY SEAL AROUND BOTTLE OR UNDER CAP IS BROKEN OR MISSING</p> <p><small>*Our pharmacists recommend the Walgreens brand. We invite you to compare to national brands.</small></p> <p><small>***This product is not manufactured or distributed by Pfizer Consumer Healthcare, distributor of Children's Dimetapp® Nighttime Cold & Congestion.</small></p> <p>DISTRIBUTED BY: WALGREEN CO. 200 WILMOT RD., DEERFIELD, IL 60015 Walgreens 100% SATISFACTION GUARANTEED Walgreens.com ©2021 Walgreen Co.</p>	Age	Dose	adults and children 12 years and over	20 mL every 4 hours	children 6 to 11 years	10 mL every 4 hours	children under 6 years	do not use	<p>DAY & NIGHT PACK NDC 0363-4590-08</p> <p>Walgreens <small>WALGREENS PHARMACY RECOMMENDS</small></p> <p>Compare to the active ingredients in Children's Dimetapp® Cold & Cough & Nighttime Cold & Congestion®</p> <p>Children's Cold & Cough Relief</p> <p>BROMPHENIRAMINE MALEATE 2 mg / ANTIHISTAMINE DEXTROMETHORPHAN HBr 10 mg / COUGH SUPPRESSANT PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT</p> <p>Children's Cold & Congestion Relief</p> <p>DIPHENHYDRAMINE HCl 12.5 mg / ANTIHISTAMINE / COUGH SUPPRESSANT PHENYLEPHRINE HCl 5 mg / NASAL DECONGESTANT</p> <p>Daytime</p> <p>Alcohol Free</p> <ul style="list-style-type: none"> Relieves nasal congestion, runny nose, itchy & watery eyes, sneezing & coughing <p>Nighttime</p> <p>Alcohol Free</p> <ul style="list-style-type: none"> Relieves stuffy nose, runny nose, sneezing, itchy & watery eyes & coughing <p>AGES 6 YEARS & OLDER Grape flavor</p> <p>AGES 6 YEARS & OLDER Grape flavor</p> <p>2 - 4 FL OZ (118 mL) BOTTLES / TOTAL 8 FL OZ (236 mL)</p> <p>ITEM 904373 W10024-0322-L</p> <p>3 1191717679 6</p> <p>PAPER BOX PLASTIC BOTTLE PLASTIC CUP</p> <p>PLD-ES910 PLD-ES914 F0008013</p> <p>Lot No: Exp. Date:</p>	<p>Walgreens Cold & Congestion Relief Drug Facts</p> <p>Drug Facts</p> <p>Active ingredients (in each 10 mL)</p> <table border="1"> <tr> <th>Active ingredients</th> <th>Purposes</th> </tr> <tr> <td>Diphenhydramine HCl 12.5 mg</td> <td>Antihistamine/Cough suppressant</td> </tr> <tr> <td>Phenylephrine HCl 5 mg</td> <td>Nasal decongestant</td> </tr> </table> <p>Uses</p> <ul style="list-style-type: none"> temporarily relieves these symptoms occurring with a cold, hay fever, or other upper respiratory allergies nasal congestion cough runny nose sneezing itchy, watery eyes itching of the nose or throat <p>Warnings</p> <p>Do not use</p> <ul style="list-style-type: none"> to sedate a child or to make a child sleepy if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. with any other product containing diphenhydramine, even one used on skin <p>Ask a doctor before use if you have</p> <ul style="list-style-type: none"> heart disease high blood pressure thyroid disease diabetes glaucoma trouble urinating due to an enlarged prostate gland cough that occurs with too much phlegm (mucus) a breathing problem or chronic cough that lasts or as occurs with smoking, asthma, chronic bronchitis, or emphysema <p>Ask a doctor or pharmacist before use if you are</p> <ul style="list-style-type: none"> taking any other oral nasal decongestant or stimulant taking sedatives or tranquilizers <p>When using this product</p> <ul style="list-style-type: none"> do not use more than directed marked drowsiness may occur avoid alcoholic drinks alcohol, sedatives, and tranquilizers may increase drowsiness be careful when driving a motor vehicle or operating machinery excitability may occur, especially in children you get nervous, dizzy, or sleepless symptoms do not get better within 7 days or occur with a fever cough lasts more than 7 days, comes back, or is accompanied by a fever, rash, or persistent headache. These could be signs of a serious condition. <p>If pregnant or breast-feeding, ask a health professional before use. Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away.</p>	Active ingredients	Purposes	Diphenhydramine HCl 12.5 mg	Antihistamine/Cough suppressant	Phenylephrine HCl 5 mg	Nasal decongestant
Age	Dose																																	
adults and children 12 years and over	20 mL every 4 hours																																	
children 6 to under 12 years	10 mL every 4 hours																																	
children under 6 years	do not use																																	
Active ingredients	Purposes																																	
Brompheniramine maleate 2 mg	Antihistamine																																	
Dextromethorphan HBr 10 mg	Cough suppressant																																	
Phenylephrine HCl 5 mg	Nasal decongestant																																	
Age	Dose																																	
adults and children 12 years and over	20 mL every 4 hours																																	
children 6 to 11 years	10 mL every 4 hours																																	
children under 6 years	do not use																																	
Active ingredients	Purposes																																	
Diphenhydramine HCl 12.5 mg	Antihistamine/Cough suppressant																																	
Phenylephrine HCl 5 mg	Nasal decongestant																																	

WALGREENS Children's Cold & Congestion Relief, Children's Cold & Cough Relief

DAYTIME COLD AND COUGH AND NIGHTTIME COLD AND CONGESTION CHILDRENS

brompheniramine maleate, dextromethorphan hbr, phenylephrine hcl, diphenhydramine hcl, phenylephrine hcl kit

Product Information

Product Type HUMAN OTC DRUG **Item Code (Source)** NDC:0363-4590

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:0363-4590-08	1 in 1 KIT; Type 0: Not a Combination Product	03/04/2022	

Quantity of Parts

Part #	Package Quantity	Total Product Quantity
Part 1	1 BOTTLE, PLASTIC	118 mL
Part 2	1 BOTTLE, PLASTIC	118 mL

NIGHT TIME COLD AND CONGESTION CHILDRENS

diphenhydramine hcl, phenylephrine hcl liquid

Product Information

Item Code (Source) NDC:0363-3130

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
DIPHENHYDRAMINE HYDROCHLORIDE (UNII: TC2D6JAD40) (DIPHENHYDRAMINE - UNII:8GTS82S83M)	DIPHENHYDRAMINE HYDROCHLORIDE	12.5 mg in 10 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
ACESULFAME POTASSIUM (UNII: 23OV73Q5G9)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
EDETATE DISODIUM (UNII: 7FLD91C86K)	
MALTITOL (UNII: D65DG142WK)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1		118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
--------------------	--	----------------------	--------------------

Part 2 of 2**DAYTIME COLD AND COUGH CHILDRENS**

brompheniramine maleate, dextromethorphan hbr, phenylephrine hcl liquid

Product Information

Item Code (Source) NDC:0363-4580

Route of Administration ORAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
BROMPHENIRAMINE MALEATE (UNII: IXA7C9ZN03) (BROMPHENIRAMINE - UNII:H57G17P2FN)	BROMPHENIRAMINE MALEATE	2 mg in 10 mL
DEXTROMETHORPHAN HYDROBROMIDE (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	10 mg in 10 mL
PHENYLEPHRINE HYDROCHLORIDE (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	5 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
GLYCERIN (UNII: PDC6A3C00X)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
WATER (UNII: 059QF0KO0R)	
SORBITOL (UNII: 506T60A25R)	
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
FD&C BLUE NO. 1 (UNII: H3R47K3TBD)	
FD&C RED NO. 40 (UNII: WZB9127XOA)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCRALOSE (UNII: 96K6UQ3ZD4)	
TRISODIUM CITRATE DIHYDRATE (UNII: B22547B95K)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	GRAPE	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
---	-----------	---------------------	----------------------	--------------------

1	118 mL in 1 BOTTLE, PLASTIC; Type 0: Not a Combination Product		
---	--	--	--

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/04/2022	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	03/04/2022	

Labeler - Walgreens (008965063)

Revised: 6/2024

Walgreens