

I-MAX LIGHTENING L- hydroquinone cream
MAXLIFE USA, INC.

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

ACTIVE INGREDIENTS:
HYDROQUINONE USP 2%

PURPOSE:

SKIN LIGHTENING

USES:

FOR THE GRADUAL FADING OF DARK AREAS OF THE SKIN.

WARNINGS:

AVOID CONTACT WITH EYES. SOME USERS MAY EXPERIENCE MILD SKIN IRRITATION.
IF IRRITATION BECOMES SEVERE, STOP USE AND CONSULT A DOCTOR.

THIS PRODUCT IS NOT INTENDED FOR USE IN THE PREVENTION OF SUNBURN AND CONTAINS AN ALPHA HYDROXY ACID (AHA) THAT MAY INCREASE YOUR SKIN'S SENSITIVITY TO THE SUN AND PARTICULARLY THE POSSIBILITY OF SUNBURN. SUN EXPOSURE SHOULD BE LIMITED BY USING A SUNSCREEN AGENT OR PROTECTIVE CLOTHING TO COVER BLEACHED SKIN AFTER TREATMENT IS COMPLETED TO PREVENT DARKENING FROM REOCCURRING.

DIRECTIONS:

ADULTS: APPLY A SMALL AMOUNT AS A THIN LAYER ON THE AFFECTED AREA TWICE DAILY, OR USE AS DIRECTED BY A DOCTOR. IF NO IMPROVEMENT IS SEEN AFTER 3 MONTHS OF TREATMENT, USE OF THIS PRODUCT SHOULD BE DISCONTINUED. LIGHTENING EFFECT OF THIS PRODUCT MAY NOT BE NOTICEABLE WHEN USED ON VERY DARK SKIN.

CHILDREN UNDER 12 YEARS OF AGE: DO NOT USE UNLESS DIRECTED BY A DOCTOR.

INACTIVE INGREDIENTS:


ALPHA ARBUTIN, C13-14 ISOPARAFFIN, CETYL ALCOHOL, ETHYLHEXYLGLYCERIN, ETHYLHEXYL STEARATE, GLYCERIN, GLYCERYL STEARATE, GLYCOLIC ACID, ISOPROPYL MYRISTATE, LAURETH-7, NIACINAMIDE, PANTOTHENIC ACID, PEG-100 STEARATE, PHENOXYETHANOL, PHENYLETHYL RESORCINOL, POLYACRYLAMIDE, PYRIDOXINE HCL, SODIUM HYDROXIDE, SODIUM METABISULFITE, TETRAHEXYLDECYL ASCORBATE, WATER (AQUA), XANTHAN GUM.

QUESTIONS? 1-323-733-7033

KEEP OUT OF REACH OF CHILDREN.

NDC 42952-203-12

Lightening Cream



AMI & PMI



for Intimate Area

2 oz/59 g

I-MaxTM EXCELLENCETM

Drug Facts

Active ingredients:

Hydroquinone USP 2%

Uses:

For the gradual fading of dark areas c

Warnings:

Avoid contact with sensitive areas. Se mild skin irritation. If irritation becom consult a doctor. Do not use on childr unless directed by a doctor.

Directions:

Adults: apply a small amount as a thi twice daily, or use as directed by a do seen after 3 months of treatment, us discontinued. Lightening effect of thi noticeable when used on very dark sl Children under 12 years old: do not u

Inactive ingredients:

Alpha Arbutin, Butylated Hydroxytolu Cetyl Alcohol, Disodium EDTA, Ethylr Stearate, Glycerin, Glyceryl Stearate, Myristate, Laureth-7, Niacinamide, P Stearate, Phenoxethanol, Phenyleth Pyridoxine HCl, Sodium Hydroxide, Sc Tetrahexylecyl Ascorbate, Tocopher Xanthan Gum.

Avoid storage at extreme temperature

Questions? 1-823-733-7033

MaxLife USA, Inc Los Ang

I-MAX LIGHTENING L

hydroquinone cream

Product Information

| | | | |
|-------------------------|----------------|--------------------|---------------|
| Product Type | HUMAN OTC DRUG | Item Code (Source) | NDC:42952-203 |
| Route of Administration | TOPICAL | | |

Active Ingredient/Active Moiety

| Ingredient Name | Basis of Strength | Strength |
|--|-------------------|--------------|
| HYDROQUINONE (UNII: XV74C1N1AE) (HYDROQUINONE - UNII:XV74C1N1AE) | HYDROQUINONE | 2 g in 100 g |

Inactive Ingredients

| Ingredient Name | Strength |
|--|----------|
| C13-14 ISOPARAFFIN (UNII: E4F12ROE70) | |
| CETYL ALCOHOL (UNII: 936JST6JCN) | |
| ETHYLHEXYLGLYCERIN (UNII: 147D247K3P) | |
| ETHYLHEXYL STEARATE (UNII: EG3PA2K3K5) | |
| GLYCERIN (UNII: PDC6A3C0OX) | |
| GLYCERYL MONOSTEARATE (UNII: 230OU9XXE4) | |
| GLYCOLIC ACID (UNII: 0WT12SX38S) | |
| ISOPROPYL MYRISTATE (UNII: 0RE8K4LNJS) | |

| |
|--|
| LAURETH-7 (UNII: Z95S6G8201) |
| NIACINAMIDE (UNII: 25X51I8RD4) |
| PANTOTHENIC ACID (UNII: 19F5HK2737) |
| PEG-100 STEARATE (UNII: YD01N1999R) |
| PHENOXYETHANOL (UNII: HIE492ZZ3T) |
| POLYACRYLAMIDE (10000 MW) (UNII: E2KR9C9V2I) |
| PYRIDOXINE HYDROCHLORIDE (UNII: 68Y4CF58BV) |
| SODIUM HYDROXIDE (UNII: 55X04QC32I) |
| SODIUM METABISULFITE (UNII: 4VON5FNS3C) |
| TETRAHEXYLDECYL ASCORBATE (UNII: 9LBV3F07AZ) |
| WATER (UNII: 059QF0K00R) |
| XANTHAN GUM (UNII: TTV12P4NEE) |
| ALPHA-ARBUTIN (UNII: 72VUP07IT5) |
| PHENYLETHYL RESORCINOL (UNII: G37UFG162O) |

Packaging

| # | Item Code | Package Description | Marketing Start Date | Marketing End Date |
|---|------------------|---|----------------------|--------------------|
| 1 | NDC:42952-203-12 | 59 g in 1 TUBE; Type 0: Not a Combination Product | 09/18/2012 | |

Marketing Information

| Marketing Category | Application Number or Monograph Citation | Marketing Start Date | Marketing End Date |
|-------------------------|--|----------------------|--------------------|
| OTC monograph not final | part358A | 09/18/2012 | |

Labeler - MAXLIFE USA, INC. (785111431)