

IBUPROFEN INFANTS DROPS - ibuprofen suspension/ drops
Harris Teeter, LLC

Ibuprofen Infants Drops

ACTIVE INGREDIENT (in each 1.25 mL)

Ibuprofen, USP 50 mg (NSAID)*
*nonsteroidal anti-inflammatory drug

PURPOSE

Pain reliever/fever reducer

USE(S)

temporarily:

- reduces fever
- relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches

WARNINGS

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:

- hives
- facial swelling
- asthma (wheezing)
- shock
- skin reddening
- rash
- blisters

If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by doctor.

DO NOT USE

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

ASK A DOCTOR BEFORE USE IF

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease or had a stroke
- child has asthma
- child is taking a diuretic

ASK A DOCTOR OR PHARMACIST BEFORE USE IF THE CHILD IS

- under a doctor's care for any serious condition
- taking any other drug

WHEN USING THIS PRODUCT

- take with food or milk if stomach upset occurs

STOP USE AND ASK DOCTOR IF

- child experiences any of the following signs of stomach bleeding:
- feels faint
- vomits blood
- has bloody or black stools
- has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
- chest pain
- trouble breathing

- weakness in one part or side of the body
- slurred speech
- leg swelling
- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

KEEP OUT OF REACH OF CHILDREN

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

DIRECTIONS

- **this product does not contain directions or complete warnings for adult use**
- **do not give more than directed**
- shake well before using
- find right dose on chart below. If possible, use weight to dose; otherwise use age.
- mL=milliliter
- measure with dosing device provided. Do not use any other dosing device.
- dispense liquid slowly into the child's mouth, toward the inner cheek
- if needed, repeat dose every **6-8 hours**
- do not use more than **4 times a day**

Dosing Chart

WEIGHT (lb)	AGE (mos)	DOSE (mL)
under 6 mos		ask a doctor
12-17 lbs	6-11 mos	1.25 mL
18-23 lbs	12-23 years	1.875 mL

OTHER INFORMATION

- store between 20-25°C (68-77°F)
- **do not use if carton is opened or if printed neckband is imprinted with "SEALED FOR YOUR PROTECTION" and "USE WITH ENCLOSED DOSING DEVICE ONLY" is broken or missing.**

INACTIVE INGREDIENTS

anhydrous citric acid, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sorbitol, sucrose, xanthan gum

QUESTIONS OR COMMENTS?

call 1-609-860-2600 (Monday - Friday 8:00 am - 4:30 pm EST)

PRINCIPAL DISPLAY PANEL

Compare to the active ingredient in Motrin[®] Infants' Drops Dye-Free*

Harris Teeter™

NDC 69256-687-09

For Ages 6 mos. to 23 mos.

Infants' Concentrated Drops

Ibuprofen

Oral Suspension

50 mg per 1.25 mL

PAIN RELIEVER/FEVERREDUCER (NSAID)

Dye-Free

- Lasts up to 8 Hours
- Use Only With Enclosed Syringe
- Non-Staining

Berry Flavor

1 FL OZ (30 mL)

Drug Facts (continued)

Warnings

Allergy alert: Ibuprofen may cause a severe allergic reaction, especially in people allergic to aspirin. Symptoms may include:
 ■ hives ■ facial swelling ■ asthma (wheezing)
 ■ shock ■ skin reddening ■ rash ■ blisters
 If an allergic reaction occurs, stop use and seek medical help right away.

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if your child:

- has had stomach ulcers or bleeding problems
- takes a blood thinning (anticoagulant) or steroid drug
- takes other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- takes more or for a longer time than directed

Heart attack and stroke warning: NSAIDs, except aspirin, increase the risk of heart attack, heart failure, and stroke. These can be fatal. The risk is higher if you use more than directed or for longer than directed.

Sore throat warning: Severe or persistent sore throat or sore throat accompanied by high fever, headache, nausea, and vomiting may be serious. Consult doctor promptly. Do not use more than 2 days or administer to children under 3 years of age unless directed by a doctor.

Do not use

- if the child has ever had an allergic reaction to ibuprofen or any other pain reliever/fever reducer
- right before or after heart surgery

Ask a doctor before use if

- stomach bleeding warning applies to your child
- child has a history of stomach problems, such as heartburn
- child has problems or serious side effects from taking pain relievers or fever reducers
- child has not been drinking fluids
- child has lost a lot of fluid due to vomiting or diarrhea
- child has high blood pressure, heart disease, liver cirrhosis, kidney disease or had a stroke
- child has asthma
- child is taking a diuretic

Ask a doctor or pharmacist before use if the child is

- under a doctor's care for any serious condition
- taking any other drug

When using this product

- take with food or milk if stomach upset occurs

Stop use and ask a doctor if

- child experiences any of the following signs of stomach bleeding:
 - feels faint ■ vomits blood ■ has bloody or black stools
 - has stomach pain that does not get better
- child has symptoms of heart problems or stroke:
 - chest pain ■ trouble breathing
 - weakness in one part or side of the body
 - slurred speech ■ leg swelling



Drug Facts (continued)

- the child does not get any relief within first day (24 hours) of treatment
- fever or pain gets worse or lasts more than 3 days
- redness or swelling is present in the painful area
- any new symptoms appear

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222)

Directions

- this product does not contain directions or complete warnings for adult use
- do not give more than directed
- shake well before using
- find right dose on chart below. If possible, use weight to dose; otherwise use age.
- mL = milliliter
- measure with the dosing device provided. Do not use any other device.
- dispense liquid slowly into the child's mouth, toward the inner cheek
- if needed, repeat dose every 6-8 hours
- do not use more than 4 times a day

Dosing Chart

WEIGHT (lb)	AGE (mos)	DOSE (mL)
under 6 mos		ask a doctor
12-17 lbs	6-11 mos	1.25 mL
18-23 lbs	12-23 mos	1.875 mL

Other information

- store between 20-25°C (68-77°F)
- do not use if carton is opened or if printed neckband imprinted with "SEALED FOR YOUR PROTECTION" and "USE WITH ENCLOSED DOSING DEVICE ONLY" is broken or missing.

Inactive ingredients anhydrous citric acid, flavors, glycerin, polysorbate 80, pregelatinized starch, purified water, sodium benzoate, sorbitol, sucrose, xanthan gum.

Questions or comments?

Call 1-(809)-860-2600 Monday - Friday, 8:00 am - 4:30 pm EST.

Drug Facts
Active ingredient (in each 1.25 mL)
 Ibuprofen 50 mg (NSAID)
Purpose
 Pain reliever/
 fever reducer
Uses temporarily: ■ reduces fever
 ■ relieves minor aches and pains due to the common cold, flu, sore throat, headaches and toothaches
 ■ nonsteroidal anti-inflammatory drug

Important: Read all product information before using.
 This product is intended for use in children ages 6 months to 23 months of age.

Compare to the active ingredient in Motrin® Infants' Drops Dye-Free*



For Ages 6 Mos. to 23 Mos.
 Infants' Concentrated Drops
Ibuprofen
 Oral Suspension

50 mg per 1.25 mL

PAIN RELIEVER/
 FEVER REDUCER (NSAID)

Dye-Free

- Lasts up to 8 Hours
- Use Only With Enclosed Syringe
- Non-Staining

Berry Flavor



1 FL OZ (30 mL)

MFR# 53041 REV 0123



0 72036 73997 1
 LOT:

* This product is not manufactured or distributed by Johnson & Johnson Consumer Inc., owner of the registered trademark Motrin®.

PROUDLY DISTRIBUTED BY:
 HARRIS TEETER, LLC
 MATTHEWS, NC 28105

1-800-432-6111
 or harristeeter.com



For Ages 6 Mos. to 23 Mos.
 Infants' Concentrated Drops
Ibuprofen
 Oral Suspension

50 mg per 1.25 mL

PAIN RELIEVER/
 FEVER REDUCER (NSAID)

- Lasts up to 8 Hours • Non-Staining
- Use Only With Enclosed Syringe

Berry Flavor

1 FL OZ (30 mL)

Actual Size

← OPEN to read all Warnings and Directions before use

Directions ■ this product does not contain directions or complete warnings for adult use ■ do not give more than directed ■ shake well before using ■ find right dose on chart below. If possible, use weight to dose; otherwise use age. ■ mL = milliliter ■ measure with the dosing device provided. Do not use any other device. ■ dispense liquid slowly into the child's mouth, toward the inner cheek ■ if needed, repeat dose every 6-8 hours ■ do not use more than 4 times a day


 For Ages 6 Mos. to 23 Mos.
 Infants' Concentrated Drops
Ibuprofen
 Oral Suspension
50 mg per 1.25 mL
 PAIN RELIEVER/
 FEVER REDUCER (NSAID)
 • Lasts up to 8 Hours • Non-Staining
 • Use Only With Enclosed Syringe
Berry Flavor

1 FL OZ (30 mL)

Weight (lb)	Age (mos)	Dose (mL)
	under 6 mos	ask a doctor
12-17 lbs	6-11 mos	1.25 mL
18-23 lbs	12-23 mos	1.875 mL

LOT:

EXP:

IBUPROFEN INFANTS DROPS

ibuprofen suspension/ drops

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69256-687
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
IBUPROFEN (UNII: WK2XY110QM) (IBUPROFEN - UNII:WK2XY110QM)	IBUPROFEN	50 mg in 1.25 mL

Inactive Ingredients

Ingredient Name	Strength
ANHYDROUS CITRIC ACID (UNII: XF417D3PSL)	
GLYCERIN (UNII: PDC6A3C0OX)	
POLYSORBATE 80 (UNII: 6OZP39ZG8H)	
STARCH, CORN (UNII: O8232NY3SJ)	
WATER (UNII: 059QF0KO0R)	
SODIUM BENZOATE (UNII: OJ245FE5EU)	
SUCROSE (UNII: C151H8M554)	
XANTHAN GUM (UNII: TTV12P4NEE)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color		Score	
Shape		Size	
Flavor	BERRY (Mixed Berry)	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69256-687-09	1 in 1 CARTON	01/29/2025	
1		30 mL in 1 BOTTLE; Type 0: Not a Combination Product		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANDA	ANDA210755	01/29/2025	

Labeler - Harris Teeter, LLC (048463103)

Revised: 2/2025

Harris Teeter, LLC