

**RITE AID REGULAR STRNGTH PINK BISMUTH- bismuth  
subsalicylate suspension  
RITE AID CORPORATION**

-----  
**RITE AID Regular Strength Pink Bismuth Subsalicylate 525 mg Drug Facts**

**Active ingredient (in each 30 mL dose cup)**

Bismuth subsalicylate 525 mg

**Purposes**

Upset stomach reliever and antidiarrheal

**Uses**

relieves

- travelers' diarrhea
- diarrhea
- upset stomach due to overindulgence in food and drink, including:
- heartburn
- indigestion
- nausea
- gas
- belching

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea or vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert: Contains salicylate.** Do not take if you are

- allergic to salicylates (including aspirin)
- taking other salicylate products

**Do not use if you have**

- an ulcer
- a bleeding problem
- bloody or black stool

**Ask a doctor before use if you have**

- fever

- mucus in the stool

### **Ask a doctor or pharmacist before use if you are**

taking any drug for

- anticoagulation (thinning the blood)
- diabetes
- gout
- arthritis

### **When using this product**

a temporary, but harmless, darkening of the stool and/or tongue may occur

### **Stop use and ask a doctor if**

- symptoms get worse or last more than 2 days
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

### **If pregnant or breast-feeding,**

ask a health professional before use.

### **Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

### **Directions**

- shake well before use
- Only use dose cup provided
- **adults and children 12 years and over:** 1 dose (30 mL) every 1/2 hour or 60 mL (2 doses) every hour as needed for diarrhea/traveler's diarrhea
- **30 mL (1 dose) every 1/2 hour as needed for overindulgence (upset stomach, heartburn, indigestion, nausea)**
- do not exceed 8 doses (240 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- **children under 12 years:** ask a doctor
- drink plenty of clear fluids to help prevent dehydration caused by diarrhea

### **Other information**

- **each 30 mL contains:** potassium 15 mg, sodium 11 mg
- **salicylate 260 mg**
- low sodium
- sugar free
- store at room temperature
- protect from freezing

- avoid excessive heat (over 104°F 40°C)

### **Inactive ingredients**

carboxymethylcellulose sodium, D&C Red No. 22, D&C Red No. 28, flavor, microcrystalline cellulose, potassium hydroxide, potassium sorbate, purified water, salicylic acid, simethicone emulsion, sodium benzoate, sucralose, xanthan gum

### **Questions or comments?**

**1-866-467-2748**

**Do not use if imprinted shrink band is missing or broken.**

### **Principal Display Panel**

meijer®

NDC# **41250-671-16**

**Compare to the active ingredient in Pepto-Bismol® Regular Strength**

**NDC 11822-7016-6**

**Regular Strength**

**Pink bismuth**

Bismuth Subsalicylate

Upset Stomach Reliever/ Antidiarrheal

Relieves:

- Nausea
- Heartburn
- Indigestion
- Upset Stomach
- Diarrhea

### **Cherry Flavor**

Naturally and Artificially Flavored

16 FL. OZ. (473 mL)

**\*This product is not manufactured or distributed by The Procter & Gamble Co., distributor of Pepto-Bismol® Regular Strength.**



## RITE AID REGULAR STRNGTH PINK BISMUTH

bismuth subsalicylate suspension

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:11822-7016
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (BISMUTH CATION - UNII:ZS9CD1I8YE, SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	525 mg in 30 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>CARBOXYMETHYLCELLULOSE SODIUM, UNSPECIFIED</b> (UNII: K679OBS311)	
<b>D&amp;C RED NO. 22</b> (UNII: 1678RKX8RT)	
<b>D&amp;C RED NO. 28</b> (UNII: 767IP0Y5NH)	
<b>MICROCRYSTALLINE CELLULOSE</b> (UNII: OP1R32D61U)	
<b>POTASSIUM HYDROXIDE</b> (UNII: WZ H3C48M4T)	
<b>POTASSIUM SORBATE</b> (UNII: 1VPU26JZ Z4)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SALICYLIC ACID</b> (UNII: O414PZ4LPZ)	
<b>DIMETHICONE, UNSPECIFIED</b> (UNII: 92RU3N3Y1O)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	

<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)				
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)				
<b>Product Characteristics</b>				
<b>Color</b>	RED	<b>Score</b>		
<b>Shape</b>		<b>Size</b>		
<b>Flavor</b>	CHERRY	<b>Imprint Code</b>		
<b>Contains</b>				
<b>Packaging</b>				
<b>#</b>	<b>Item Code</b>	<b>Package Description</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
1	NDC:11822-7016-6	473 mL in 1 BOTTLE; Type 0: Not a Combination Product	01/02/2023	
<b>Marketing Information</b>				
<b>Marketing Category</b>		<b>Application Number or Monograph Citation</b>	<b>Marketing Start Date</b>	<b>Marketing End Date</b>
OTC Monograph Drug		M008	01/02/2023	

**Labeler -**
RITE AID CORPORATION (014578892)