

MAXIMUM STRENGTH ANTACID - aluminum hydroxide, magnesium hydroxide, simethicone liquid
Harris Teeter, LLC

Maximum Strength Antacid Mint

ACTIVE INGREDIENTS (in each 10 mL)

Aluminum hydroxide (equiv. to dried gel, USP) 800 mg
Magnesium hydroxide 800 mg
Simethicone 80 mg

PURPOSE

Antacid
Antacid
Antigas

USE(S)

relieves:

- acid indigestion
- heartburn
- sour stomach
- upset stomach and gas associated with these symptoms

WARNINGS

Do not take more than 60 mL in a 24-hour period, or use the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a doctor.

ASK A DOCTOR BEFORE USE IF YOU HAVE

- kidney disease
- a magnesium-restricted diet

ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE

- presently taking a prescription drug. Antacids may interact with certain prescription drugs.

STOP USE AND ASK DOCTOR

if symptoms last more than two weeks

KEEP OUT OF REACH OF CHILDREN

DIRECTIONS

- **shake well before use**
- adults and children 12 years and older: take 10 mL to 20 mL two times a day, or as directed by a doctor
- children under 12 years: consult a doctor
- mL = milliliter

OTHER INFORMATION

- **each 10 mL contains:** magnesium 340 mg, sodium 5 mg
- store at controlled room temperature 20°C-25°C (68°-77°F)
- do not freeze

INACTIVE INGREDIENTS

ethyl alcohol, flavor, glycerin, hydroxyethyl cellulose, methylparaben, propylene glycol, propylparaben, purified water, saccharin sodium, simethicone emulsion, sorbitol

PRINCIPAL DISPLAY PANEL

Harris Teeter™

NDC 69256-517-05

MAXIMUM STRENGTH

ANTACID

Antacid & Antigas

Heartburn & Acid Indigestion


Pressure & Bloating

Compare to the active ingredients in Mylanta® Maximum Strength*

Mint Flavored Liquid

12 FL OZ (355 mL)

Alcohol Content 0.2% v/v



Drug Facts **DO NOT USE IF PRINTED SEAL UNDER CAP IS BROKEN OR MISSING**

Active ingredients (in each 10 mL)	Purpose
Aluminum hydroxide (equiv. to dried gel, USP) 800mg	Antacid
Magnesium hydroxide 800 mg	Antacid
Simethicone 80 mg	Antigas

Uses relieves: ■ acid indigestion ■ heartburn ■ sour stomach ■ upset stomach and gas associated with these symptoms

Warnings
Do not take more than 60 mL in a 24-hour period, or use the maximum dosage of this product for more than 2 weeks, except under the advice and supervision of a doctor.

Ask a doctor before use if you have ■ kidney disease ■ a magnesium-restricted diet

Ask a doctor or pharmacist before use if you are ■ presently taking a prescription drug. Antacids may interact with certain prescription drugs.

Stop use and ask a doctor if symptoms last more than two weeks

Keep out of reach of children.

Directions ■ shake well before use ■ adults and children 12 years and older: take 10 - 20 mL two times a day, or as directed by a doctor ■ children under 12 years: consult a doctor ■ mL = milliliter

Other information ■ each 10 mL contains: magnesium 340 mg, sodium 5 mg ■ store at controlled room temperature 20°C-25°C (68°F-77°F) ■ do not freeze

Inactive ingredients ethyl alcohol, flavor, glycerin, hydroxyethyl cellulose, methylparaben, propylene glycol, propylparaben, purified water, saccharin sodium, simethicone emulsion, sorbitol.

*This product is not manufactured or distributed by Infirst Healthcare Inc., the distributor of MYLANTA® Maximum Strength.

PROUDLY DISTRIBUTED BY:
HARRIS TEETER, LLC
MATTHEWS, NC 28105

1-800-432-6111
or harristeeter.com

MFR# 53041 REV 0121



MAXIMUM STRENGTH ANTACID

aluminum hydroxide, magnesium hydroxide, simethicone liquid

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:69256-517
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ALUMINUM HYDROXIDE (UNII: 5QB0T2IUN0) (ALUMINUM HYDROXIDE - UNII:5QB0T2IUN0)	ALUMINUM HYDROXIDE	800 mg in 10 mL
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838)	MAGNESIUM HYDROXIDE	800 mg in 10 mL
DIMETHICONE (UNII: 92RU3N3Y10) (DIMETHICONE - UNII:92RU3N3Y10)	DIMETHICONE	80 mg in 10 mL

Inactive Ingredients

Ingredient Name	Strength
ALCOHOL (UNII: 3K9958V90M)	
GLYCERIN (UNII: PDC6A3C0OX)	
HYDROXYETHYL CELLULOSE, UNSPECIFIED (UNII: T4V6TWG28D)	
METHYLPARABEN (UNII: A2I8C7HI9T)	
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)	
PROPYLPARABEN (UNII: Z8IX2SC1OH)	
WATER (UNII: 059QF0KO0R)	
SACCHARIN SODIUM (UNII: SB8ZUX40TY)	
SORBITOL (UNII: 506T60A25R)	

Product Characteristics

Color	WHITE	Score	
Shape		Size	
Flavor	MINT	Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69256-517-05	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/06/2025	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M001	02/06/2025	

Labeler - Harris Teeter, LLC (048463103)

Establishment

Name	Address	ID/FEI	Business Operations
------	---------	--------	---------------------

Guardian Drug Company

119210276

MANUFACTURE(69256-517)

Revised: 2/2025

Harris Teeter, LLC