

**STOMACH RELIEF - bismuth subsalicylate liquid**  
**Harris Teeter, LLC**

-----  
**Regular Strength Stomach Relief**

**ACTIVE INGREDIENT(in each 30 mL)**

Bismuth subsalicylate 525 mg

**PURPOSE**

Upset stomach reliever and anti-diarrheal

**USE(S)**

relieves:

- diarrhea
- heartburn
- indigestion
- nausea
- upset stomach associated with these symptoms

**WARNINGS**

**Reye's Syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's Syndrome, a rare but serious illness.

**Allergy alert: Contains salicylate. Do not take if you are**

- allergic to salicylates (including aspirin)
- taking other salicylate products

**DO NOT USE IF YOU HAVE**

- an ulcer
- bloody or black stool
- a bleeding problem

**ASK A DOCTOR BEFORE USE IF YOU HAVE**

- fever
- mucus in the stool

## **ASK A DOCTOR OR PHARMACIST BEFORE USE IF YOU ARE**

taking any drug for

- anticoagulation (thinning of the blood)
- diabetes
- gout
- arthritis

## **WHEN USING THIS PRODUCT**

a temporary, but harmless darkening of the stool and/or tongue may occur

## **STOP USE AND ASK DOCTOR IF**

- symptoms get worse
- ringing in the ears or loss of hearing occurs
- diarrhea lasts more than 2 days

## **IF PREGNANT OR BREAST-FEEDING,**

ask a health professional before use.

## **KEEP OUT OF REACH OF CHILDREN**

In case of overdose, get medical help or contact a Poison Control Center immediately.

## **DIRECTIONS**

- **shake well before use**
- mL = milliliter
- TBSP = tablespoon
- adults and children 12 years and over: 1 dose (2 TBSP or 30 mL) every 1/2 to 1 hour as needed
- do not exceed 8 doses (16 TBSP or 240 mL) in 24 hours
- use until diarrhea stops but not more than 2 days
- children under 12 years: ask a doctor
- drink plenty of fluids to help prevent dehydration caused by diarrhea

## **OTHER INFORMATION**

- **each 30 mL or 2 TBSP contains:**
- potassium 25 mg
- salicylate 260 mg
- sodium 8 mg

- protect from freezing
- avoid excessive heat (over 104°F or 40°C)
- dosage cup provided

## **INACTIVE INGREDIENTS**

benzoic acid, D&C red # 22, D&C red # 28, flavor, hydroxyethyl cellulose, potassium hydroxide, purified water, saccharin sodium, salicylic acid, simethicone, xanthan gum

## **PRINCIPAL DISPLAY PANEL**

Harris Teeter™

NDC 69256-527-04

## **STOMACH RELIEF BISMUTH SUBSALICYLATE LIQUID**

### **Upset Stomach Reliever/Antidiarrheal**

Each 30mL contains:

525 mg bismuth subsalicylate

### **5-Symptom Digestive Relief:**

- Heartburn
- Indigestion
- Nausea
- Diarrhea
- Upset stomach

Compare to active ingredient of Pepto-Bismol\*

Original Flavor

8 FL OZ (236 mL)

**Drug Facts** **DO NOT USE IF IMPRINTED SHRINKBAND IS MISSING OR BROKEN.**

**Active ingredient (in each 30 mL) Purpose**  
Bismuth subsalicylate 525 mg .....Upset stomach reliever and anti-diarrheal

**Uses** relieves: ■ diarrhea ■ heartburn ■ indigestion ■ nausea ■ upset stomach associated with these symptoms

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert:** Contains salicylate. Do not take if you are ■ allergic to salicylates (including aspirin) ■ taking other salicylate products  
**Do not use if you have** ■ an ulcer ■ bloody or black stool ■ a bleeding problem  
**Ask a doctor before use if you have** ■ fever ■ mucus in the stool  
**Ask a doctor or pharmacist before use if you are** taking any drug for ■ anticoagulation (thinning of the blood) ■ diabetes ■ gout ■ arthritis  
**When using this product** a temporary, but harmless darkening of the stool and/or tongue may occur  
**Stop use and ask a doctor if** ■ symptoms get worse ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days  
**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center immediately. **Drug Facts continued**



**STOMACH RELIEF**  
**BISMUTH SUBSALICYLATE LIQUID**

**Upset Stomach Reliever/Antidiarrheal**

Each 30mL contains:  
525 mg bismuth subsalicylate

**5-Symptom Digestive Relief:**

- Heartburn • Indigestion
- Nausea • Upset stomach
- Diarrhea

Compare to the active ingredient in Pepto-Bismol\*

Original Flavor

**8 FL OZ**  
(236 mL)

**Drug Facts (continued)**

**Directions** ■ shake well before use  
■ mL = milliliter ■ TBSP = tablespoon ■ adults and children 12 years and over: 1 dose (2 TBSP or 30 mL) every 1/2 to 1 hour as needed ■ do not exceed 8 doses (16 TBSP or 240 mL) in 24 hours ■ use until diarrhea stops but not more than 2 days ■ children under 12 years: ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Other information** ■ each 30 mL or 2 TBSP contains:  
• potassium 25 mg • salicylate 260 mg • sodium 8 mg ■ protect from freezing. ■ avoid excessive heat (over 104°F or 40°C). ■ dosage cup provided

**Inactive ingredients** benzoic acid, D&C red # 22, D&C red # 28, flavor, hydroxyethyl cellulose, potassium hydroxide, purified water, saccharin sodium, salicylic acid, simethicone, xanthan gum.

\*This product is not manufactured or distributed by Procter & Gamble, Inc., the distributor of Pepto-Bismol®.

**PROUDLY DISTRIBUTED BY:**  
HARRIS TEETER, LLC  
MATTHEWS, NC 28105

1-800-432-6111  
or harristeeter.com

MFR# 53041 REV 0121



0 72036 73537 9

Harris Teeter™

NDC 69256-527-05

**STOMACH RELIEF**  
**BISMUTH SUBSALICYLATE LIQUID**

**Upset Stomach Reliever/Antidiarrheal**

Each 30mL contains:

525 mg bismuth subsalicylate

**5-Symptom Digestive Relief:**

- Heartburn
- Indigestion
- Nausea
- Diarrhea
- Upset stomach

Compare to active ingredient of Pepto-Bismol\*

Original Flavor

12 FL OZ (354 mL)

**Drug Facts** **DO NOT USE IF IMPRINTED SHRINKBAND IS MISSING OR BROKEN.**

**Active ingredient (in each 30 mL)** Purpose  
Bismuth subsalicylate 525 mg .....Upset stomach reliever and anti-diarrheal

**Uses** relieves: ■ diarrhea ■ heartburn ■ indigestion ■ nausea ■ upset stomach associated with these symptoms

**Warnings**  
**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.  
**Allergy alert: Contains salicylate. Do not take if you are**  
■ allergic to salicylates (including aspirin)  
■ taking other salicylate products  
**Do not use if you have** ■ bloody or black stool ■ an ulcer ■ a bleeding problem  
**Ask a doctor before use if you have** ■ fever ■ mucus in the stool  
**Ask a doctor or pharmacist before use if you are taking any drug for**  
■ anticoagulation (thinning of the blood)  
■ diabetes ■ gout ■ arthritis  
**When using this product** a temporary, but harmless darkening of the stool and/or tongue may occur.  
**Stop use and ask a doctor if** ■ symptoms get worse ■ ringing in the ears or loss of hearing occurs ■ diarrhea lasts more than 2 days  
**If pregnant or breast-feeding,** ask a health professional before use. **Keep out of reach of children.** In case of overdose, get medical help or contact a Poison Control Center immediately.

◀ **Drug Facts continued**



# STOMACH RELIEF

## BISMUTH SUBSALICYLATE LIQUID

**Upset Stomach Reliever/Antidiarrheal**

Each 30mL contains:  
525 mg bismuth subsalicylate

**5-Symptom Digestive Relief:**

- Heartburn • Indigestion
- Nausea • Upset stomach
- Diarrhea

Compare to the active ingredient in Pepto-Bismol®

**Original Flavor**

**12 FL OZ**  
(354 mL)

**Drug Facts (continued)**

**Directions** ■ shake well before use ■ mL = milliliter  
■ Tbsp = tablespoon ■ adults and children 12 years and over: 1 dose (2 Tbsp or 30 mL) every 1/2 to 1 hour as needed  
■ do not exceed 8 doses (16 Tbsp or 240 mL) in 24 hours  
■ use until diarrhea stops but not more than 2 days  
■ children under 12 years: ask a doctor ■ drink plenty of clear fluids to help prevent dehydration caused by diarrhea

**Other information** ■ each 30 mL or 2 Tbsp contains:  
• potassium 25 mg • salicylate 260 mg • sodium 8 mg  
■ protect from freezing. ■ avoid excessive heat (over 104°F or 40°C). ■ dosage cup provided

**Inactive ingredients** benzoic acid, D&C red # 22, D&C red # 28, flavor, hydroxyethyl cellulose, potassium hydroxide, purified water, saccharin sodium, salicylic acid, simethicone, xanthan gum.

\*This product is not manufactured or distributed by Procter & Gamble, Inc., the distributor of Pepto-Bismol®.

**PROUDLY DISTRIBUTED BY:**  
**HARRIS TEETER, LLC** 1-800-432-6111  
**MATTHEWS, NC 28105** or harristeeter.com



## STOMACH RELIEF

bismuth subsalicylate liquid

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:69256-527
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>BISMUTH SUBSALICYLATE</b> (UNII: 62TEY51RR1) (SALICYLIC ACID - UNII:O414PZ4LPZ)	BISMUTH SUBSALICYLATE	525 mg in 30 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>BENZOIC ACID</b> (UNII: 8SKN0B0MIM)	
<b>D&amp;C RED NO. 22</b> (UNII: 1678RKX8RT)	
<b>D&amp;C RED NO. 28</b> (UNII: 767IP0Y5NH)	
<b>POTASSIUM HYDROXIDE</b> (UNII: WZ H3C48M4T)	
<b>WATER</b> (UNII: 059QF0KO0R)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SALICYLIC ACID</b> (UNII: O414PZ4LPZ)	
<b>HYDROXYETHYL CELLULOSE (1500 MPA.S AT 1%)</b> (UNII: L605B5892V)	
<b>DIMETHICONE</b> (UNII: 92RU3N3Y1O)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

**Product Characteristics**

<b>Color</b>	PINK	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>		<b>Imprint Code</b>	
<b>Contains</b>			

**Packaging**

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:69256-527-04	236 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/06/2025	
2	NDC:69256-527-05	354 mL in 1 BOTTLE; Type 0: Not a Combination Product	02/06/2025	

**Marketing Information**

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M008	02/06/2025	

**Labeler** - Harris Teeter, LLC (048463103)

Revised: 2/2025

Harris Teeter, LLC