

SALIGARD 60- sodium salicylate solution
Aurora Pharmaceutical, Inc.

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click [here](#).

SaliGard™
Sodium Salicylate
Concentrate 60% w/v

ACTIVE INGREDIENTS

Sodium Salicylate 60% w/v

OTHER INGREDIENTS

Water, sodium phosphate dibasic,
hydrochloric acid

INDICATIONS

For use in the drinking water of poultry and swine as an aid in reducing pain, fever and inflammation.

DIRECTIONS FOR USE

For Analgesic and Antipyretic Use

Water Proportioner Use:

Add 6.5 ounces (192 mL) of Sodium Salicylate 60% Concentrate to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 11.3 mg/lb (25 mg/kg) body weight daily.

Livestock Tank Use:

Add 6.5 ounces (192 mL) of Sodium Salicylate 60% Concentrate to 128 gallons of drinking water. This will achieve a target dose of 11.3 mg/lb (25 mg/kg) body weight daily.

For Anti-Inflammatory/Anti-Prostaglandin Use — Day 1

Water Proportioner Use:

Add 14 ounces (414 mL) of Sodium Salicylate 60% Concentrate to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 22.7 mg/lb (50 mg/kg) body weight daily.

Livestock Tank Use:

Add 14 ounces (414 mL) of Sodium Salicylate 60% Concentrate to 128 gallons of drinking water. This will achieve a target dose of 22.7 mg/lb (50 mg/kg) body weight daily.

Day 2 through 7

Water Proportioner Use:

Add 8 ounces (236 mL) of Sodium Salicylate 60% Concentrate to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 13.6 mg/lb (30 mg/kg) body weight daily.

Livestock Tank Use:

Add 8 ounces (236 mL) of Sodium Salicylate 60% Concentrate to 128 gallons of drinking water. This will achieve a target dose of 13.6 mg/lb (30 mg/kg) body weight daily.

Prepare fresh solutions daily. Repeat as necessary.

WARNING

Do not administer concentrated solution by direct oral administration — gastro-intestinal irritation or overdose may occur.

Do not use in piglets less than 3 weeks of age.

Do not use if allergic or sensitive to the active ingredients.

STORAGE

Store upright up to 25°C (77°F) with brief excursions to 40°C (104°F).

CAUTION

Keep container closed when not in use. Product may solidify at cold storage temperatures. Place container in room temperature storage, which will thaw the solution. If precipitate remains after thawing, place in warm water. Gently invert container to ensure dissolution and uniformity of product. Gradual darkening will not affect product stability.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away.



MANUFACTURED IN THE USA

REORDER NO: 21002

MANUFACTURED BY:

Aurora Pharmaceutical, Inc.

NORTHFIELD, MINNESOTA 55057
1-888-215-1256
www.aurorapharmaceutical.com
IN 50-1714 12/2024

PRINCIPAL DISPLAY PANEL - 3.79 Liter Bottle Label

NDC 51072-117-00

SaliGard™

**Sodium Salicylate
Concentrate 60% w/v**

For Use in Livestock Only

Keep Out of Reach of Children

Net Contents:

1 Gallon (3.79 Liters)

AURORA PHARMACEUTICAL®

ACTIVE INGREDIENT
Sodium Salicylate..... 60% w/v

OTHER INGREDIENTS
Water, sodium phosphate dibasic, hydrochloric acid,

INDICATIONS: For use in the drinking water of poultry and swine as an aid in reducing pain, fever and inflammation.

DIRECTIONS FOR USE:
For Analgesic and Antipyretic Use
Water Proportioner Use: Add 6.5 ounces (192 mL) of Sodium Salicylate 60% Concentrate to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 11.3 mg/lb (25 mg/kg) body weight daily.
Livestock Tank Use: Add 6.5 ounces (192 mL) of Sodium Salicylate 60% Concentrate to 128 gallons of drinking water. This will achieve a target dose of 11.3 mg/lb (25 mg/kg) body weight daily.
(continued on next panel)

MANUFACTURED BY:
Aurora Pharmaceutical, Inc.
NORTHFIELD, MINNESOTA 55057
1-888-215-1256
www.aurorapharmaceutical.com
IN 50-1714 12/2024

REORDER NO: 21602

TAKE TIME

OBSERVE LABEL DIRECTIONS

MANUFACTURED IN THE USA

NDC 51072-117-00

SaliGard 60™

**Sodium Salicylate
Concentrate 60% w/v**

For Use in Livestock Only

Keep Out of Reach of Children

Net Contents:
1 Gallon (3.79 Liters)



DIRECTIONS FOR USE (continued)

For Anti-inflammatory/Anti-Prostaglandin Use — Day 1
Water Proportioner Use: Add 14 ounces (414 mL) of Sodium Salicylate 60% Concentrate to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 22.7 mg/lb (50 mg/kg) body weight daily.
Livestock Tank Use: Add 14 ounces (414 mL) of Sodium Salicylate 60% Concentrate to 128 gallons of drinking water. This will achieve a target dose of 22.7 mg/lb (50 mg/kg) body weight daily.

Day 2 through 7
Water Proportioner Use: Add 8 ounces (236 mL) of Sodium Salicylate 60% Concentrate to make 1 gallon of stock solution and administer through a medicator metered at 1:128 (1 ounce per gallon). This will achieve a target dose of 13.6 mg/lb (30 mg/kg) body weight daily.
Livestock Tank Use: Add 8 ounces (236 mL) of Sodium Salicylate 60% Concentrate to 128 gallons of drinking water. This will achieve a target dose of 13.6 mg/lb (30 mg/kg) body weight daily.
Prepare fresh solutions daily. Repeat as necessary.

WARNING
Not for use in lactating dairy cattle.
Do not administer concentrated solution by direct oral administration — gastro-intestinal irritation or overdose may occur.
Do not use in piglets less than 3 weeks of age.
Do not use if allergic or sensitive to the active ingredients.

STORAGE
Store upright up to 25°C (77°F) with brief excursions to 40°C (104°F).

CAUTION
Keep container closed when not in use. Product may solidify at cold storage temperatures. Place container in room temperature storage, which will thaw the solution. If a precipitate remains after thawing, place in warm water. Gently invert container to ensure dissolution and uniformity of product. Gradual darkening will not affect product stability.

Keep out of reach of children.
In case of overdose, get medical help or contact a Poison Control Center right away.

SALIGARD 60

sodium salicylate solution

Product Information

Product Type	OTC ANIMAL DRUG	Item Code (Source)	NDC:51072-117
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of	Strength
-----------------	----------	----------

Ingredient Name		Strength	Strength	
SODIUM SALICYLATE (UNII: MQ1H85SYP) (SALICYLIC ACID - UNII:O414PZ4LPZ)		SODIUM SALICYLATE	60 g in 100 mL	
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:51072-117-00	3790 mL in 1 BOTTLE		
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
unapproved drug other			05/04/2022	

Labeler - Aurora Pharmaceutical, Inc. (832848639)

Establishment

Name	Address	ID/FEI	Business Operations
Aurora Pharmaceutical, Inc.		832848639	manufacture

Revised: 5/2025

Aurora Pharmaceutical, Inc.