

MILK OF MAGNESIA- magnesium hydroxide suspension

HEB

Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.

HEB Milk of Magnesia Drug Facts

Active ingredient (in each 15 mL)

Magnesium hydroxide 1200 mg

Purpose

Saline laxative

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in ½ to 6 hours

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

Ask a doctor or pharmacist before use if you are

taking any other drug. Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.
- you need to use a laxative for more than 1 week

If pregnant or breast-feeding,

ask a health professional before use.

Keep out of reach of children.

In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not exceed the maximum recommended daily dose in a 24 hour period

- shake well before use
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
- mL = milliliter

adults and children 12 years and older	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	ask a doctor

Other information

- **each 15 mL contains:** calcium 20 mg and magnesium 595 mg
- does not meet USP requirements for preservative effectiveness
- store at 20°-25°C (68°-77°F)
- close cap tightly after use
- avoid freezing

Inactive ingredient

purified water

Questions or comments?

1-800-719-9260

Principal Display Panel

Compare to Phillips'® Milk of Magnesia Original active ingredient

MILK OF MAGNESIA

Magnesium Hydroxide, 1200 mg per 15 mL

Saline Laxative

Comfortable, Cramp Free Relief

Stimulant Free

Sugar Free

Original Flavor

#1 DOCTOR RECOMMENDED saline laxative active ingredient

GLUTEN FREE

12 FL OZ (355 mL)

Compare to Phillips®
Milk of Magnesia Original
active ingredient*

NDC 37808-396-40



MILK OF MAGNESIA

Magnesium Hydroxide, 1200 mg per 15 mL

Saline Laxative

- Comfortable, Cramp Free Relief
- Stimulant Free
- Sugar Free



Original Flavor

GLUTEN FREE 12 FL OZ (355 mL)

: 39640 1J F6

Drug Facts

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING.

Active ingredient (in each 15 mL)

Magnesium hydroxide 1200 mg.....Saline laxative

Purpose

Uses

- relieves occasional constipation (irregularity)
- generally produces bowel movement in ½ to 6 hours

Warnings

Ask a doctor before use if you have

- kidney disease
- a magnesium-restricted diet
- stomach pain, nausea, or vomiting
- a sudden change in bowel habits that lasts more than 2 weeks

Ask a doctor or pharmacist before use if you are taking any other drug.

Take this product two or more hours before or after other drugs. Laxatives may affect how other drugs work.

Stop use and ask a doctor if

- you have rectal bleeding or no bowel movement after using this product. These could be signs of a serious condition.

- you need to use a laxative for more than 1 week

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222).

Directions

- do not exceed the maximum recommended daily dose in a 24 hour period
- shake well before use
- dose may be taken once a day preferably at bedtime, in divided doses, or as directed by a doctor
- drink a full glass (8 oz) of liquid with each dose
- for accurate dosing, use dose cup provided
- mL = milliliter

adults and children 12 years and older	30 mL to 60 mL
children 6 to 11 years	15 mL to 30 mL
children under 6 years	ask a doctor

Other information

- each 15 mL contains: calcium 20 mg and magnesium 595 mg
- does not meet USP requirements for preservative effectiveness
- store at 20°-25°C (68°-77°F)
- close cap tightly after use
- avoid freezing

Inactive ingredient purified water

Questions or comments? 1-800-719-9260

*This product is not manufactured or distributed by Bayer HealthCare LLC, distributor of Phillips® Milk of Magnesia Original.

**In the U.S. Source: AlphalimpactRx

MADE WITH PRIDE & CARE FOR H-E-B®
SAN ANTONIO, TX 78204

1601

100% GUARANTEE PROMISE:
If you aren't completely pleased
with this product, we'll be happy
to replace it or refund your money.
You have our word on it.



: 39640 1J F6

MILK OF MAGNESIA

magnesium hydroxide suspension

Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:37808-396
Route of Administration	ORAL		

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
MAGNESIUM HYDROXIDE (UNII: NBZ3QY004S) (MAGNESIUM CATION - UNII:T6V3LHY838, HYDROXIDE ION - UNII:9159UV381P)	MAGNESIUM HYDROXIDE	1200 mg in 15 mL

Inactive Ingredients

Ingredient Name	Strength
WATER (UNII: 059QF0KO0R)	

Product Characteristics

Color	WHITE (opaque)	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:37808-396-40	355 mL in 1 BOTTLE; Type 0: Not a Combination Product	06/09/1992	

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC monograph not final	part334	06/09/1992	

Labeler - HEB (007924756)