

EUTHANASIA SOLUTION- pentobarbital sodium and phenytoin sodium solution

MWI Veterinary Supply

Euthanasia Solution

Pentobarbital Sodium and Phenytoin Sodium

FOR DOGS ONLY

CAUTION:

Federal law restricts this drug to use by or on the order of a licensed veterinarian.

DESCRIPTION:

A non-sterile solution containing pentobarbital sodium and phenytoin sodium as the active ingredients. Rhodamine B, a bluish-red fluorescent dye, is included in the formulation to help distinguish it from parenteral drugs intended for therapeutic use. Although the solution is not sterile, benzyl alcohol, a bacteriostat, is included to retard the growth of microorganisms.

Each mL contains: *Active ingredients:* 390 mg pentobarbital sodium (barbituric acid derivative), 50 mg phenytoin sodium; *Inactive ingredients:* 10% ethyl alcohol, 18% propylene glycol, 0.003688 mg rhodamine B, 2% benzyl alcohol (preservative), water for injection q.s. Sodium hydroxide and/or hydrochloric acid may be added to adjust pH.

ACTIONS:

Euthanasia Solution contains two active ingredients which are chemically compatible but pharmacologically different. Each ingredient acts in such a manner so as to cause humane, painless, and rapid euthanasia. Euthanasia is due to cerebral death in conjunction with respiratory arrest and circulatory collapse. Cerebral death occurs prior to cessation of cardiac activity.

When administered intravenously, pentobarbital sodium produces rapid anesthetic action. There is a smooth and rapid onset of unconsciousness. At the lethal dose, there is depression of vital medullary respiratory and vasomotor centers.

When administered intravenously, phenytoin sodium produces toxic signs of cardiovascular collapse and/or central nervous system depression. Hypotension occurs when the drug is administered rapidly.

Pharmacodynamic Activity: The sequence of events leading to humane, painless, and rapid euthanasia following intravenous injection of Euthanasia Solution is similar to that following intravenous injection of pentobarbital sodium or other barbituric acid derivatives. Within seconds, unconsciousness is induced with simultaneous collapse of the dog. This stage rapidly progresses to deep anesthesia with concomitant reduction in the blood pressure. A few seconds later, breathing stops, due to depression of the medullary respiratory center; encephalographic activity becomes isoelectric, indicating cerebral death, and then cardiac activity ceases. Phenytoin sodium exerts its effect during the deep anesthesia stage caused by the pentobarbital sodium. This ingredient,

due to its cardiotoxic properties, hastens the stoppage of electrical activity in the heart.

INDICATIONS:

For use in dogs for humane, painless, and rapid euthanasia.

WARNING:

For canine euthanasia only. Must not be used for therapeutic purposes. Do not use in animals intended for food.

<p>ENVIRONMENTAL HAZARD: This product is toxic to wildlife. Birds and mammals feeding on treated animals may be killed. Euthanized animals must be properly disposed of by deep burial, incineration, or other method in compliance with state and local laws, to prevent consumption of carcass material by scavenging wildlife.</p>

HUMAN WARNING:

Caution should be exercised to avoid contact of the drug with open wounds or accidental self-inflicted injections. Keep out of reach of children. If eye contact, flush with water and seek medical advice/attention.

PRECAUTIONS:

Euthanasia may sometimes be delayed in dogs with severe cardiac or circulatory deficiencies. This may be explained by the impaired movement of the drug to its site of action. An occasional dog may elicit reflex responses manifested by motor movement; however, an unconscious animal does not experience pain, because the cerebral cortex is not functioning.

When restraint may cause the dog pain, injury, or anxiety, or danger to the person making the injection, prior use of tranquilizing or immobilizing drugs may be necessary.

Dosage and Administration:

Dosage: Dogs, 1 mL for each 10 pounds of body weight.

Administration: Intravenous injection is preferred. Intracardiac injection may be made when intravenous injection is impractical, as in a very small dog, or in a comatose dog with impaired vascular functions. Good injection skill is necessary for intracardiac injection.

The calculated dose should be given in a single bolus injection.

For intravenous injection, a needle of sufficient gauge to ensure intravenous placement of the entire dose should be used. The use of a Luer-Lok® syringe is recommended to prevent accidental exposure due to needle/syringe separation.

ADVERSE REACTION:

To report suspected adverse events, for technical assistance or to obtain a copy of the Safety Data Sheet (SDS), contact the technical help-line at 866-489-1344. For additional information about adverse drug experience reporting for animal drugs, contact FDA at 1-888-FDA-VETS or online at www.fda.gov/reportanimalae.

HOW SUPPLIED:

Euthanasia Solution (Pentobarbital Sodium and Phenytoin Sodium) is available in 100 ml multi-dose vials in package of one, NDC 86136-138-31.

Manufactured by a nonsterilizing process.

STORAGE:

Store at controlled room temperature 20° to 25°C (68° to 77°F). Use contents within 60 days of first puncture. Approved by FDA under ANADA #200-614

Distributed by: MWI
Boise, ID, 83705
www.VetOne.net

Iss. 10/2025

Luer-Lok is a registered trademark of Becton, Dickinson and Company.

<p>NDC 86136-138-31</p> <p> Euthanasia Solution (Pentobarbital Sodium and Phenytoin Sodium)</p> <p>FOR DOGS ONLY</p> <p>Caution: Federal law restricts this drug to use by or on the order of a licensed veterinarian.</p> <p>Approved by FDA under ANADA # 200-614</p> <p>100 mL V1 503017</p>	<p>Manufactured by a nonsterilizing process.</p> <p>Multi-Dose Vial</p> <p>Each mL contains: <i>active ingredients:</i> 390 mg pentobarbital sodium (barbituric acid derivative), 50 mg phenytoin sodium; <i>inactive ingredients:</i> 10% ethyl alcohol, 18% propylene glycol, 0.003688 mg rhodamine B, 2% benzyl alcohol (preservative), purified water qs. Sodium hydroxide and/or hydrochloric acid may be added to adjust pH.</p> <p>For Intravenous or Intracardiac Use.</p> <div style="border: 1px solid black; padding: 5px;"><p>Environmental Hazard: This product is toxic to wildlife. Birds and mammals feeding on treated animals may be killed. Euthanized animals must be properly disposed of by deep burial, incineration, or other method in compliance with state and local laws, to prevent consumption of carcass material by scavenging wildlife.</p></div> <p>Warning: For canine euthanasia only. Must not be used for therapeutic purpose. Do not use in animals intended for food.</p> <p>Human Warning: Refer to package insert for important human risk information.</p> <p>Store at controlled room temperature 20° to 25°C (68° to 77°F). Use contents within 60 days of first puncture.</p> <p>Read package insert carefully. See Warnings above.</p> <p>Distributed by: MWI, Boise, ID 83705, www.VetOne.net</p>	 <p>3 86136 13831 3</p>
----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------	----------------------------------------------------------------------------------------------------------------------



EUTHANASIA SOLUTION

pentobarbital sodium and phenytoin sodium solution

Product Information

Product Type	PRESCRIPTION ANIMAL DRUG	Item Code (Source)	NDC:86136-138
Route of Administration	INTRAVENOUS	DEA Schedule	CIII

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
-----------------	-------------------	----------

pentobarbital sodium (UNII: NJJ0475N0S) (pentobarbital - UNII:I4744080IR)	pentobarbital sodium	390 mg in 1 mL
phenytoin sodium (UNII: 4182431BJH) (phenytoin - UNII:6158TKW0C5)	phenytoin sodium	50 mg in 1 mL

Product Characteristics

Color	pink	Score	
Shape		Size	
Flavor		Imprint Code	
Contains			

Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:86136-138-31	100 mL in 1 VIAL, MULTI-DOSE		

Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
ANADA	ANADA200614	12/15/2025	

Labeler - MWI Veterinary Supply (019926120)

Revised: 12/2025

MWI Veterinary Supply