STOP HAIR - citric acid monohydrate spray STOP HAIR PTY LTD

Disclaimer: This drug has not been found by FDA to be safe and effective, and this labeling has not been approved by FDA. For further information about unapproved drugs, click here.

Active Ingredient

Citric Acid Monohydrate 0.7 % w/v

Purpose

DEPILATORY/HAIR INHIBITOR

Keep Out of Reach of Children.

Keep Out of Reach of Children.

Indication and Usage

DEPILATORY/HAIR INHIBITOR

Dosage and Administration

Step 1 Remove Hair:

The first is step isof the process is to remove the hair. Any method of hair removal can be used, such as shaving, waxing, etc., but as stated above, methods that remove the hair completely from the follicle are more effective.

Step 2: Apply STOPHAIR - Hair Inhibiting Spray

The second step of the process is to apply the STOPHAIR - Hair Inhibiting Spray. Lightly spray the STOPHAIR solution on the area the hair was removed from and massage in.

This process is to be repeated twice a day for five days after the hair was removed.

Step 3: Continue Applying STOPHAIR - Inhibiting Spray once a day

The third step of the process is to continue applying the STOPHAIR - Hair Inhibiting Spray once a day to the area the hair was removed from. As in Step 2, simply spray the STOPHAIR solution on the area the hair was removed from and lightly massage in.

Continue this process until the next time the hair needs to be removed. This step will vary in legth of time depending on the individual.

Step4: Repeat Steps 2 and 3

The fourth step of the process is to repeat Steps 2 and 3 until the hair permanently stops growing. As above, the number of times this step will have to be repeated depends on the hair properties of the individual.

SPRAY LIGHTLY ONTO SKIN TWICE DAILY FOR 5 DAYS.

Product Label

Manufactured in Australia by StopHair Pty Ltd info@stophair.com.au 1-888-304-3197 (toll free) Contains: Inhibitor 2000 (Proprietary Blend of Plant Extracts) Moisturisers (Glycerol, Glycerine), Almond Extract



DIRECTIONS FOR USE

Apply STOPHAIR - Hair Inhibiting Spray - directly to the skin twice a day (morning and night) for five consecutive days after hair removal and continue once a day thereafter. Repeat this process every time after hair removal.

stophair can be used with any conventional methods of hair removal, including waxing, shaving, depilatory creams or electrolysis.

www.stophair.com.au

125 ml / 4.30 fl. oz.

STOP HAIR

citric acid monohydrate spray

-						-		- 4	r				. •		
u	м	n		П	ct		П	71	hn	и	m		t I	O	n
		u	u	u	L		ш	ш	w		•	a	ш	v	и

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:50965-123

Route of Administration TOPICAL

Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
	Citric Acid Monohydrate	0.7 mg in 100 mL

Inactive Ingredients					
Ingredient Name	Strength				
GLYCERIN (UNII: PDC6 A3C0 OX)					
WATER (UNII: 059QF0KO0R)					
ALMOND (UNII: 3Z252A2K9G)					

Packaging									
#	Item Code	Package Description	Marketing Start Date	Marketing End Date					
1	NDC:50965-123-10	125 mL in 1 BOTTLE, SPRAY							

Marketing Information							
Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date				
unapproved drug other		09/01/2012					

Labeler - STOP HAIR PTY LTD (745058011)

Establishment							
Name	Address	ID/FEI	Business Operations				
STOP HAIR PTY LTD		745058011	manufacture, pack				

Revised: 6/2012 STOP HAIR PTY LTD