

**NIGHTTIME SEVERE COLD AND FLU- acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl solution  
Kroger Company**

-----  
**Kroger Co. NightTime Severe Cold & Flu Drug Facts**

**Active ingredients (in each 30 mL)**

Acetaminophen 650 mg

Dextromethorphan HBr 20 mg

Doxylamine succinate 12.5 mg

Phenylephrine HCl 10 mg

**Purpose**

Pain reliever/fever reducer

Cough suppressant

Antihistamine

Nasal decongestant

**Uses**

temporarily relieves common cold/flu symptoms:

- sinus congestion and pressure
- nasal congestion
- minor aches and pains
- headache
- runny nose and sneezing
- sore throat
- cough to help you sleep
- fever
- cough due to minor throat and bronchial irritation
- reduces swelling of nasal passages
- promotes nasal and/or sinus drainage
- temporarily restores freer breathing through the nose

**Warnings**

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen

- 3 or more alcoholic drinks every day while using this product

**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

### **Do not use**

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients

### **Ask a doctor before use if you have**

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough that occurs with too much phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- trouble urinating due to an enlarged prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema
- a sodium-restricted diet

### **Ask a doctor or pharmacist before use if you are**

- taking sedatives or tranquilizers
- taking the blood thinning drug warfarin

### **When using this product**

- **do not use more than directed**
- excitability may occur, especially in children
- marked drowsiness may occur
- avoid alcoholic drinks

- be careful when driving a motor vehicle or operating machinery
- alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if**

- you get nervous, dizzy or sleepless
- pain, nasal congestion, or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,**

ask a health professional before use.

**Keep out of reach of children.**

**Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions**

- take only as directed – see Overdose warning
- only use the dose cup provided
- do not exceed 4 doses per 24 hrs

adults & children 12 yrs & over	30 mL every 4 hrs
children 4 to under 12 yrs	ask a doctor
children under 4 yrs	do not use

**Other information**

- **each 30 mL contains:** sodium 41 mg
- store at 20-25°C (68-77°F)

**Inactive ingredients**

anhydrous citric acid, edetate disodium, FD&C blue #1, FD&C red #40, flavor, glycerin, propylene glycol, purified water, saccharin sodium, sodium benzoate, sodium chloride, sodium citrate, sorbitol solution, sucralose, xanthan gum

**Questions?**

**1-800-632-6900**

**Package/Label Principal Display Panel**

COMPARE TO the active ingredients of VICKS<sup>®</sup> NYQUIL<sup>®</sup> SEVERE COLD & FLU

See back panel

Kroger<sup>®</sup>

Maximum Strength

NightTime Severe COLD & FLU

Acetaminophen

Dextromethorphan HBr

Doxylamine Succinate

Phenylephrine HCl

OUR PHARMACIST RECOMMENDED

Pain Reliever

Fever Reducer

Cough Suppressant

Antihistamine

Nasal Decongestant

Berry Flavor

12 FL OZ (354 mL)

COMPARE TO the active ingredients of VICKS® NYQUIL® SEVERE COLD & FLU  
\*See back panel

NDC 30142-763-40



Maximum Strength

# NightTime Severe COLD & FLU

Acetaminophen  
Dextromethorphan HBr  
Doxylamine Succinate  
Phenylephrine HCl

Pain Reliever  
Fever Reducer  
Cough Suppressant  
Antihistamine  
Nasal Decongestant



Berry Flavor

12 FL OZ (354 mL)

: 76340 45 FL

DISTRIBUTED BY THE KROGER CO.  
CINCINNATI, OHIO 45202  
QUALITY GUARANTEE  
www.kroger.com

GLUTEN FREE

## PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org

DO NOT USE IF PRINTED NECKBAND IS BROKEN OR MISSING

### Drug Facts

Active ingredients (in each 30 mL)	Purpose
Acetaminophen 650 mg	Pain reliever/fever reducer
Dextromethorphan HBr 20 mg	Cough suppressant
Doxylamine succinate 12.5 mg	Antihistamine
Phenylephrine HCl 10 mg	Nasal decongestant

**Uses** temporarily relieves common cold/flu symptoms: ■ sinus congestion and pressure ■ nasal congestion ■ minor aches and pains ■ headache ■ runny nose and sneezing ■ sore throat ■ cough to help you sleep ■ fever ■ cough due to minor throat and bronchial irritation ■ reduces swelling of nasal passages ■ promotes nasal and/or sinus drainage ■ temporarily restores freer breathing through the nose

### Warnings

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take ■ more than 4,000 mg of acetaminophen in 24 hours ■ with other drugs containing acetaminophen ■ 3 or more alcoholic drinks every day while using this product  
**Allergy alert:** Acetaminophen may cause severe skin reactions. Symptoms may include: ■ skin reddening ■ blisters ■ rash  
If a skin reaction occurs, stop use and seek medical help right away.

PEEL BACK AT CORNER FOR MORE INFORMATION

: 76340 45 B1



### Drug Facts (continued)

**Sore throat warning:** If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

**Do not use** ■ with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist. ■ if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product. ■ if you have ever had an allergic reaction to this product or any of its ingredients  
Ask a doctor before use if you have

ESIVE AREA  
NO VARNISH • NO TYPE

### Drug Facts (continued)


When using this product ■ do not use more than directed ■ excitability may occur, especially in children ■ marked drowsiness may occur ■ avoid alcoholic drinks ■ be careful when driving a motor vehicle or operating machinery ■ alcohol, sedatives, and tranquilizers may increase drowsiness

**Stop use and ask a doctor if** ■ you get nervous, dizzy or sleepless ■ pain, nasal congestion, or cough gets worse or lasts more than 7 days ■ fever gets worse or lasts more than 3 days ■ redness or swelling is present ■ new symptoms occur ■ cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.

**If pregnant or breast-feeding,** ask a health professional before use.  
**Keep out of reach of children. Overdose warning:** In case of overdose, get medical help or contact a Poison Control Center right away (1-800-222-1222). Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

**Directions** ■ take only as directed - see Overdose warning ■ only use the dose con

ADHESIVE AREA  
NG • NO VARNISH • NO TYPE



## NIGHTTIME SEVERE COLD AND FLU

acetaminophen, dextromethorphan hbr, doxylamine succinate, phenylephrine hcl solution

### Product Information

<b>Product Type</b>	HUMAN OTC DRUG	<b>Item Code (Source)</b>	NDC:30142-763
<b>Route of Administration</b>	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
<b>ACETAMINOPHEN</b> (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	650 mg in 30 mL
<b>DEXTROMETHORPHAN HYDROBROMIDE</b> (UNII: 9D2RTI9KYH) (DEXTROMETHORPHAN - UNII:7355X3ROTS)	DEXTROMETHORPHAN HYDROBROMIDE	20 mg in 30 mL
<b>DOXYLAMINE SUCCINATE</b> (UNII: V9B19B5Y12) (DOXYLAMINE - UNII:95QB77JKPL)	DOXYLAMINE SUCCINATE	12.5 mg in 30 mL
<b>PHENYLEPHRINE HYDROCHLORIDE</b> (UNII: 04JA59TNSJ) (PHENYLEPHRINE - UNII:1WS297W6MV)	PHENYLEPHRINE HYDROCHLORIDE	10 mg in 30 mL

### Inactive Ingredients

Ingredient Name	Strength
<b>ANHYDROUS CITRIC ACID</b> (UNII: XF417D3PSL)	
<b>EDETATE DISODIUM</b> (UNII: 7FLD91C86K)	
<b>FD&amp;C BLUE NO. 1</b> (UNII: H3R47K3TBD)	
<b>FD&amp;C RED NO. 40</b> (UNII: WZB9127XOA)	
<b>GLYCERIN</b> (UNII: PDC6A3C0OX)	
<b>PROPYLENE GLYCOL</b> (UNII: 6DC9Q167V3)	
<b>WATER</b> (UNII: 059QF0KOOR)	
<b>SACCHARIN SODIUM</b> (UNII: SB8ZUX40TY)	
<b>SODIUM BENZOATE</b> (UNII: OJ245FE5EU)	
<b>SODIUM CHLORIDE</b> (UNII: 451W47IQ8X)	

<b>SODIUM CITRATE, UNSPECIFIED FORM</b> (UNII: 1Q73Q2JULR)	
<b>SORBITOL</b> (UNII: 506T60A25R)	
<b>SUCRALOSE</b> (UNII: 96K6UQ3ZD4)	
<b>XANTHAN GUM</b> (UNII: TTV12P4NEE)	

### Product Characteristics

<b>Color</b>	RED (clear, dark)	<b>Score</b>	
<b>Shape</b>		<b>Size</b>	
<b>Flavor</b>	BERRY	<b>Imprint Code</b>	
<b>Contains</b>			

### Packaging

#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:30142-763-40	354 mL in 1 BOTTLE; Type 0: Not a Combination Product	08/13/2014	

### Marketing Information

Marketing Category	Application Number or Monograph Citation	Marketing Start Date	Marketing End Date
OTC Monograph Drug	M012	08/13/2014	

**Labeler** - Kroger Company (006999528)

Revised: 10/2024

Kroger Company